


Baterun Kunsah

Bioactivity of Phytochemicals in Curcuma Domestica on Lymphocyte Cell Count in Mice (Mus Musculus)

 Quick Submit

 Quick Submit

 Universitas Muhammadiyah Surabaya

Document Details

Submission ID

trn:oid::1:3196266430

Submission Date

Mar 27, 2025, 8:22 PM GMT+7

Download Date

Mar 27, 2025, 8:24 PM GMT+7

File Name

9_Bioactivity_of_Phytochemicals_in_Curcuma.pdf

File Size

252.2 KB

8 Pages

3,235 Words

16,947 Characters

10% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- ▶ Bibliography
- ▶ Quoted Text
- ▶ Small Matches (less than 8 words)




Exclusions

- ▶ 5 Excluded Sources

Match Groups

-  **24 Not Cited or Quoted 10%**
Matches with neither in-text citation nor quotation marks
-  **1 Missing Quotations 0%**
Matches that are still very similar to source material
-  **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 10%  Internet sources
- 0%  Publications
- 0%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- **24 Not Cited or Quoted 10%**
Matches with neither in-text citation nor quotation marks
- **1 Missing Quotations 0%**
Matches that are still very similar to source material
- **0 Missing Citation 0%**
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**
Matches with in-text citation present, but no quotation marks

Top Sources

- 10% Internet sources
- 0% Publications
- 0% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	pusbangdik.poltekkesdepkes-sby.ac.id	2%
2	Internet	garuda.kemdikbud.go.id	1%
3	Internet	e-journal.stiekusumanegara.ac.id	1%
4	Internet	www.britishjournalofnursing.com	<1%
5	Internet	perpustakaan.poltekkes-malang.ac.id	<1%
6	Internet	innovareacademics.in	<1%
7	Internet	repository.unusa.ac.id	<1%
8	Internet	valleyinternational.net	<1%
9	Internet	siladikti.hangtuah.ac.id	<1%
10	Internet	dspace.uii.ac.id	<1%

11 Internet

ojs.unimal.ac.id <1%

12 Internet

ejournal.ung.ac.id <1%

13 Internet

jte.uho.ac.id <1%

Bioactivity of Phytochemicals in *Curcuma Domestica* on Lymphocyte Cell Count in Mice (*Mus Musculus*)

Nastiti Kartikorini¹(✉), Rahma Widyastuti², Diah Ariana¹, Baterun Kunsah¹, and Florencia Dita Permata¹

- ¹ D3 Medical Laboratory Technology Study Program, Faculty of Health Sciences, Universitas Muhammadiyah Surabaya, Surabaya, Indonesia
nastitikartikorini4956@gmail.com
- ² Applied Bachelor of Medical Laboratory Technology Study Program, Faculty of Health Sciences, Universitas Muhammadiyah Surabaya, Surabaya, Indonesia

Abstract. The immune system or immunity is a system that can protect the human body against foreign objects that enter the body. People can maintain their immune system by consuming healthy and nutritious food, exercising regularly, maintaining personal health, taking supplements or immune-boosting vitamins, and consuming traditional herbs, such as juice from the phytochemical *Curcuma domestica* (turmeric). *Curcuma domestica* is a tropical plant, and it can find in many parts of Asia, one of which is Indonesia. The main components of the phytochemical *Curcuma domestica* are curcumin and essential oils. In contrast, the phytochemicals of curcumin and essential oils can boost the immune system because they are proven to increase lymphocyte cells. This study aimed to determine the bioactivity of phytochemicals in *Curcuma domestica* on the number of lymphocyte cells in mice (*Mus musculus*). This study was experimental research with the pre and post-test design. The sample used in this study was 32 mice divided into two groups, including control and treatment groups. Data of lymphocyte cells were analyzed using a manual method, namely peripheral blood smear. The data were analyzed using the SPSS (Statistical Program Social Science) application using the independent t-test. The results were $p = 0.000$ ($p < 0.05$), so H_0 was rejected, and H_a was accepted. The results obtained in the control group showed that the average number of lymphocyte cells before treatment was 42.5%, and after treatment was 47.5%. While in the treatment group, the average number of lymphocyte cells before being given the phytochemical extract of *Curcuma domestica* was 41.75%, and after being given the phytochemical extract of *Curcuma domestica* was 57.25%. The conclusion is that giving the phytochemical extract of *Curcuma domestica* affects the number of lymphocyte cells in mice (*Mus musculus*).

Keywords: Bioactivity of phytochemicals · *Curcuma domestica* (Turmeric) · lymphocyte cell count · Mice (*Mus musculus*)

1 Introduction

The immune system is a system that forms the ability in the body to fight bacteria, viruses, and other microbes so that the body avoids a disease. Immunity in the human body is very important. If the body's immunity decreases, the ability to protect our body also decreases. Thus, bacteria, viruses, and other microbes or pathogens can grow and multiply in the human body. Several diseases associated with a lack of immunity include TB, HIV, THYPUS, and FLU, and what is often discussed at this time is COVID-19 [1].

East Java became the second province with the largest TB cases in Indonesia, reaching 22,585 cases with 208 TB cases in children, and the city of Surabaya became the highest city with TB cases reaching up to 3,093 cases [2]. According to a report from the Ministry of Health, Surabaya was ranked first in the highest HIV and AIDS cases in East Java province, with a total of 7,000 cases [3]. In the case of COVID-19 in Surabaya, the total of Patient Under Investigation (ODP) was 1,398 people, the total of Patient Under Surveillance (PDP) was 523 people, and the total of confirmed cases was 208 people [4].

In healing the above diseases, strong immunity is needed to avoid or recover from the disease. One of the cells that play an important role in the immune system or commonly called immunity, is a lymphocyte cell. B lymphocytes are derived from bone marrow cells and are important in forming humoral antibodies in the blood. In contrast, T lymphocyte cells come from the thymus (thymus gland) and have an important role in forming antibodies cellularly. The normal value of lymphocyte cells in humans is 25–40% of the total number of leukocytes [5]. There are various ways to strengthen the immune system, including taking medicines that contain vitamin C, herbal ingredients, and traditional herbs. Plants that can be consumed and can be made into herbs to strengthen the immune system include ginger and turmeric [6].

Curcuma domestica (Turmeric) is a tropical plant, and it can be found in many parts of Asia, one of which is Indonesia. *Curcuma domestica* plants are easier to find or obtain, easy to plant, relatively inexpensive, and have many benefits. It is often considered alternative medicine and used as a spice, dye, and herbal medicine [7]. It also has a variety of shapes, including long, short, thick, and straight and some are curved. The main components of *Curcuma domestica* are the phytochemicals of curcumin and essential oils. The phytochemical content of curcumin in turmeric has many medicinal properties, such as antioxidant, anti-inflammatory, and antibacterial, and it can maintain our immune system [8].

Maintaining immunity is very important for humans. There are many ways to maintain the body's immunity to stay awake, one of which is by consuming traditional herbs, such as juice from *Curcuma domestica*, containing phytochemicals. The phytochemical of *Curcuma domestica* can boost the immune system and increase the number of leukocytes. The phytochemical content of curcumin in *Curcuma domestica* can increase the number of leukocytes, one of which is lymphocyte cells, because it functions as an antigen against disease [9].

The phytochemical curcumin in *Curcuma domestica* is naturally yellow. Curcumin belongs to a group of polyphenolic compounds that can cause protein denaturation and damage cell membranes. Phenolic compounds contained in the phytochemical of

Curcuma domestica can damage and penetrate bacterial cell walls and then precipitate microbial cell proteins [10].

5 Population and Research Sample

The population in this study was a group of mice (*Mus musculus*) obtained from the Central Agency of Veterinary Farma (PUSVETMA), A. Yani Street, 68–70 Surabaya. Meanwhile, the sample from this study was 32 mice (*Mus musculus*) divided into two groups, each consisting of 16 mice (*Mus musculus*) in healthy condition with shining eyes, not dull fur, active, good appetite, and approximately 20–30 g.

Research Variables and Operational Definitions of Variables

There are two types of variables used for this research, namely:

1. Independent variable: Administration of Phytochemical extract of *Curcuma domestica* (Turmeric)
2. Dependent variable: Total lymphocyte cell

The operational definitions of variables are as follows:

1. Provision of Phytochemical Extract of *Curcuma domestica*
Giving the phytochemical extract of *Curcuma domestica* to mice (*Mus musculus*) as much as 1 ml every one time a day for 14 days using the probe through the mouth of mice (*Mus musculus*). Meanwhile, mice (*Mus musculus*) not given turmeric juice (*Curcuma domestica*) were replaced with aquades.
2. The number of lymphocyte cells was calculated using the manual method, namely making a peripheral blood smear that was read under a microscope. The number of lymphocyte cells was obtained from the calculation of the number of lymphocyte cells in the blood smear of mice (*Mus musculus*). The normal value of lymphocyte cells in humans is around 20–50% [11].

2 Methods of Data Collection and Data Analysis

The method used to examine lymphocyte cells is a blood smear. The blood smear method is intended to quantitatively determine the number of lymphocyte cells in the body. An easy way to do this is by using a microscope. Meanwhile, the data analysis used was the free T-test to determine the difference in the number of mice's lymphocytes before and after administrating the phytochemical extract of *Curcuma domestica*.

3 Results

The results of the study entitled “Bioactivity of Phytochemical in *Curcuma domestica* (Turmeric) on the Lymphocyte Cells Count in Mice (*Mus musculus*) used 32 mice (*Mus musculus*), which were carried out with two treatments and 16 repetitions. The examination was carried out by looking at the number of lymphocyte cells in mice (*Mus musculus*) before and after administration of the phytochemical extract of *Curcuma domestica* between the control group (TP) and the treatment group (P) to obtain the data in Table 1 as follows:

Table 1. Research Results of Phytochemical of Curcuma domestica on Lymphocyte Cell Count in Mice (*Mus musculus*)

Control Group				Treatment Group			
Sample Code	Lymphocyte Cell Number (%)		Difference (%)	Sample Code	Lymphocyte Cell Number (%)		Difference (%)
	Before	After			Before	After	
K1	44	52	8	P1	25	44	19
K2	40	45	5	P2	41	60	19
K3	40	46	6	P3	47	65	18
K4	44	46	2	P4	46	58	12
K5	42	50	8	P5	48	65	17
K6	41	42	1	P6	50	63	13
K7	41	48	7	P7	49	65	16
K8	40	49	9	P8	30	49	19
K9	49	50	1	P9	30	45	15
K10	40	48	8	P10	43	55	12
K11	43	46	3	P11	47	60	13
K12	44	47	3	P12	42	55	13
K13	44	50	6	P13	42	58	16
K14	44	52	8	P14	43	58	15
K15	42	44	2	P15	39	56	17
K16	42	45	3	P16	46	60	14
Quantity	680	760	80	Quantity	668	916	248
Average	42,5	47,5	5	Average	41,75	57,25	15,5
SD	2,37	2,89	2,83	SD	7,39	6,55	2,50

Information: K: Control; P: Treatment

The average increase in the number of lymphocyte cells in the Table 1 shows a difference between the average of the control group (TP) and the average of the treatment group (P). In contrast, the average difference between the treatment groups was higher than the average difference between the control group. In the control group, the average number of lymphocyte cells before treatment was 42.5%, and after treatment was 47.5%. While in the treatment group, the average number of lymphocyte cells before being given the phytochemical extract of *Curcuma domestica* was 41.75%, and after being given the phytochemical extract of *curcuma domestica* was 57.25%. The following is a graph of the average difference in the number of lymphocyte cells in the control and treatment groups (Fig. 1).




Fig. 1. Bioactivity of Phytochemical in *Curcuma domestica* (Turmeric) on Lymphocyte Cell Count in Mice (*Mus musculus*)

3.1 Data Analysis

The research data contained in Table 1 were processed using the SPSS (Statistical Program Social Science) application. then, the normality test was carried out using the Kolmogorov-Smirnova test, and the data were normally distributed with $p > 0.05$ (attached in the attachment). Furthermore, the data was tested using the homogeneity test, and the data was declared homogeneous with a result of $p > 0.05$ (attached in the appendix). After the data was declared normal and homogeneity, the independent T-test (Independent Sample T-test) was carried out. A free T-test was conducted to determine the effect of increasing the number of lymphocyte cells in mice (*Mus musculus*), which were given the phytochemical juice of *curcuma domestica* and without the phytochemical extract of *Curcuma domestica*. based on the tests carried out, it can be seen that the significant value was $(p) = 0.000 (<0.05)$, so H_0 was rejected, and H_a was accepted, so there was a significant effect on the number of lymphocytes between mice (*Mus musculus*) given turmeric juice (*Curcuma domestica*) with mice (*Mus musculus*) that were not squeezed with turmeric (*Curcuma domestica*).

4 Discussion

Based on the results of research on the phytochemical bioactivity of *curcuma domestica* on the number of lymphocyte cells in mice (*Mus musculus*). This research shows that the effect of turmeric (*Curcuma domestica*) juice has been shown to increase the number of lymphocyte cells in mice (*Mus musculus*). This can be seen from the results of the difference in the average calculation, which shows that there is a difference in the amount between the control group without turmeric juice (*Curcuma domestica*) with an average difference of 5% and the treatment group given turmeric juice (*Curcuma domestica*) with an average difference 15.5%. The t-test results show that the significant result is 0.000 (<0.05), so H_0 is rejected, and H_a is accepted.

12 Administration of phytochemical extract of *Curcuma domestica* increases the number of lymphocyte cells in mice (*Mus musculus*). *Curcuma domestica* contains an active compound, namely curcumin, where curcumin has many benefits, including as an antibacterial, antioxidant, anti-inflammatory, and very good antiviral used to increase our immune system to stay healthy [6, 12]. The phytochemical content of *Curcuma domestica* functions as an immune stimulant and as an antigen against disease [9, 13]. The phytochemical *Curcuma domestica* has anti-inflammatory properties that can inhibit the production of prostaglandins in the cyclooxygenase cycle through the process of inhibiting the activity of the cyclooxygenase enzyme and suppressing leukotriene biosynthesis by inhibiting lipoxygenase and binding free radicals that can cause the inflammatory process [14, 15].

7 In addition to the curcumin content, according to Yuliati et al. [16], essential oils can also be used as antibacterial agents. This is because essential oils contain hydroxyl and carbonyl functional groups, including phenol derivatives. These hydroxyl and carbonyl functional groups can inhibit the growth of bacteria by damaging the process of cell membrane formation so that the cell membrane is not formed perfectly.

6 *Curcuma domestica* is a plant that is very easy to find or people often encounter in our area, and the price is relatively low. This *Curcuma domestica* plant is often used as a spice or kitchen spice, as a coloring agent, can also be made herbal medicine, and is often considered as an alternative medicine. This *Curcuma domestica* plant has many benefits, including increasing our body's resistance to avoid disease. Some researchers have also tested the benefits of *Curcuma domestica* and found many benefits to help our body's health needs, especially during a pandemic like this.

6 Based on the above results, it can be proven by the results of the number of lymphocyte cells in mice (*Mus musculus*) after being given the phytochemical juice of *Curcuma domestica* (turmeric) experienced a significant increase. Due to the content of curcumin and essential oils in *Curcuma domestica*, it functions as an immune system. Increasing the body's resistance can help us avoid contracting a disease and avoid bacteria, viruses, and other microbes or pathogens. So that the phytochemical extract of *Curcuma domestica* can be used as a choice of herbal medicine to increase human endurance was 57.25%.

Suggestion

1. For Further Researchers.

For further research, it can be done by adding other phytochemicals, such as ginger, and java ginger, to increase body resistance and maximize the number of lymphocyte cells.

2. For Educational Institutions

This research can be used to add broader knowledge and as input for learning media in developing science in the field of basic chemistry for health and hematology in educational institutions, especially medical laboratory technology.

3. For Society

People can consume turmeric juice to increase endurance because it contains curcumin and essential oils, so it can be a substitute for chemical drugs to increase endurance.

References

1. A. A. Hidayat, Sopyan., Syahputa, "Sistem imun tubuh pada manusia," *Vis. Herit. J. Kreasi Seni dan Budaya*, vol. 2, no. 03, pp. 144–149, 2020.
2. Kementerian Kesehatan RI, "Profil Kesehatan Indonesia Tahun 2017," Kementerian Kesehatan Republik Indonesia, Jakarta, 2017.
3. Huda, Jesica Miftakhul., Prasetyo, Iwan Joko., Fitriyah, Irmia, "Komunikasi Interpersonal Antar ODHA untuk Menumbuhkan Motivasi Kembali Hidup Normal di Yayasan Mahameru Surabaya," *J. Komun. Prof.*, vol. 3, no. 1, pp. 12–22, 2019.
4. S. Albana, Abduh Sayid., Azhari, "Prediksi Penyebaran COVID-19 Kota Surabaya dengan Simulasi Monte Carlo," *J. Adv. Inf. Ind. Technol.*, vol. 2, no. 1, pp. 36–42, 2020.
5. Tiara, Dhea., Tiho, Murniati., Mewo, Yanti M., "Gambaran kadar limfosit pada pekerja bangunan," *J. e-Biomedik*, vol. 4, no. 2, pp. 2–5, 2016.
6. S. S. Kusumo, Adristy Ratna., Wiyoga, Farrel Yumna., Perdana, Haekal Putra., Khairunnisa, Izzatidiva., Suhandi, Raihan Ibadurrohman., Prastika, "Traditional Indonesian Jamu: Natural Way to Boost Immune System During Pandemic," *J. Public Serv.*, vol. 4, no. 2, pp. 1–7, 2020.
7. Y. Shan, Chu Yuan., Iskandar, "Studi Kandungan Kimia dan Aktivitas Farmakologi Tanaman Kunyit (*Curcuma Longa L.*)," *Farmaka*, vol. 16, no. 2, pp. 547–555, 2018.
8. Azis, Abdul, "KUNYIT (*Curcuma domestica Val*) SEBAGAI OBAT ANTIPIRETIK," *J. Ilmu Kedokt. dan Kesehat.*, vol. 6, no. April, pp. 116–120, 2019.
9. U. N. Susantie, Darna., Manurung., "ADDITION OF BAKER'S YEAST (*Saccaromyces cereviceae*) AND TURMERIC (*Curcumae domesticae Val*) TO FISH FEED FOR INCREASE GROWTH AND IMMUNITY OF FISH IN KAWIO ISLAND REGENCY OF SANGIHE," *J. Ilmu Tatengkorang*, vol. 3, pp. 66–71, 2019.
10. Nobiola, Rezica Kanza., Triwahyuni, Tusy., Triswanti, Nia., Warganegara, Efrida, "Uji Sensitivitas Kunyit Kuning dan Kunyit Putih Terhadap Bakteri Pencemar Susu," *Arter. J. Ilmu Kesehat.*, vol. 1, no. 4, pp. 263–269, 2020.
11. Unawekla, Julyan V., Moeis, Emma Sy., Langi, Yuanita A, "Hubungan antara Status Gizi dan Sistem Imun Seluler pada Subyek Penyakit Ginjal Kronik Stadium V Hemodialisis di Instalasi Tindakan Hemodialisis RSUP Prof Dr. R. D. Kandou Manado," *J. e-Clinic*, vol. 6, no. 1, pp. 16–21, 2018.
12. D. D. Cahyani, Y. Armiyanti, C. Komariyah, B. Hermansyah, and Y. Nurdian, "Description of Leukocytes Differential Count in Coffee Plantation Workers Silo Subdistrict that Infected by Soil-transmitted Helminths," *J. Agromedicine Med. Sci. Vol 6 No 1 (2020)DO* - <https://doi.org/10.19184/ams.v6i1.9620>, Jan. 2020.
13. Wiyanti, Wiwik Rosi., Endrawati, Susi, "The Influence of Tonicum Effect Test of Ethanol Extract Temu Giring Rhizome (*Curcuma Heyneana Val*) to Mice.," *IJMS (Indonesian J. Med. Sci.*, vol. 4, no. 1, pp. 25–32, 2017.
14. Wulandari, Ratna., Puspitasari, Puspitasari, "Pengaruh Infusa Rimpang Temu Putih (*Curcuma zedoaria* (Berg.) Roscoe) Terhadap Jumlah Leukosit Dan Differential Counting (Diffcount) Pada Kesembuhan Luka Laparatomi Pasca Bedah," *J. Med. Lab. Sci. Technol.*, vol. 2, no. 1, pp. 22–27, 2019.
15. A. A. Rahmah, "The Effectiveness of Turmeric (*Curcuma domestica*) in Decreasing The Risk of Atherosclerosis," *Prev. J. Kesehat. Masy.*, vol. 10, no. 2 SE-Articles, Aug. 2020.
16. Yuliati et al, "Uji Efektivitas Ekstrak Kunyit sebagai Antibakteri dalam Pertumbuhan *Bacillus sp* dan *Shigella Dysentriae* Secara In Vitro," *J. Profesi Med.*, vol. 10, no. 1, pp. 26–32, 2016.

Open Access This chapter is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.

