

Gastric mucosal status

by Muhammad Miftahussurur

Submission date: 24-Oct-2019 04:56PM (UTC+0800)

Submission ID: 1199377950

File name: Gastric_mucosal_status_in_populations.pdf (2.49M)

Word count: 9240

Character count: 46647

RESEARCH ARTICLE

Gastric mucosal status in populations with a low prevalence of *Helicobacter pylori* in Indonesia

Muhammad Miftahussurur^{1,2,3,4}, Iswan Abbas Nusi³, Fardah Akil⁵, Ari Fahrial Syam⁶, I. Dewa Nyoman Wibawa⁷, Yudith Annisa Ayu Rezkitha⁴, Umami Maimunah³, Phawinee Subsomwong¹, Muhammad Luthfi Parewangi⁶, I. Ketut Mariadi⁷, Pangestu Adi⁴, Tomohisa Uchida⁸, Herry Purbayu³, Titong Sugihartono³, Langgeng Agung Waskito^{1,4}, Hanik Badriyah Hidayati⁴, Maria Inge Lusida⁴, Yoshio Yamaoka^{1,2*}

1 Department of Environmental and Preventive Medicine, Oita University Faculty of Medicine, Yufu, Japan, **2** Department of Medicine, Gastroenterology and Hepatology Section, Baylor College of Medicine, Houston, Texas, United States of America, **3** Division of Gastroentero-hepatology, Department of Internal Medicine, Faculty of Medicine-Dr. Soetomo Teaching Hospital, Universitas Airlangga, Surabaya, Indonesia, **4** Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia, **5** Center of Gastroentero-hepatology, Department of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia, **6** Division of Gastroenterology, Department of Internal Medicine, Faculty of Medicine, University of Indonesia, Jakarta, Indonesia, **7** Division of Gastroentero-hepatology, Department of Internal Medicine, Faculty of Medicine University of Udayana, Denpasar, Indonesia, **8** Department of Molecular Pathology, Oita University Faculty of Medicine, Hasama-machi, Yufu-City, Oita, Japan

* yyamaoka@oita-u.ac.jp



OPEN ACCESS

Citation: Miftahussurur M, Nusi IA, Akil F, Syam AF, Wibawa IDN, Rezkitha YAA, et al. (2017) Gastric mucosal status in populations with a low prevalence of *Helicobacter pylori* in Indonesia. PLoS ONE 12(5): e0176203. <https://doi.org/10.1371/journal.pone.0176203>

Editor: Niyaz Ahmed, International Centre for Diarrhoeal Disease Research Bangladesh (icddr), BANGLADESH

Received: December 20, 2016

Accepted: April 6, 2017

Published: May 2, 2017

Copyright: © 2017 Miftahussurur et al. This is an open access article distributed under the terms of [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files (S1 Table).

Funding: This work was supported in part by grants from the National Institutes of Health (R01NS082813) and Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan (25293126, 26640114, and 15H02657) (YY). Support was also provided by the Japan Society for

Abstract

In Indonesia, endoscopy services are limited and studies about gastric mucosal status by using pepsinogens (PGs) are rare. We measured PG levels, and calculated the best cutoff and predictive values for discriminating gastric mucosal status among ethnic groups in Indonesia. We collected gastric biopsy specimens and sera from 233 patients with dyspepsia living in three Indonesian islands. When ≥ 5.5 U/mL was used as the best cutoff value of *Helicobacter pylori* antibody titer, 8.6% (20 of 233) were positive for *H. pylori* infection. PG I and II levels were higher among smokers, and PG I was higher in alcohol drinkers than in their counterparts. PG II level was significantly higher, whereas PG I/II ratios were lower in *H. pylori*-positive than in *H. pylori*-negative patients. PG I/II ratios showed a significant inverse correlation with the inflammation and atrophy scores of the antrum. The best cutoff values of PG I/II were 4.05 and 3.55 for discriminating chronic and atrophic gastritis, respectively. PG I, PG II, and PG I/II ratios were significantly lower in subjects from Bangli than in those from Makassar and Surabaya, and concordant with the ABC group distribution; however, group D (*H. pylori* negative/PG positive) was the lowest in subjects from Bangli. In conclusion, validation of indirect methods is necessary before their application. We confirmed that serum PG level is a useful biomarker determining chronic gastritis, but a modest sensitivity for atrophic gastritis in Indonesia. The ABC method should be used with caution in areas with a low prevalence of *H. pylori*.

the Promotion of Science (JSPS) Institutional Program for Young Researcher Overseas Visits (YY), the Strategic Funds for the Promotion of Science and Technology from the Japan Science and Technology Agency (JST and YY). MM was a previous PhD student, and PS and LAW are PhD students supported by the Japanese government (MEXT) scholarship program for 2012, 2013, and 2015, respectively.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: PGs, pepsinogens; OLGA, Operative Link for Gastritis Assessment; ELISA, enzyme-linked immunosorbent assay; AUC, area under curve; ROC, receiver-operating characteristic; NSAID, nonsteroidal anti-inflammatory drug; PPV, positive predictive value; NPV, negative predictive value; OR, odds ratio; CI, confidence interval.

Introduction

Helicobacter pylori has a unique capacity to persistently colonize the extremely acidic environment of the stomach and cause progressive gastric mucosal inflammation. Long-term infection induces a multistep histological cascade, from chronic non-atrophic gastritis that progresses to chronic atrophic gastritis, intestinal metaplasia, and adenocarcinoma [1]. Currently, chronic atrophic gastritis characterized by chronic inflammation with loss of gastric glandular cells is an established precursor lesion to gastric adenocarcinoma [2]. Although gastric mucosal biopsy is the reference method for determining the grade and topographical distribution of gastritis [3], this method is uncomfortable and expensive for patients. In contrast, serology is a cheaper, acceptable, and easily repeated method. Recent reports confirmed that serum pepsinogens (PGs) are a valuable biomarker of the gastric mucosal status, including inflammation, atrophic gastritis, and gastric cancer [4], even before the discovery of *H. pylori* [5].

Although the most part of PGs are secreted by the gastric cells and, in low levels, are permeated into serum [6]. PG I and PG II, the two main types of PGs are produced in different regions of the stomach [7]. Serum PG I, which is secreted purely on the fundus, decreases progressively, whereas PG II level decreases less markedly or remain stable owing to the additional production in non-gastric glands such as the duodenal bulb; therefore, PG I/II ratios serve as a gastric mucosal biomarker [7] and can be applied to gastric cancer screening [4, 8, 9]. The cut-off points of ≤ 70 $\mu\text{g/L}$ and ≤ 3 for PG I level and PG I/II ratio, respectively, were used in Japan for identifying the risk of gastric cancer [10]. However, the discriminative baseline PG levels may not be applicable to other countries and therefore should be recalculated, because several factors including geographic area, race, age, sex, smoking and drinking habits, and *H. pylori* infection also contribute to the levels [11–13]. In addition, there is evidence that PG II level increases and the PG I/II ratio decreases with the grade of inflammation [14], providing an opportunity for the application PG levels in patients with chronic gastritis who also frequent in countries with a lower prevalence of gastric cancer than that in Japan [10].

As we explained in detail previously [15–19], Indonesia is a multiethnic nation in Southeast Asia with a relatively low risk of gastric cancer (age-standardized incidence rate [ASR]: 2.8 in 100,000; GLOBOCAN2012, <http://globocan.iarc.fr/>) with the overall prevalence of *H. pylori* infection in the five largest islands of Indonesia was 22.1% [17]. The predominant ethnic, Javanese, has a low prevalence of *H. pylori* infection (2.4%) [16, 17], however, several ethnic groups have a much higher risk of *H. pylori* infection [17]. According to Indonesia as the fourth most populous country worldwide with a high prevalence of *H. pylori* antibiotic resistance [19, 20], and the strains harbor more virulent *H. pylori* genotypes [18], consequently, *H. pylori*-associated diseases are an important problem in Indonesia. Moreover, although dyspepsia was sixth and fifth of the 10 most prevalent outpatient and inpatient diseases in Indonesia, respectively [21], hospitals that provide gastrointestinal endoscopy services in the country are very limited. Therefore, noninvasive methods such as measurement of PG values are the best choice for determining the gastric mucosal status in Indonesia, especially in remote areas.

To our knowledge, studies examining the gastric mucosal status in Indonesia by using PG levels are rare [22, 23]. Two previous studies determined the accuracy of PG levels in a single city that was predominantly inhabited by the Javanese people: Purwokerto [22] and Jakarta [23]. These studies also did not compare the PG data with histological grades (e.g., updated Sydney system) [24]. The study performed in Purwokerto reported that the specificity of PG I level to predict atrophic chronic gastritis was only 50%, with a sensitivity of 70% [22]. In contrast, the study in Jakarta suggested a low sensitivity (43%) of PG I/II as the biomarker of *H. pylori* chronic gastritis, with a specificity of 83% [23]. In the current study, we measured the PG levels of three ethnic groups from different islands. We also calculated the best cutoff and

predictive values for discriminating chronic and atrophic gastritis based on PG levels among ethnic groups in Indonesia.

Materials and methods

Study population

We performed a prospective study between January and August 2015 in Surabaya, Java Island; Makassar, Sulawesi Island; and Bangli, Bali Island. We excluded subjects with a history of *H. pylori* eradication therapy and partial/total gastrectomy, nonfasted subjects, and those with contraindication for upper endoscopy. Experienced endoscopists acquired two gastric biopsy specimens during each endoscopy procedure: one from the lesser curvature of the antrum approximately 3 cm from the pyloric ring, and another from the greater curvature of the corpus, which were used for histological examination. Fasting serum was collected on the day of endoscopy and then stored at -20°C. Socio-demographic data including body mass index, smoking and drinking habits, and use of nonsteroidal anti-inflammatory drugs (NSAIDs) were collected during the interview. Written informed consent was obtained from all participants, and the study protocol was approved by the ethics committees of Dr. Soetomo Teaching Hospital (Surabaya, Indonesia), Dr. Wahidin Sudirohusodo Teaching Hospital (Makassar, Indonesia), and Oita University Faculty of Medicine (Yufu, Japan).

Determination of *H. pylori* serology and PG levels

The separated sera were used for the measurement of the *H. pylori* antibody titers and PG levels. The anti-*H. pylori* IgG levels were quantified by using an ELISA kit (Eiken, Co. Ltd., Tokyo, Japan), and the PG I and II levels were measured by using PG ELISA (Eiken), according to the manufacturer's instructions. Subjects with serum *H. pylori* antibody titers ≥ 10 IU/mL were classified as *H. pylori* positive per the manufacturer's instructions. Those with PG I level ≤ 70 ng/mL and PG I/II ratio ≤ 3.0 were classified as PG positive according to the Japanese guidelines [4]. According to the ABC method, we categorized the subjects into four groups: *H. pylori* negative/PG negative (group A), *H. pylori* positive/PG negative (group B), *H. pylori* positive/PG positive (group C), and *H. pylori* negative/PG positive (group D) [4].

Histology and immunohistochemistry

All biopsy materials for histological testing were fixed in 10% buffered formalin and embedded in paraffin. Serial sections were stained with hematoxylin and eosin and May-Giemsa stain. The degree of inflammation, atrophy, and bacterial density were classified into four grades according to the updated Sydney system: 0, normal; 1, mild; 2, moderate; and 3, marked [24]. Samples with bacterial loads \geq grade 1 were considered positive for *H. pylori*.

To increase the accuracy for detecting *H. pylori*, we performed immunohistochemical confirmation, as previously described [25]. Briefly, after antigen retrieval and inactivation of endogenous peroxidase activity, tissue sections were incubated with anti- α -*H. pylori* antibody (DAKO, Glostrup, Denmark) overnight at 4°C. After washing, the sections were incubated with biotinylated goat anti-rabbit IgG (Nichirei Co., Tokyo, Japan), followed by incubation with an avidin-conjugated horseradish peroxidase solution (Vectastain Elite ABC Kit; Vector Laboratories Inc., Burlingame, CA, USA). Peroxidase activity was detected by using an H₂O₂/diaminobenzidine substrate solution. To minimize potential bias, the same experienced pathologist (TU) who also performed experiments for Myanmar, Vietnam, Bhutan, Dominican Republic, and Indonesia [16, 26–30] evaluated all the specimens in this study.

Data analyses

Discrete variables were tested by using the chi-square test; continuous variables were tested with the Mann-Whitney *U* and Kruskal–Wallis test. The Spearman rank coefficients (*r*) were determined to evaluate the association between PG levels and gastric mucosal inflammation and atrophy. A multivariate logistic regression model was used to calculate the ORs of the clinical outcomes by including age, sex, NSAID use, alcohol drinking, smoking, *H. pylori* infection, and gastritis type. All determinants with $P < 0.10$ were entered together in the full model logistic regression, and the model was reduced by excluding variables with $P > 0.10$. The OR and 95% confidence interval (CI) were used to estimate the risk. $P < 0.05$ was accepted as statistically significant. Receiver-operating characteristic (ROC) curves were used to calculate the best cutoff, including the area under curve (AUC) and predictive values for discriminating chronic and atrophic gastritis. The SPSS statistical software package version 18.0 (SPSS Inc., Chicago, IL, USA) was used for all statistical analyses.

Results


H. pylori infection rate based on serology

We recruited 233 patients with dyspepsia comprising 110 women and 123 men with a mean age of 46.1 ± 12.3 years (range, 14–82 years), including 99 patients from Surabaya, 75 from Makassar, and 59 from Bangli (S1 Table). There were 19 patients with peptic ulcers. On the basis of histology confirmed by immunohistochemistry, 15 patients (6.4%) were positive for *H. pylori* infection.

By using the cutoff value from the manufacturer's instructions (positive if ≥ 10 U/mL), we found the sensitivity and specificity of the ELISA kit for *H. pylori* infection to be 66.7% and 97.2%, respectively, compared with histology confirmed by immunohistochemistry as a gold standard. As the low sensitivity was due to a large number of false negatives (5 of 15, 33.3%), we determined the best cutoff values of *H. pylori* antibody titers in ELISA kit with an ROC curve. We found that the cutoff of ≥ 5.5 U/mL was the best value to determine *H. pylori* positivity (sensitivity, specificity, positive predictive value [PPV], and negative predictive value [NPV]: 86.7%, 96.8%, 65.0%, and 99.1%, respectively; Fig 1), with an AUC of 0.913 (95% CI, 0.811–1.000). When we used the new cutoff value of ≥ 5.5 U/mL, 8.6% (20 of 233) were positive for *H. pylori* infection (S1 Table). The *H. pylori* infection rate by age group was 5.6% (1 of 18), 5.2% (3 of 58), 10.3% (6 of 58), 7.5% (5 of 67), and 15.6% (5 of 32) for patients aged ≤ 29 , 30–39, 40–49, 50–59, and ≥ 60 years, respectively. Patients with peptic ulcer had a higher prevalence of *H. pylori* infection than those without ulcer (5 of 19, 26.3% vs. 15 of 214, 7.0%; $P = 0.004$). According to location, Bangli (13.6%, 8 of 59) and Makassar (12.0%, 9 of 75) had a higher prevalence of *H. pylori* infection than Surabaya (3.0%, 3 of 99) ($P = 0.01$ and 0.02 , respectively).

Socio-demographic data and PG levels

Women had lower PG I, PG II, and PG I/II ratio than men ($P < 0.001$, $P < 0.001$, $P = 0.008$, respectively). PG I and PG II were increased with age ($r = 0.34$ and $r = 0.32$, $P < 0.001$, respectively), but not PG I/II ratio. There was no difference in the PG I and II levels and PG I/II ratios in patients with gastritis and those with peptic ulcer ($P = 0.10$, $P = 0.08$, $P = 0.85$, respectively). When we considered patients with any inflammation and atrophy in the antrum or corpus on histological examination as a symptomatic patient, PG II levels was higher and PG I/II ratio was lower among symptomatic than asymptomatic patients (15.4 ± 11.3 vs. 12.4 ± 9.6 , $P = 0.05$ and 6.0 ± 1.8 vs. 6.6 ± 2.0 , $P = 0.017$, respectively). There was no significant



Diagonal segments are produced by ties.

Fig 1. Receiver-operating characteristic (ROC) curve for determining the optimal cutoff of *Helicobacter pylori* antibody titers. The sensitivity delineated in X-axis (86.7%) and the Y-axis characterized value of 1-specificity (96.8%). PPV was a positive predictive value and NPV was a negative predictive value.

<https://doi.org/10.1371/journal.pone.0176203.g001>

association between the PG levels and NSAID use and body mass index ($P > 0.05$). Smoking was significantly higher in subjects from Makassar and Bangli than in those from Surabaya (28.0% [21 of 75], 22.0% [13 of 59] vs. 10.1% [10 of 99], $P = 0.002$ and $P = 0.04$). PG I and II levels were higher in smokers than in nonsmokers (114.9 ± 87.2 vs. 79.1 ± 67.6 , $P = 0.001$ and 18.9 ± 13.3 vs. 12.6 ± 9.4 , $P < 0.0001$, respectively). There was no significant difference in drinking habits in all populations ($P > 0.05$). PG I level was also higher in drinking than in nondrinking subjects (111.8 ± 91.8 vs. 84.0 ± 70.9 , $P = 0.043$).

H. pylori infection status and PG levels

PG I level tended to be higher in *H. pylori*-positive than in *H. pylori*-negative patients ($P = 0.056$). Additionally, PG II level was significantly higher in *H. pylori*-positive than in *H. pylori*-negative patients ($P < 0.0001$, Table 1), whereas the PG I/II ratios were significantly

Table 1. Demographics and *Helicobacter pylori* antibody status (mean ± SD).

Parameter	<i>H. pylori</i> positive	<i>H. pylori</i> negative	P-value
Male/female	10/10	100/113	
Age	49.75 ± 12.1	45.80 ± 12.3	0.14
Body mass index (kg/m ²)	21.82 ± 2.7	23.02 ± 3.8	0.18
Pepsinogen I (ng/mL)	113.73 ± 80.1	83.28 ± 71.8	0.056
Pepsinogen II (ng/mL)	24.09 ± 13.9	12.83 ± 9.6	<0.0001
Pepsinogen I/II ratio	4.70 ± 1.4	6.43 ± 1.9	<0.0001
Chronic gastritis (%)	20/20 (100.0)	69/213 (32.4)	<0.0001
Atrophic gastritis (%)	15/20 (75.0)	45/213 (21.2)	<0.0001

<https://doi.org/10.1371/journal.pone.0176203.t001>

lower in *H. pylori*-positive than in *H. pylori*-negative patients ($P < 0.0001$, Fig 2). When we considered patients with any inflammation and atrophy in the antrum or corpus on histological examination as positive for chronic and atrophic gastritis, respectively, the prevalence of chronic and atrophic gastritis was higher in *H. pylori*-positive than in *H. pylori*-negative patients (both $P < 0.0001$). Only *H. pylori* infection was an independent risk factor for the development of chronic gastritis in the corpus but not in the antrum (OR, 13.12; 95% CI, 2.3–73.8) after adjustment for age, sex, NSAID use, alcohol drinking, and smoking. After adjustment, *H. pylori* infection also was an independent risk factor for the development of atrophy in the antrum (OR, 9.87; 95% CI, 1.8–54.4).

PG levels and chronic gastritis

Inflammation in the antrum had a correlation with atrophy, similar to that in the corpus ($P < 0.001$, $r = 0.382$ and $P < 0.001$, $r = 0.384$, respectively). We examined the correlation between the severity of chronic gastritis and PG levels (Table 2). Overall, in patients with chronic gastritis, PG I level was almost stable in conjunction with the increase of PG II level and decrease of I/II ratio ($P < 0.05$). Although PG I level did not correlate with the inflammation scores ($P = 0.55$ in the antrum and $P = 0.31$ in the corpus), PG II level showed a weak correlation in the antrum and corpus (both $P = 0.01$, $r = 0.16$) and the association was disappeared in the multivariate analysis after adjusted with age, sex, smoking, drinking habits and positivity of *H. pylori* ($P = 0.21$). Additionally, PG I/II ratio showed a significant inverse correlation with the inflammation scores in the antrum and corpus ($P = 0.0001$, $r = -0.24$ and $P = 0.001$, $r = -0.22$, respectively). Moreover, when we analyzed only the *H. pylori*-positive patients, the PG I/II ratios showed a significant inverse correlation with the inflammation scores in the antrum ($P = 0.02$, $r = -0.61$) but not in the corpus ($P = 0.52$). In the multivariate analysis, the inverse association of PG I/II ratios and antral inflammation was still significant although adjusted with age, sex, smoking consumption and drinking habits (OR, 0.34; 95% CI, 0.21–0.56).

Next, we determined the best cutoff values for discriminating chronic gastritis. By using the ROC curve, PG II level was not found to be a good discriminatory marker for antral chronic gastritis (data not shown). For stages \geq II in the inflammation score, considered as positive for chronic gastritis, the best cutoff value of PG I/II was 5.65 (sensitivity, specificity, PPV, NPV, and accuracy: 83.3%, 63.7%, 16.1%, 97.9%, and 65.2%, respectively). The AUC was 0.832 (95% CI, 0.721–0.942). For stages \geq III in the inflammation score, the best cutoff value of PG I/II was 4.05 (sensitivity, specificity, PPV, NPV, and accuracy: 80.0%, 94.3%, 23.5%, and 99.5%, respectively). The overall accuracy was 94.0% with an AUC of 0.857 (95% CI, 0.664–1.000). The PG I best cutoff value was 47.5 (the AUC were 0.674 [95% CI, 0.585–0.762]) with sensitivity, specificity, PPV, NPV, and accuracy were 80.0%, 64.9%, 4.8%, 99.3%, 65.2%, respectively




Fig 2. Pepsinogens levels among *H. pylori* positive and negative patients. PG II level was significantly higher and PG I/II ratios were significantly lower in *H. pylori*-positive than in *H. pylori*-negative patients.

<https://doi.org/10.1371/journal.pone.0176203.g002>

for stages \geq III in the inflammation score. When we used a combination PG I and PG I/II for stages \geq III in the antral chronic gastritis, the outcomes practically the same as when utilizing just PG I/II ratio although had a higher PPV than only PG I/II ratio marker (sensitivity, specificity, PPV, NPV, and accuracy were 80.0%, 97.3%, 40.0%, 99.6%, and 97.0%, respectively).

PG levels and atrophic gastritis

According to the updated Sydney system, we determined that 74.3% of patients (173 of 233) had no mucosal atrophy in both the antrum and corpus, and these patients were categorized as the nonatrophic gastritis group. In addition, only 21.9% (51 of 233), 0.4% (1 of 233), and 3.4% (8 of 233) of patients had antral atrophy, corpus atrophy, and multifocal atrophy, respectively. There were differences in the PG II level and PG I/II ratio according to the pattern of atrophic

43

Table 2. Levels of pepsinogen (PG) I, PG II, and PG I/II in chronic and atrophic gastritis (mean ± SD).

	Grade	n	PG I	PG II	PG I/II
Chronic gastritis					
Antrum	0	147	85.3 ± 77.9	12.7 ± 9.7	6.5 ± 1.9
	1	68	89.3 ± 68.3	14.9 ± 11.4*	6.2 ± 2.0*
	2	13	89.7 ± 44.9	21.4 ± 13.2*	4.6 ± 1.4*
	3	5	46.7 ± 6.4	12.1 ± 2.8	4.1 ± 1.4
Corpus	0	187	84.5 ± 74.5	12.8 ± 9.7	6.4 ± 1.8
	1	42	90.2 ± 66.0	17.2 ± 12.7*	5.7 ± 2.4*
	2	4	106.5 ± 73.1	22.5 ± 13.5*	4.7 ± 2.0*
	3	0	NA	NA	NA
Atrophic gastritis					
Antrum	0	174	83.5 ± 72.0	12.8 ± 9.7	6.5 ± 2.0
	1	53	97.4 ± 78.5	17.2 ± 12.8*	5.7 ± 1.5*
	2	5	55.4 ± 10.8	13.6 ± 4.6	4.5 ± 1.8
	3	1	18	13.8	3.1
Corpus	0	224	84.6 ± 72.4	13.4 ± 10.2	6.3 ± 2.0
	1	8	112.9 ± 84.4	22.1 ± 13.8*	5.3 ± 1.9
	2	1	168.0	32.3*	5.2
	3	0	NA	NA	NA

* P < 0.05 vs. grade 0.

<https://doi.org/10.1371/journal.pone.0176203.t002>

gastritis (P = 0.009 and P = 0.023, respectively; Table 3). Compared with nonatrophic gastritis, antral atrophy had significantly lower PG/II ratio (P = 0.01). Moreover, PG I tended to have higher levels and PG I/II ratio tended to have lower levels in subjects with nonatrophic gastritis than those with multifocal atrophic gastritis (P = 0.053 and P = 0.057, respectively). PG II had a significantly lower level in subjects with nonatrophic gastritis than in those with multifocal atrophic gastritis (P = 0.002).

Although the PG I level did not correlate with the atrophy scores (P = 0.49 in the antrum and P = 0.11 in the corpus), the PG II level and PG I/II ratios showed a significant correlation in the antrum (P = 0.02, r = 0.16 and P = 0.002, r = -0.20, respectively; Table 2) and only PG II showed a significant correlation in the corpus (P = 0.02, r = 0.16). However, there was no correlation between the PG levels and atrophy scores among *H. pylori*-positive patients. This may be due to the low number of *H. pylori*-positive patients with atrophic gastritis. When we analyzed only *H. pylori*-negative patients, PG I and PG II levels showed a weak correlation with the atrophy scores in the corpus (P = 0.04, r = 0.14 and P = 0.03, r = 0.15, respectively).

Table 3. Pepsinogen level according to atrophic gastritis pattern.

Pepsinogens	No Atrophy	Antral atrophy	Corpus atrophy	Multifocal atrophy	P-value
N	173	51	1	8	
Pepsinogen I (ng/mL)	83.72 ± 72.1	87.42 ± 73.9	133.0	128.41 ± 81.3	0.24
Pepsinogen II (ng/mL)	12.79 ± 9.7	15.53 ± 11.7	17.9	25.30 ± 12.6*, **	0.009
Pepsinogen I/II ratio	6.51 ± 2.0	5.67 ± 1.5*	7.4	5.01 ± 1.9	0.023

Multifocal atrophy means the subjects had atrophy in both the antrum and corpus.

* P < 0.05 vs. no atrophy.

** P < 0.05 vs. antral atrophy.

<https://doi.org/10.1371/journal.pone.0176203.t003>

When we used the Japanese standards to define the PG-positive status (cutoff values: ≤ 70 ng/mL and ≤ 3.0 for PG I level and PG I/II ratio, respectively), we found that the overall prevalence of PG-positive subjects was very low (1.3%, 3 of 233). Furthermore, when we used the cutoff value only for PG I/II ratios ≤ 3.0 , the sensitivity, specificity, PPV, NPV, and accuracy were 5.1%, 99.4%, 75.0%, 75.6%, and 75.5%, respectively, for antral atrophic scores ≥ 1 . In cases with an antral atrophic score ≥ 2 , the values were 16.7%, 98.7%, 25.0%, 97.8%, and 96.6%, respectively.

Owing to the low sensitivity of the Japanese standard, we determined the best cutoff value of PG I/II ratio with an ROC curve. For stages $\geq I$ in the antral atrophic score, considered as atrophy positive, the best cutoff value of PG I/II ratio was 6.05 (sensitivity, specificity, PPV, NPV, and accuracy: 61.0%, 55.2%, 31.6%, 80.7%, and 56.7%, respectively). The AUC was 0.629 (95% CI, 0.544–0.713). For stages $\geq II$ in the antral atrophic score, considered as atrophy positive, the best cutoff value of PG I/II was 3.55 (sensitivity, specificity, PPV, NPV, and accuracy: 66.7%, 96.5%, 33.3%, 99.1%, and 95.7%, respectively), and the AUC was 0.793 (95% CI, 0.580–1.000).

The PG I and PG II best cutoff value for stages $\geq II$ in the antral atrophic score were 65.0 (the AUC were 0.583 [95% CI, 0.482–0.684] with sensitivity, specificity, PPV, NPV, and accuracy was 83.3%, 47.1%, 96.0%, 99.1%, 48.1%, respectively. while PG II level was not observed to be a sufficient marker for antral chronic gastritis by using the ROC curve (data not shown). When we used a combination PG I and PG I/II ratio for stages $\geq II$ in the antral atrophic score, the results almost the same as when using only PG I/II ratio (sensitivity, specificity, PPV, NPV, and accuracy were 66.7%, 97.8%, 44.4%, 99.1%, and 96.9%, respectively).

PG levels and the ABC method among ethnic groups

The comparison of PG levels among the ethnic groups is shown in Table 4. Among the *H. pylori*-negative subjects, PG I and II levels were significantly lower in subjects from Bangli than in those from Makassar (both $P < 0.001$). Furthermore, the PG I/II ratios were significantly lower in subjects from Bangli than in those from Makassar ($P = 0.03$). Subjects from Bangli also had significantly lower PG I/II ratios than those from Surabaya ($P = 0.004$). Although the PG I and II levels in subjects from Surabaya were significantly lower than in those from Makassar, the difference in PG I/II ratios was not significant (Table 4). The similar results also showed among the *H. pylori*-positive subjects, PG I and II levels were significantly lower in subjects from Bangli than in those from Makassar (56.3 ± 18.8 vs. 165.4 ± 87.6 , $P = 0.007$ and 14.5 ± 6.5 vs. 31.6 ± 14.3 , $P = 0.012$, respectively). However, there was no difference for the PG I/II ratios and PGs levels between Surabaya and Makassar subjects.

Table 4. Characteristics of pepsinogens among the ethnic groups.

Pepsinogens	Surabaya	Makassar	Bangli	P* value	P** value	P*** value
N	96	66	51			
PG I (ng/mL)	79.5 ± 58.1	113.2 ± 96.1	61.3 ± 54.2	0.01	0.07	<0.001
PG II (ng/mL)	12.0 ± 8.1	17.3 ± 13.0	10.6 ± 9.0	0.004	0.36	<0.001
PG I/II ratio	6.6 ± 1.8	6.5 ± 2.4	5.9 ± 1.3	0.56	0.004	0.03

PG, pepsinogen.

* Surabaya vs. Makassar.

** Bangli vs. Surabaya.

*** Bangli vs. Makassar.

<https://doi.org/10.1371/journal.pone.0176203.t004>

Table 5. ABC method among ethnic groups.

ABC groups	Surabaya (%)	Makassar (%)	Bangli (%)
Group A	94/99 (94.9)	63/75 (84.0)	50/59 (84.7)
Group B	3/99 (3.0)	7/75 (9.3)	4/59 (6.8)
Group C	0/99 (0.0)	2/75 (2.7)	4/59 (6.8)
Group D	2/99 (0.2)	3/75 (4.0)	1/59 (1.7)

<https://doi.org/10.1371/journal.pone.0176203.t005>

By using ≥ 5.5 U/mL and ≤ 3.55 as the best cutoff values of *H. pylori* antibody titers and PG I/II ratio, respectively, we determined the distribution of the four groups in each age category, according to the ABC method. Overall, group A (207 of 233, 88.8%) was the most predominant, followed by groups B (14 of 233, 6.0%), C (6 of 233, 2.6%), and D (6 of 233, 2.6%). A comparison of the ABC method among the ethnic groups is shown in Table 5. Consistent with the results obtained for PG I/II ratios, subjects from Bangli had the highest proportion of group C and lowest proportion of group A compared with those from Makassar and Surabaya. Interestingly, the percentage of group D in subjects from Bangli was the lowest among all groups.

Discussion

Although the ELISA kit for serology, which was developed by using Japanese *H. pylori* strains, it reached a high sensitivity and specificity when used in Japan and Nepal (95.2–100% and 76.2–80.0% [31] and 89.1% and 83.5% [32], respectively), we confirmed the low accuracy of the ELISA kit when used in the Indonesian population. Therefore, we calculated the best cutoff values of IgG ELISA. The same kits also showed a low sensitivity in Myanmar [33] and a low specificity when used in the Bhutan population [34]. *H. pylori* antibody titers vary greatly depending on the test kit used [35], and in general showed a lower performance in general population, but an outstanding result on the selected samples [36]. The discrepancy of *H. pylori* positivity number using ELISA and histology-immunohistochemistry associated with the advantages and disadvantages of several tests, thus the selection of the test should be performed based on these considerations [37, 38]. Importantly, the low PPV of ELISA indicated that although the kits could become the first diagnostic test, it should be confirmed with other test in positive case, especially with a high PPV value test. The Maastricht III and IV consensus suggested that “some serological tests with good sensitivity and specificity can be used to perform the initial diagnosis of infection with *H. pylori*” [39], and only validated commercial tests should be used [38, 40]. Thus, an appraisal of validation has to be performed on every serology test to the particular study population, and the antigen has to be wisely selected [41]. It might be important to develop ELISA kits using native *H. pylori* strains to the study population.

Similar to our results, previous studies [42, 43] reported that PG levels have a positive correlation with aging, particularly in men, which may be related to the increase of renal function loss and influence of hormones [44, 45]. In concordance with our data, smoking was reported to stimulate PG secretion [13]. However, long-term stimulation may exhaust the chief cells and promote atrophy of the gastric mucosa, reflected as lower PG I/II ratios in smokers [46]. In our study, drinking habits increased the PG I and II levels, in contrast to previous results [13]. Our results also confirmed the involvement of *H. pylori* as a key factor in the development of chronic and atrophic gastritis even after adjustment for other factors. *H. pylori* infection induces mucosal inflammation and the lipopolysaccharide of bacteria directly stimulates PG secretion [47], which significantly decreases after eradication concomitant with the improvement in inflammation scores [48]. The stimulation is mediated by calcium, which increases histamine and dibutyl-cyclic adenosine monophosphate [49].

Several studies suggested PG II as a good marker of gastric inflammation [45, 48, 50, 51]. Our study also showed the association between PG II and inflammation scores; however, it showed a weak correlation and was not a good discriminating marker for chronic gastritis. On the other hand, PG I/II ratios showed a high accuracy in predicting chronic gastritis, in line with previous results [14]. As Indonesia is a large country consisting of thousands of islands with a lack of endoscopy facilities, the validated PG I/II ratios offer the benefit of discriminating Indonesian patients with chronic gastritis. In addition, the PG I/II ratios also could discriminate antral atrophic gastritis, although with a lower sensitivity than that for chronic gastritis and the combination with PG I showed almost similar results. Although, overall, the PG I/II ratios had a high accuracy for predicting chronic and atrophic gastritis, they had low PPVs (23.5% and 33.3%, respectively) and high NPVs. It is suggested that PG I/II ratios have the advantages of excluding patients with chronic or atrophic gastritis (i.e., a negative result is usually a true negative result). However, the modest sensitivity result reduced the role of PG I/II ratio as valuable marker for atrophic gastritis screening. In contrast, a positive PG I/II result should be confirmed by using other diagnostic modalities. We found a low PPV when we measured the PG I/II ratio in the population with a low prevalence of atrophic gastritis; although the test had a high sensitivity and specificity [52].

In general, we found that the prevalence of atrophic gastritis in Indonesia was low, similar to our previous study [16] and in line with the incidence of gastric cancer. For example, an atrophic score >1 was observed in the antrum in only six patients and in the corpus in only one patient (Table 2), and 74.3% of patients had nonatrophic gastritis (Table 3). The next analysis based on ethnic groups revealed that the PG I/II ratios were significantly lower in subjects from Bangli than in those from Makassar and Surabaya. It is suggested that inhabitants of Bangli have a higher risk of gastric mucosal diseases than the two other ethnic groups. Although these results were in concordance with the prevalence of *H. pylori* infection among the ethnic groups, in this study, we analyzed PG levels only in *H. pylori*-negative patients. In fact, smoking but not NSAID use and alcohol drinking was significantly higher in subjects from Makassar and Bangli than in those from Surabaya. Dietary habits such as consumption of fermented fish, local spicy sauce, seafood, and fresh fruits may be risk factors, as reported in the Thai population [53].

Similar findings were also observed in the distribution of the four groups according to the ABC method. Subjects from Bangli had the highest proportion of group C and the lowest proportion of group A compared with those from Makassar and Surabaya. However, group D was the lowest in subjects from Bangli compared with the other groups. The ability of the ABC method to discriminate groups at a high risk for the development of gastric cancer was reported in a study on 8286 Japanese patients who underwent endoscopic screening for gastric cancer [4]. The authors suggested that the risk of gastric diseases is very low in group A; the risk of peptic ulcer is elevated in group B; and groups C and D had a higher risk of developing diseases resulting from atrophy of the gastric mucosa, such as gastric cancer, gastric adenoma, and hyperplastic polyps. The risk of gastric cancer is the highest in group D, followed by groups C, B, and A [4]. There are several problems in adopting the ABC method for screening for primary gastric cancer in Indonesia. First, in addition to the problem about the best cutoff for the kits, the criteria were developed for the Japanese population, which has an around 10 times higher incidence of gastric cancer than the Indonesian population (ASR, 29.9 in 100,000). Therefore, Japan has a higher prevalence of atrophic gastritis and the ABC method has a greater accuracy to exclude Japanese patients positive for atrophic gastritis (high PPV). Second, generally, Indonesia has a low prevalence of *H. pylori* infection. Therefore, there is a high possibility that a false-positive patient will be included in group D, the group with the highest risk for gastric cancer. There was no a similar concern in our previous study in a low of gastric cancer risk country, Nepal due to the modest prevalence of *H. pylori* infection in this country [32]. A validated PGs

levels and ABC method could discriminate gastric cancer risk between two group ethnics in Nepal. In addition, we found that the same kits of anti-*H. pylori* IgG reached a sufficient sensitivity and specificity among Nepalese population [32]. Further multicenter studies recruiting subjects on a large scale and from various ethnicities are required to confirm the benefits of the ABC criteria to discriminate atrophic gastritis in the Indonesian population.

The number of samples in this study was relatively low, which certainly becomes a study limitation. In addition, we included patients from three cities in three islands. Therefore, our results cannot be generalized to all of Indonesia. Recently, we are continuing our surveys to add to the sample numbers and expanding our investigations to other islands. Next, we included only dyspeptic patients in our study population, and not members of general population or health subjects. The two previous studies also only included dyspeptic patients [22, 23]. In general, gastric mucosal status was severer in the dyspeptic patients than general population or health subjects. Further studies using PGs in general population are substantial as part of a disease prevention strategy in population with low *H. pylori* prevalence, even in Indonesia.

Conclusions

Validation of indirect methods is essential before their application. We confirmed that serum PG levels are a useful biomarker for determining chronic gastritis, but a modest sensitivity for atrophic gastritis in Indonesia. The ABC method should be used with caution in areas with a low prevalence of *H. pylori*.

Supporting information

S1 Table. Socio demographic and serology data. Positivity of ELISA was determined with cut-off point ≥ 5.5 U/ml with the sensitivity, specificity, positive predictive value, and negative predictive value: 86.7%, 96.8%, 65.0%, and 99.1%, respectively. (PDF)

Author Contributions

Conceptualization: MM YY AFS.

Data curation: MM YAAR PS TU HBH MIL LAW YY.

Formal analysis: MM YY.

Funding acquisition: YY.

Investigation: MM AFS YY IAN FA IDNW UM MLP IKM PA HP TS.

Methodology: MM YY.

Project administration: MM YY.

Resources: MM AFS YY IAN FA IDNW UM MLP IKM PA HP TS.

Supervision: MM AFS YY.

Validation: MM YY.

Visualization: MM AFS YY.

Writing – original draft: MM YY.

Writing – review & editing: MM YY.

References

1. Kuipers EJ. Review article: exploring the link between *Helicobacter pylori* and gastric cancer. *Alimentary pharmacology & therapeutics*. 1999; 13 Suppl 1:3–11.
2. Correa P. Human gastric carcinogenesis: a multistep and multifactorial process—First American Cancer Society Award Lecture on Cancer Epidemiology and Prevention. *Cancer research*. 1992; 52(24):6735–40. PMID: 1458460
3. Carpenter HA, Talley NJ. Gastroscopy is incomplete without biopsy: clinical relevance of distinguishing gastropathy from gastritis. *Gastroenterology*. 1995; 108(3):917–24. PMID: 7875496
4. Miki K. Gastric cancer screening by combined assay for serum anti-*Helicobacter pylori* IgG antibody and serum pepsinogen levels—"ABC method". *Proc Jpn Acad Ser B Phys Biol Sci*. 2011; 87(7):405–14. <https://doi.org/10.2183/pjab.87.405> PMID: 21785258
5. Mirsky A, Futterman P, Kaplan S. Blood plasma pepsinogen. II. The activity of the plasma from normal subjects, patients with duodenal ulcer, and patients with pernicious anemia. *The Journal of laboratory and clinical medicine*. 1952; 40(2):188–99. PMID: 14946456
6. Samloff IM. Pepsinogens I and II: purification from gastric mucosa and radioimmunoassay in serum. *Gastroenterology*. 1982; 82(1):26–33. PMID: 7053333
7. Gritti I, Banfi G, Roi GS. Pepsinogens: physiology, pharmacology pathophysiology and exercise. *Pharmacological research: the official journal of the Italian Pharmacological Society*. 2000; 41(3):265–81.
8. Kim N, Jung HC. The role of serum pepsinogen in the detection of gastric cancer. *Gut Liver*. 2010; 4(3):307–19. PubMed Central PMCID: PMC2956341. <https://doi.org/10.5009/gnl.2010.4.3.307> PMID: 20981206
9. Sipponen P, Graham DY. Importance of atrophic gastritis in diagnostics and prevention of gastric cancer: application of plasma biomarkers. *Scand J Gastroenterol*. 2007; 42(1):2–10. <https://doi.org/10.1080/00365520600863720> PMID: 17190755
10. Miki K. Gastric cancer screening using the serum pepsinogen test method. *Gastric Cancer*. 2006; 9(4):245–53. <https://doi.org/10.1007/s10120-006-0397-0> PMID: 17235625
11. Ang TL, Fock KM, Dhamodaran S, Teo EK, Tan J. Racial differences in *Helicobacter pylori*, serum pepsinogen and gastric cancer incidence in an urban Asian population. *J Gastroenterol Hepatol*. 2005; 20(10):1603–9. <https://doi.org/10.1111/j.1440-1746.2005.03898.x> PMID: 16174081
12. Bornschein J, Selgrad M, Wex T, Kuester D, Malfertheiner P. Serological assessment of gastric mucosal atrophy in gastric cancer. *BMC Gastroenterol*. 2012; 12:10. PubMed Central PMCID: PMC3280182. <https://doi.org/10.1186/1471-230X-12-10> PMID: 22289789
13. Kikuchi S, Inaba Y, Wada O, Miki K, Tenjin H, Kaneko E, et al. The association of smoking and drinking habits with serum pepsinogens. *Int J Epidemiol*. 1995; 24(2):346–53. PMID: 7635595
14. Kiyohira K, Yoshihara M, Ito M, Haruma K, Tanaka S, Chayama K. Serum pepsinogen concentration as a marker of *Helicobacter pylori* infection and the histologic grade of gastritis; evaluation of gastric mucosa by serum pepsinogen levels. *Journal of gastroenterology*. 2003; 38(4):332–8. <https://doi.org/10.1007/s005350300060> PMID: 12743772
15. Miftahussurur M, Tuda J, Suzuki R, Kido Y, Kawamoto F, Matsuda M, et al. Extremely low *Helicobacter pylori* prevalence in North Sulawesi, Indonesia and identification of a Maori-tribe type strain: a cross sectional study. *Gut pathogens*. 2014; 6(1):42. PubMed Central PMCID: PMC4189669. <https://doi.org/10.1186/s13099-014-0042-0> PMID: 25299127
16. Miftahussurur M, Shiota S, Suzuki R, Matsuda M, Uchida T, Kido Y, et al. Identification of *Helicobacter pylori* infection in symptomatic patients in Surabaya, Indonesia, using five diagnostic tests. *Epidemiology and infection*. 2015; 143(5):986–96. <https://doi.org/10.1017/S095026881400154X> PMID: 25034254
17. Syam AF, Miftahussurur M, Makmun D, Nusi IA, Zain LH, Zulkhairi, et al. Risk Factors and Prevalence of *Helicobacter pylori* in Five Largest Islands of Indonesia: A Preliminary Study. *PloS one*. 2015; 10(11): e0140186. PubMed Central PMCID: PMC4658100. <https://doi.org/10.1371/journal.pone.0140186> PMID: 26599790
18. Miftahussurur M, Syam AF, Makmun D, Nusi IA, Zein LH, Zulkhairi, et al. *Helicobacter pylori* virulence genes in the five largest islands of Indonesia. *Gut pathogens*. 2015; 7:26. PubMed Central PMCID: PMC4594740. <https://doi.org/10.1186/s13099-015-0072-2> PMID: 26442711
19. Miftahussurur M, Syam AF, Nusi IA, Makmun D, Waskito LA, Zein LH, et al. Surveillance of *Helicobacter pylori* Antibiotic Susceptibility in Indonesia: Different Resistance Types among Regions and with Novel Genetic Mutations. *PloS one*. 2016; 11(12):e0166199. PubMed Central PMCID: PMC5131997. <https://doi.org/10.1371/journal.pone.0166199> PMID: 27906990

20. Kumala W, Rani A. Patterns of *Helicobacter pylori* isolate resistance to fluoroquinolones, amoxicillin, clarithromycin and metronidazoles. *The Southeast Asian journal of tropical medicine and public health*. 2006; 37(5):970–4. PMID: [17333742](#)
21. Makmun D. Present status of endoscopy, therapeutic endoscopy and the endoscopy training system in Indonesia. *Digestive endoscopy*. 2014; 26 Suppl 2:2–9.
22. Arinton IG. Serum gastrin level and pepsinogen I/II ratio as biomarker of *Helicobacter pylori* chronic gastritis. *Acta medica Indonesiana*. 2010; 42(3):142–6. PMID: [20724767](#)
23. Aulia D, Manz GO, Simadibrata M. Pepsinogen I concentration in organic dyspepsia patients at Gastroenterology Division, Department of Internal Medicine, Cipto Mangunkusumo Hospital. *Acta medica Indonesiana*. 2009; 41(3):107–14. PMID: [19752481](#)
24. Dixon M, Genta R, Yardley J, Correa P. Classification and grading of gastritis. The updated Sydney System. International Workshop on the Histopathology of Gastritis, Houston 1994. *Am J Surg Pathol*. 1996; 20(10):1161–81. PMID: [8827022](#)
25. Uchida T, Kanada R, Tsukamoto Y, Hijiya N, Matsuura K, Yano S, et al. Immunohistochemical diagnosis of the *cagA*-gene genotype of *Helicobacter pylori* with anti-East Asian *CagA*-specific antibody. *Cancer science*. 2007; 98(4):521–8. Epub 2007/02/08. <https://doi.org/10.1111/j.1349-7006.2007.00415.x> PMID: [17284255](#)
26. Vilaichone RK, Mahachai V, Shiota S, Uchida T, Ratanachu-ek T, Tshering L, et al. Extremely high prevalence of *Helicobacter pylori* infection in Bhutan. *World journal of gastroenterology*. 2013; 19(18):2806–10. PubMed Central PMCID: [PMC3653155](#). <https://doi.org/10.3748/wjg.v19.i18.2806> PMID: [23687418](#)
27. Shiota S, Murakami K, Fujioka T, Yamaoka Y. Population-based strategies for *Helicobacter pylori*-associated disease management: a Japanese perspective. Expert review of gastroenterology & hepatology. 2010; 4(2):149–56. PubMed Central PMCID: [PMC2896743](#).
28. Shiota S, Cruz M, Abreu JA, Mitsui T, Terao H, Disla M, et al. Virulence genes of *Helicobacter pylori* in the Dominican Republic. *Journal of medical microbiology*. 2014; 63(Pt 9):1189–96. PubMed Central PMCID: [PMC4140083](#). <https://doi.org/10.1099/jmm.0.075275-0> PMID: [24965801](#)
29. Nguyen TL, Uchida T, Tsukamoto Y, Trinh DT, Ta L, Mai BH, et al. *Helicobacter pylori* infection and gastroduodenal diseases in Vietnam: a cross-sectional, hospital-based study. *BMC gastroenterology*. 2010; 10:114. PubMed Central PMCID: [PMC2959090](#). <https://doi.org/10.1186/1471-230X-10-114> PMID: [20920280](#)
30. Nguyen LT, Uchida T, Tsukamoto Y, Trinh TD, Ta L, Mai HB, et al. Clinical relevance of *cagPAI* intactness in *Helicobacter pylori* isolates from Vietnam. *European journal of clinical microbiology & infectious diseases: official publication of the European Society of Clinical Microbiology*. 2010; 29(6):651–60. PubMed Central PMCID: [PMC3137892](#).
31. Matsuo K, Hamajima N, Suzuki T, Nakamura T, Matsuura A, Tominaga S. Better ROC Curves for a Regionally Developed *Helicobacter Pylori* Antibody Test. *Asian Pac J Cancer Prev*. 2001; 2(2):155–6. PMID: [12718648](#)
32. Miftahussurur M, Sharma RP, Shrestha PK, Maharjan RK, Shiota S, Uchida T, et al. *Helicobacter pylori* Infection and Gastric Mucosal Atrophy in Two Ethnic Groups in Nepal. *Asian Pacific journal of cancer prevention: APJCP*. 2015; 16(17):7911–6. PMID: [26625820](#)
33. Myint T, Shiota S, Vilaichone RK, Ni N, Aye TT, Matsuda M, et al. Prevalence of *Helicobacter pylori* infection and atrophic gastritis in patients with dyspeptic symptoms in Myanmar. *World journal of gastroenterology*. 2015; 21(2):629–36. PubMed Central PMCID: [PMC4296025](#). <https://doi.org/10.3748/wjg.v21.i2.629> PMID: [25605987](#)
34. Shiota S, Mahachai V, Vilaichone RK, Ratanachu-ek T, Tshering L, Uchida T, et al. Seroprevalence of *Helicobacter pylori* infection and gastric mucosal atrophy in Bhutan, a country with a high prevalence of gastric cancer. *Journal of medical microbiology*. 2013; 62(Pt 10):1571–8. PubMed Central PMCID: [PMC3799224](#). <https://doi.org/10.1099/jmm.0.060905-0> PMID: [23831768](#)
35. Burucoa C, Delchier JC, Courillon-Mallet A, de Korwin JD, Megraud F, Zerbib F, et al. Comparative evaluation of 29 commercial *Helicobacter pylori* serological kits. *Helicobacter*. 2013; 18(3):169–79. <https://doi.org/10.1111/hel.12030> PMID: [23316886](#)
36. Laheij RJ, Straatman H, Jansen JB, Verbeek AL. Evaluation of commercially available *Helicobacter pylori* serology kits: a review. *Journal of clinical microbiology*. 1998; 36(10):2803–9. PubMed Central PMCID: [PMC105068](#). PMID: [9738024](#)
37. Chey WD, Wong BC, Practice Parameters Committee of the American College of G. American College of Gastroenterology guideline on the management of *Helicobacter pylori* infection. *The American journal of gastroenterology*. 2007; 102(8):1808–25. <https://doi.org/10.1111/j.1572-0241.2007.01393.x> PMID: [17608775](#)

38. Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F, et al. Management of Helicobacter pylori infection—the Maastricht IV/ Florence Consensus Report. *Gut*. 2012; 61(5):646–64. <https://doi.org/10.1136/gutjnl-2012-302084> PMID: 22491499
39. Malfertheiner P, Megraud F, O'Morain C, Bazzoli F, El-Omar E, Graham D, et al. Current concepts in the management of Helicobacter pylori infection: the Maastricht III Consensus Report. *Gut*. 2007; 56(6):772–81. PubMed Central PMCID: PMC1954853. <https://doi.org/10.1136/gut.2006.101634> PMID: 17170018
40. Malfertheiner P, Megraud F, O'Morain CA, Gisbert JP, Kuipers EJ, Axon AT, et al. Management of Helicobacter pylori infection—the Maastricht V/Florence Consensus Report. *Gut*. 2016.
41. Miftahussurur M, Yamaoka Y. Diagnostic Methods of Helicobacter pylori Infection for Epidemiological Studies: Critical Importance of Indirect Test Validation. *BioMed research international*. 2016; 2016:4819423. PubMed Central PMCID: PMC4745376. <https://doi.org/10.1155/2016/4819423> PMID: 26904678
42. Mossi S, Meyer-Wyss B, Renner EL, Merki HS, Gamboni G, Beglinger C. Influence of Helicobacter pylori, sex, and age on serum gastrin and pepsinogen concentrations in subjects without symptoms and patients with duodenal ulcers. *Gut*. 1993; 34(6):752–6. PubMed Central PMCID: PMC1374256. PMID: 8314506
43. Pals G, Defize J, Pronk JC, Frants RR, Eriksson AW, Westerveld BD, et al. Relations between serum pepsinogen levels, pepsinogen phenotypes, ABO blood groups, age and sex in blood donors. *Annals of human biology*. 1985; 12(5):403–11. PMID: 4062235
44. Veenendaal RA, Biemond I, Pena AS, van Duijn W, Kreuning J, Lamers CB. Influence of age and Helicobacter pylori infection on serum pepsinogens in healthy blood transfusion donors. *Gut*. 1992; 33(4):452–5. PubMed Central PMCID: PMC1374057. PMID: 1582585
45. Kitamura Y, Yoshihara M, Ito M, Boda T, Matsuo T, Kotachi T, et al. Diagnosis of Helicobacter pylori-induced gastritis by serum pepsinogen levels. *Journal of gastroenterology and hepatology*. 2015; 30(10):1473–7. <https://doi.org/10.1111/jgh.12987> PMID: 25974661
46. Kikuchi S, Kurosawa M, Sakiyama T, Tenjin H. Long-term effect of smoking on serum pepsinogen values. *Journal of epidemiology / Japan Epidemiological Association*. 2002; 12(5):351–6.
47. Young GO, Stemmet N, Lastovica A, van der Merwe EL, Louw JA, Modlin IM, et al. Helicobacter pylori lipopolysaccharide stimulates gastric mucosal pepsinogen secretion. *Alimentary pharmacology & therapeutics*. 1992; 6(2):169–77.
48. Wagner S, Haruma K, Gladziwa U, Soudah B, Gebel M, Bleck J, et al. Helicobacter pylori infection and serum pepsinogen A, pepsinogen C, and gastrin in gastritis and peptic ulcer: significance of inflammation and effect of bacterial eradication. *The American journal of gastroenterology*. 1994; 89(8):1211–8. PMID: 8053437
49. Lorente S, Doiz O, Trinidad Serrano M, Castillo J, Lanas A. Helicobacter pylori stimulates pepsinogen secretion from isolated human peptic cells. *Gut*. 2002; 50(1):13–8. PubMed Central PMCID: PMC1773080. PMID: 11772960
50. Di Mario F, Cavallaro LG, Moussa AM, Caruana P, Merli R, Maini A, et al. Usefulness of serum pepsinogens in Helicobacter pylori chronic gastritis: relationship with inflammation, activity, and density of the bacterium. *Digestive diseases and sciences*. 2006; 51(10):1791–5. PMID: 17203556
51. Leja M, Wex T, Malfertheiner P. Markers for gastric cancer premalignant lesions: where do we go? *Digestive diseases*. 2012; 30(3):268–76. <https://doi.org/10.1159/000336990> PMID: 22722551
52. Altman DG, Bland JM. Diagnostic tests 2: Predictive values. *Bmj*. 1994; 309(6947):102. PubMed Central PMCID: PMC2540558. PMID: 8038641
53. Atisook K, Kachinthorn U, Luengrojanakul P, Tanwandee T, Pakdirat P, Puapairoj A. Histology of gastritis and Helicobacter pylori infection in Thailand: a nationwide study of 3776 cases. *Helicobacter*. 2003; 8(2):132–41. PMID: 12662381

Gastric mucosal status

ORIGINALITY REPORT

19%

SIMILARITY INDEX

11%

INTERNET SOURCES

17%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Atsushi Takeoka, Jun Tayama, Hironori Yamasaki, Masakazu Kobayashi et al. "Impact of Helicobacter pylori Immunoglobulin G Levels and Atrophic Gastritis Status on Risk of Metabolic Syndrome", PLOS ONE, 2016
Publication 2%

- 2** Submitted to iGroup
Student Paper 1%

- 3** lib.bioinfo.pl
Internet Source 1%

- 4** www.science.gov
Internet Source 1%

- 5** archive.rsna.org
Internet Source 1%

- 6** Hyun Young Kim. "Clinical meaning of pepsinogen test and Helicobacter pylori serology in the health check-up population in Korea", European Journal of Gastroenterology & Hepatology, 06/2009
Publication 1%

7

Kitamura, Yoko, Masaharu Yoshihara, Masanori Ito, Tomoyuki Boda, Taiji Matsuo, Takahiro Kotachi, Shinji Tanaka, and Kazuaki Chayama. "Diagnosis of Helicobacter pylori -induced gastritis by serum pepsinogen levels : Diagnosis of chronic gastritis by serum markers", Journal of Gastroenterology and Hepatology, 2015.

Publication

1%

8

Katsunori Iijima, Hitoshi Sekine, Tomoyuki Koike, Akira Imatani, Shuichi Ohara, Tooru Shimosegawa. "Serum pepsinogen concentrations as a measure of gastric acid secretion in Helicobacter pylori-negative and -positive Japanese subjects", Journal of Gastroenterology, 2005

Publication

1%

9

Yukitoshi Sakao, Mitsushige Sugimoto, Hitomi Ichikawa, Shu Sahara et al. "Severity of Gastric Mucosal Atrophy Is the Major Determinant of Plasma Ghrelin Level in Hemodialysis Patients", American Journal of Nephrology, 2016

Publication

<1%

10

Burucoa, Christophe, Jean-Charles Delchier, Anne Courillon-Mallet, Jean-Dominique de Korwin, Francis Mégraud, Frank Zerbib, Josette Raymond, and Jean-Louis Fauchère. "Comparative Evaluation of 29 Commercial

<1%

Helicobacter pylori Serological Kits",
Helicobacter, 2013.

Publication

11

Leung, Wai K., Enders K. W. Ng, Candice C. H. Lam, Kui-Fat Chan, Wing Y. Chan, Alex C. M. Auyeung, Justin C. Y. Wu, Jessica Y. L. Ching, James Y. W. Lau, and Joseph J. Y. Sung. "Helicobacter pylori infection in 1st degree relatives of Chinese gastric cancer patients", Scandinavian Journal of Gastroenterology, 2006.

Publication

<1%

12

S. Sanduleanu. "Double gastric infection with Helicobacter pylori and non-Helicobacter pylori bacteria during acid-suppressive therapy: increase of pro-inflammatory cytokines and development of atrophic gastritis", Alimentary Pharmacology and Therapeutics, 8/22/2001

Publication

<1%

13

www.tandfonline.com

Internet Source

<1%

14

Yoshiko Nakayama, Yingsong Lin, Minoru Hongo, Hiroya Hidaka, Shogo Kikuchi. "infection and its related factors in junior high school students in Nagano Prefecture, Japan ", Helicobacter, 2017

Publication

<1%

15

Takeishi Shimamoto, Nobutake Yamamichi, Shinya Kodashima, Yu Takahashi et al. "No Association of Coffee Consumption with Gastric Ulcer, Duodenal Ulcer, Reflux Esophagitis, and Non-Erosive Reflux Disease: A Cross-Sectional Study of 8,013 Healthy Subjects in Japan", PLoS ONE, 2013

Publication

<1%

16

Yasuhiko Abe. "Barrett's esophagus is characterized by the absence of *Helicobacter pylori* infection and high levels of serum pepsinogen I concentration in Japan", Journal of Gastroenterology and Hepatology, 01/2009

Publication

<1%

17

Banno, Shogo, Yoshifuji Matsumoto, Yoshiki Sugiura, Takeo Yoshinouch, Hideo Shibata, and Ryuzo Ueda. "Seroprevalence of *Helicobacter pylori* and association with atrophic gastritis in patients with Sjögren's syndrome", Japanese Journal of Rheumatology, 1999.

Publication

<1%

18

Svetlana Kurilovich, Anna Belkovets, Oleg Reshetnikov, Tatyana Openko, Yulia Ragino. "Sa1952a The First Steps in the Development of Gastric Cancer Riskometry: The Siberian Retrospective Case-Control Study. Novosibirsk State Medical University, Russian Federation",

<1%

Gastroenterology, 2014

Publication

19

Xian-Zhe Chen, Cheng-Zhi Huang, Wei-Xian Hu, Ying Liu, Xue-Qing Yao. "Gastric Cancer Screening by Combined Determination of Serum Helicobacter pylori Antibody and Pepsinogen Concentrations", Chinese Medical Journal, 2018

Publication

<1%

20

Helicobacter pylori, 1998.

Publication

<1%

21

ainfo.cnptia.embrapa.br

Internet Source

<1%

22

bm Cresnotes.biomedcentral.com

Internet Source

<1%

23

Fumiaki Kitahara, Ryoichi Shimazaki, Tadashi Sato, Yuichiro Kojima, Atsuro Morozumi, Masayuki A. Fujino. "Severe atrophic gastritis with Helicobacter pylori infection and gastric cancer", Gastric Cancer, 1998

Publication

<1%

24

medlib.mef.hr

Internet Source

<1%

25

eprints.lib.hokudai.ac.jp

Internet Source

<1%

26

refubium.fu-berlin.de

Internet Source

<1%

27

docplayer.net

Internet Source

<1%

28

"Stratification of Gastric Cancer Risk by H. pylori Infection", *Helicobacter pylori*, 2016.

Publication

<1%

29

Ko Miura, Hiroyuki Okada, Yoshiyasu Kouno, Hiromitsu Kanzaki et al. "Actual Status of Involvement of *Helicobacter pylori* Infection That Developed Gastric Cancer from Group A of ABC (D) Stratification - Study of Early Gastric Cancer Cases That Underwent Endoscopic Submucosal Dissection", *Digestion*, 2016

Publication

<1%

30

www.biohit.cn

Internet Source

<1%

31

Dipti Chourasia, Asha Misra, Shweta Tripathi, Narendra Krishnani, Uday C. Ghoshal. "Patients with *Helicobacter pylori* infection have less severe gastroesophageal reflux disease: a study using endoscopy, 24-hour gastric and esophageal pH metry", *Indian Journal of Gastroenterology*, 2011

Publication

<1%

32

mafiadoc.com

Internet Source

<1%

Kato, M., and M. Asaka. "Recent Development

33

of Gastric Cancer Prevention", Japanese Journal of Clinical Oncology, 2012.

Publication

<1%

34

Kikuchi, Shogo, Michiko Kurosawa, Tsuguo Sakiyama, and Hiroshi Tenjin. "Long-term Effect of Smoking on Serum Pepsinogen Values.", Journal of Epidemiology, 2002.

Publication

<1%

35

Y Nagahata. "Helicobacter pylori may cause "reflux" gastritis after gastrectomy", Journal of Gastrointestinal Surgery, 1997

Publication

<1%

36

Naoyo Ishikura, Yoshiaki Usui, Hidemi Ito, Yumiko Kasugai et al. "Helicobacter pylori (HP) infection alone, but not HP-induced atrophic gastritis, increases the risk of gastric lymphoma: a case-control study in Japan", Annals of Hematology, 2019

Publication

<1%

37

T. Ohkusa. "Improvement in serum pepsinogens and gastrin in long-term monitoring after eradication of Helicobacter pylori: comparison with H. pylori-negative patients", Alimentary Pharmacology and Therapeutics, 7/2004

Publication

<1%

38

Young Woon Chang. "Clinical parameters including serum pepsinogen level and

<1%

management strategy in patients with premalignant gastric dysplasia :", European Journal of Gastroenterology & Hepatology, 05/2011

Publication

39

P. Zentilin. "Helicobacter pylori infection is not involved in the pathogenesis of either erosive or non-erosive gastro-oesophageal reflux disease", Alimentary Pharmacology and Therapeutics, 4/2003

Publication

<1%

40

Karpouza, Angeliki, Elisabeth Samouilidou, Stefanos Karagiannis, Vasiliki Kostopoulou, Maria Sotiropoulou, Eleftheria Roma, Kalliopi Petraki, and Spyros Michopoulos. "Patients with duodenal ulcer have lower levels of serum cholesterol compared to other dyspeptic patients independently of Helicobacter pylori status", Scandinavian Journal of Gastroenterology, 2008.

Publication

<1%

41

bmcinfectdis.biomedcentral.com

Internet Source

<1%

42

Masanori Ito, Tomoyuki Boda, Takahiro Kotachi, Mariko Kiso et al. "Chapter 6 Serum Markers", Springer Nature America, Inc, 2019

Publication

<1%

43 Zheng, Kui-Cheng, Kazuo Aoki, Xiao-Qing Li, Shu-Guang Lin, Bing-Shan Wu, Wen-Ling Zhong, Tie-Hui Chen, Song Lin, Jian-Wang You, and Chang Su. "Serum Pepsinogens, Gastrin-17 and Helicobacter pylori Antibody in the Residents of Two Cities in China with Distinct Mortality Rates of Gastric Cancer", The Tohoku Journal of Experimental Medicine, 2012.
Publication

44 www.pepsinogen.org
Internet Source

45 Akira Yokoyama. "Helicobacter pylori, chronic atrophic gastritis, inactive aldehyde dehydrogenase-2, macrocytosis and multiple upper aerodigestive tract cancers and the risk for gastric cancer in alcoholic Japanese men", Journal of Gastroenterology and Hepatology, 2/2007
Publication

46 www0.nih.go.jp
Internet Source

47 www.wjgnet.com
Internet Source

48 obgyn.onlinelibrary.wiley.com
Internet Source

bmcpregnancychildbirth.biomedcentral.com

49

Internet Source

<1%

50

Cara L Cooke, Javier Torres, Jay V Solnick. " Biomarkers of associated gastric cancer ", Gut Microbes, 2014

Publication

<1%

51

Marcis Leja. "The Validity of a Biomarker Method for Indirect Detection of Gastric Mucosal Atrophy Versus Standard Histopathology", Digestive Diseases and Sciences, 09/03/2009

Publication

<1%

52

my-oscar.nhs.uk

Internet Source

<1%

53

www.thieme-connect.de

Internet Source

<1%

54

www.scielo.br

Internet Source

<1%

55

Mohamed Ramelah, Ahmad Aminuddin, Hanafiah Alfizah, Mohd Rose Isa et al. " gene variants in Malaysian strains isolated from patients of different ethnic groups ", FEMS Immunology & Medical Microbiology, 2005

Publication

<1%

56

link.springer.com

Internet Source

<1%

57

www.id-press.eu

Internet Source

<1%

58

www.physiology.org

Internet Source

<1%

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On