

THE POTENCY OF WATER  
LEAVES EXTRACT OF BAY  
(*Syzygium polyanthum*) AND  
PAPAYA (*Carica papaya*) AS  
LARVICIDES TO FILARIA  
VECTOR *Culex*  
*queinquefasciatus* (Diptera:  
*Culicidae*)

---

**Submission date:** 16-Jun-2023 11:05AM (UTC+0700)  
by Rinza Rahmawati Samsudin

**Submission ID:** 2117073164

**File name:** 2022\_OKTOBER\_JURNALL\_BIOMA\_AR\_VL\_RR\_4\_point.pdf (351.08K)

**Word count:** 2227

**Character count:** 12302



**1**  
**THE POTENCY OF WATER LEAVES EXTRACT OF BAY (*Syzygium polyanthum*) AND PAPAYA (*Carica papaya*) AS LARVICIDES TO FILARIA VECTOR *Culex quinquefasciatus* (Diptera: Culicidae)**

Vella Rohmayani<sup>1\*</sup>, Anindita Riesti Retno Arimurti<sup>2</sup>, Rinza Rahma<sup>2</sup>, ti Samsudin<sup>2</sup>

<sup>1</sup>Department of Medical Laboratory Technologist, Vocational Program, Faculty of Health Sciences, University of Muhammadiyah Surabaya.

Jl. Sutorejo No.59, Surabaya, East Java, Indonesia 60112

<sup>2</sup>Medical Laboratory Technology, Study Program DIII, Faculty of Health Sciences, University of Muhammadiyah Surabaya.

Jl. Sutorejo No.59, Surabaya, East Java, Indonesia 60113

\*Corresponding author: [vella\\_rohmayani@um-surabaya.ac.id](mailto:vella_rohmayani@um-surabaya.ac.id)

**ARTICLE INFO**

**Article history**

Submission

December 26<sup>th</sup>, 2021

Revision

March 14<sup>th</sup>, 2022

Accepted

June 9<sup>th</sup>, 2022

**Keywords:**

Bay leaves (*Syzygium polyanthum*)

*Culex quinquefasciatus*

Larvacide

Papaya leaves (*Carica papaya*)

**ABSTRACT**

**11**  
*Culex quinquefasciatus* is a mosquito that acts as a vector of filariasis disease. In Indonesia filariasis is still one of the endemic diseases. This proves that the handling of vector *Culex quinquefasciatus* is still not done optimally. One way that can be done to control mosquito vectors one of them is to use larvacide. Larvacide which is widely used today is still synthetic. Although very efficient in causing mosquito mortality, larvacide synthesis harms the environment and can cause vectors resistance. Then another larvacide alternative is needed that is effective and does not cause negative effects. Plants can be a choice of larvacide from nature. The study aims to find out the potential of the water extract of bay leaves and papaya leaves in killing the larvae of the *Cx. quinquefasciatus* mosquito as well as the value of  $LC_{50}$  after 24-hour observation. The type of research used is experimental research. The concentration of the water extract of bay leaves and papaya leaves used is 10%, 30%, 50%, 70%, and 90% with three times replication.  $LC_{50}$  is calculated using the Finney formula (1971), the calculation is carried out using application SPSS statistical version 26. Based on the research conducted, the value of bay leaf water extract  $LC_{50} = 62.89$  g/ml and papaya leaf water extract  $LC_{50} = 45.10$  g/ml. So based on the  $LC_{50}$  value, it can be concluded that papaya leaf water extract is more effective in causing mortality *Cx. quinquefasciatus* compared with bay leaf water extract.

**INTRODUCTION**

Filariasis is a disease that transmitted through mosquito vector intermediaries. In Indonesia, filariasis disease is still one of the endemic diseases that cause health problems

in the community until now. There were 10,681 cases in Indonesia in 2018 that were spread in 236 districts/cities from 28 provinces are declared as filariasis endemic. The highest cases of filariasis in Indonesia occurred in Papua Province with 3,615 cases, then East Nusa Tenggara with 1,542 cases, and followed by West Java province with the 985 cases (Indah, 2019).

The main step that can be done to prevent the transmission of transmitted diseases through vector intermediaries is controlling the disease-transmitting vector (Fouque & Reeder, 2019). One of the ways to control the disease vectors is chemical using larvicides. However, the larvicides that are widely used today are still synthetic larvicides that cause various negative effects <sup>5</sup> such as causing environmental pollution, and resistance and can be toxic to non-target organisms (Kumar *et al.*, 2014; Princess & Kurnia, 2018). Therefore it is necessary to develop natural larvicides that are cheaper, eco-friendly does not toxic to non-target organisms, biodegradable, and do not destroy ecosystems (Astriani & Widawati, 2016; Kumar *et al.*, 2014).

Plants that may potentially be used as natural larvicides are bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*) because they contain secondary metabolite compounds. According to Utami & Puspaningtyas (2014), papaya leaves contain essential oils, lactones, phenols, saponins, and polyphenols. According to Cahyati *et al.* (2019), papaya leaves contain the enzymes papain, saponins, tannins, flavonoids, steroids, karpaina alkaloids, glycosids, karposids, saccharides, dextrose, and fructose.

Tannin compounds, saponins, flavonoids, and alkaloids have toxic properties for animals, so they can be used as larvicides (Ariesta, 2013; Gutierrez *et al.*, 2014). Tannins can damage the digestive part of insects, as well as cause disruptions in the function of organism water absorption (Mardiana *et al.*, 2009; Pradani, 2009). Saponins can inhibit growth, and development and interfere insect reproductive system. Alkaloids and flavonoids can inhibit insect-eating, besides essential oils and flavonoids can work as respiratory toxins (Pradani, 2009).

Research of potential testing by bay leaf extract and papaya leaves as natural larvicides has been done by researchers before. This research aims to increase information about the potential of bay leaves and papaya leaves using the water extract method, the use of squeeze method can be directly applied by the general public, in

contrast to extract method which must be carried out in laboratory. Therefore, it can be acknowledged the potential or effectiveness of each water <sup>1</sup> extract of bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*) towards the death rate of *Cx. quenequefasciatus* mosquitoes.

## MATERIALS AND METHODS

### *Tools and Materials*

The tools used in this study are blenders, measuring cups, digital scales, beaker glass, petri dishes, filters, stirrers, rubber, stationery, and la bel paper. While the ingredients used are bay leaves (*Syzygium polyanthum*), papaya leaves (*Carica papaya*), aquades, and *Cx. quenequefasciatus* mosquito larvae. As for the leaves used are dark green leaves.

### *Research Procedures*

First, make sharpening bay leaves (*Syzygium polyanthum*) and papaya (*Carica papaya*). Papaya leaves (*Syzygium polyanthum*) 100 grams. Second, washed thoroughly, dried, and cut into small pieces to facilitate the smoothing process. Then, papaya leaves (*Syzygium polyanthum*) are put in a blender and added 100 ml of akuades. The blender results are water extractd and filtered. Furthermore, the result of this water extract is a 100% concentration. The result of this water extract is diluted using the formula:

$$V_1 \times M_1 = V_2 \times M_2$$

<sup>6</sup>escription:

V1 = Initial volume

M1 = Initial concentration

V2 = Final volume

M2 = Final concentration

(Nurhaifah & Sukesi, 2015)

The way to make a water extract of papaya leaf (*Carica papaya*) is similar to how to make a water extract of bay leaves (*Syzygium polyanthum*). Next, make a water extract on each leave with concentrations of <sup>7</sup> 10% (P1), 30% (P2), 50% (P3), 70% (P4), and 90% (P5). After that, the *Cx. quenequefasciatus* instar III larvae are put into a petri dish with as many as 25 individu for each treatment. Then, the petri dish treatment was closed


and incubated for 24 hours after treatment. The experiment are repeated 3 times. <sup>5</sup> After 24 hours, the number of dead larvae was calculated and recorded for each treatment on the data tabulation table. Mosquito larvae are claimed as dead larvae are larvae if the condition is not moving anymore and/or the body is destroyed.

### Data Analysis

Calculated LC50 following to Probit analysis by formula Finney 1971). The calculating using SPSS application version 26.

## RESULTS AND DISCUSSION

The deaths number of *Cx. quinquefasciatus* larvae after being treated by bay leaf and papaya leaves can be seen in **Figure 1**. Based on **Figure 1** it can be observed that in treatment 1, 10% of the water extract of bay leaves and papaya leaves causes the death of larvae of *Cx. quinquefasciatus*, the percentage of death in mosquito larvae will increase proportionally to the amount of concentration of water extract given.



**Figure 1.** Death average of *Cx. quinquefasciatus* larvae after 24 hours giving a water extract of bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*)

The lowest percentage of deaths occurred in the treatment with a 10% concentration and the highest percentage of *Cx. quinquefasciatus* mosquito larvae deaths occurred in the treatment with a 90% concentration both in the treatment of bay leaf water extract or papaya leaf water extract. Based on these results, it can be seen that the higher the concentration or content of active compounds in bay leaves (*Syzigium polyanthum*) and papaya leaves, the mortality rate in *Cx. quinquefasciatus* larvae will increase too.

LC<sub>50</sub> for bay leaves = 62.89 g/ml and LC<sub>50</sub> for papaya leaves = 45.10 g/ml. Based on these results, the LC<sub>50</sub> papaya leaves is smaller than the LC 50 bay leaves, it can be concluded that the papaya leaf water extract is more effective than the bay leaf water extract. Both Bay leaf (*Syzygium Polyanthum*) and papaya leaves (*Carica papaya*) contain various secondary metabolite compounds, such as flavonoids, tannins, saponins, phenolic hydroquinon alkaloids, steroids, and triterpenoids that are toxic to animals (Cahyati *et al.*, 2019; Sulistiyani *et al.*, 2014; Utami & Puspaningtyas, 2014).

<sup>9</sup> Based on the results of the research, tannin compounds can damage the digestive to some parts of insects and able to make disturb the function of water absorption in organisms (Mardiana *et al.*, 2009; Pradani, 2009). Saponins can cause a decrease in insect food intake, causing stunted growth, development, and disruption of the insect reproductive system. In addition, other active compounds such as alkaloids and flavonoids are also able to act as compounds that inhibit eating insects, essential oils and flavonoids can also work as respiratory toxins (Pradani, 2009).

## CONCLUSION

The LC<sub>50</sub> of papaya leaves (*Carica papaya*) is smaller than the LC<sub>50</sub> of bay leaves (*Syzygium polyanthum*). It can be concluded that the papaya leaves water extract is more effective than the bay leaf water extract as larvacide to *Cx. quinquefasciatus*.

## REFERENCES

- Ariesta, A. A. (2013). *Uji Efektifitas Larutan Daun Pepaya (Carica papaya) Sebagai Larvasida Terhadap Kematian Larva Nyamuk Aedes aegypti di Laboratorium B2P2VRP*. Balai Besar Litbang Vektor dan Reservoir Penyakit. Salatiga.
- Astriani, Y., & Widawati, M. (2016). Potensi tanaman di Indonesia sebagai larvasida

- alami untuk *Aedes aegypti*. *SPIRAKEL*, 8(2), 37–46. <https://doi.org/10.22435/spirakel.v8i2.6166>
- Cahyati, W. H., Asmara, W., Umniyati, S. R., & Mulyaningsih, B. (2017). The phytochemical analysis of hay infusions and papaya leaf juice as an attractant containing insecticide for *Aedes aegypti*. *Jurnal Kesehatan Masyarakat*, 12(2), 218–224. <https://doi.org/10.15294/kemas.v12i2.6223>
- Cahyati, W. H., Asmara, W., Umniyati, S. R., & Mulyaningsih, B. (2019). Biolarvicidal effects of papaya leaves juice against *Aedes aegypti* Linn larvae. *Journal of International Dental and Medical Research*, 12(2), 780–785. [http://www.jidmr.com/journal/wp-content/uploads/2019/07/70\\_M17\\_449\\_Widya\\_Hary\\_Cahyati\\_Layout.pdf](http://www.jidmr.com/journal/wp-content/uploads/2019/07/70_M17_449_Widya_Hary_Cahyati_Layout.pdf)
- Dwiyanti, R. D., Dediq, R., & Thuraidah, A. (2017). Daya bunuh ekstrak air daun salam (*Syzygium polyanthum*) terhadap larva *Aedes* sp. *Medical Laboratory Technology Journal*, 3(1), 17. <https://doi.org/10.31964/mltj.v3i1.151>
- Fouque, F., & Reeder, J. C. (2019). Impact of past and on-going changes on climate and weather on vector-borne diseases transmission: A look at the evidence. *Infectious Diseases of Poverty*, 8(1), 1–9. <https://doi.org/10.1186/s40249-019-0565-1>
- Gutierrez, P. M., Antepuesto, A. N., Eugenio, B. A. L., & Santos, M. F. L. (2014). Larvicidal activity of selected plant extracts against the dengue vector *Aedes aegypti* mosquito. *International Research Journal of Biological Sciences*, 3(4), 23–32.
- Indah, I. S. (2019). *Situasi Filariasis di Indonesia*. Pusat Data dan Informasi Kementerian Kesehatan RI (p. 12). <https://pusdatin.kemkes.go.id/article/view/20010200002/infodatin-situasi-filariasis-di-indonesia.html>
- Kumar, S., Mishra, M., Wahab, N., & Warikoo, R. (2014). Larvicidal, repellent, and irritant potential of the seed-derived essential oil of *Apium graveolens* against dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *Frontiers in Public Health*, 2(SEP), 1–6. <https://doi.org/10.3389/fpubh.2014.00147>
- Mardiana, Supraptini, & Aminah, N. S. (2009). *Datura metel* Linnaeus sebagai insektisida dan larvasida botani serta bahan baku obat tradisional. *Media Penelitian dan Pengembangan Kesehatan*, XIX. <https://agris.fao.org/agris-search/search.do?recordID=DJ2020000225>
- Pradani, F. Y. (2009). Indeks pertumbuhan larva *Aedes aegypti* L. yang terdedah dalam ekstrak air kulit jengkol (*Pithecellobium lobatum*). *Aspirator: Journal of Vector-borne Disease Studies*, 1(2). <https://ejournal2.litbang.kemkes.go.id/index.php/aspirator/issue/archive>
- Sulistiyani, Falah, S., Wahyuni, W. T., Sugahara, T., Tachibana, S., & Syaefudin. (2014). Cellular mechanism of the cytotoxic effect of extracts from *Syzygium polyanthum* leaves. *American Journal of Drug Discovery and Development*, 4(2), 90–101. <https://doi.org/10.3923/ajdd.2014.90.101>

- Utami, P., & Puspaningtyas, D. E. (2014). *The Miracle Of Herbs*. Agro Media Pustaka.
- Waskito, P. E., & Cahyati, W. H. (2018). Efektivitas granul daun salam (*Eugenia polyantha* Wight) sebagai larvasida nyamuk *Aedes aegypti*. *SPIRAKEL*, 10(1), 12–20. <http://ejournal2.litbang.kemkes.go.id/index.php/spirakel/article/view/546>



# THE POTENCY OF WATER LEAVES EXTRACT OF BAY (*Syzygium polyanthum*) AND PAPAYA (*Carica papaya*) AS LARVICIDES TO FILARIA VECTOR *Culex quinquefasciatus* (Diptera: Culicidae)

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

13%

INTERNET SOURCES

%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://repository.um-surabaya.ac.id">repository.um-surabaya.ac.id</a> Internet Source	4%
2	<a href="http://www.panafrican-med-journal.com">www.panafrican-med-journal.com</a> Internet Source	2%
3	<a href="http://journal.unismuh.ac.id">journal.unismuh.ac.id</a> Internet Source	1%
4	<a href="http://journal.unnes.ac.id">journal.unnes.ac.id</a> Internet Source	1%
5	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	1%
6	<a href="http://www.coursehero.com">www.coursehero.com</a> Internet Source	1%
7	<a href="http://sinta.unud.ac.id">sinta.unud.ac.id</a> Internet Source	1%
8	<a href="http://journal-center.litpam.com">journal-center.litpam.com</a> Internet Source	1%

[www.scribd.com](http://www.scribd.com)

9

Internet Source

1 %

---

10

[journal.poltekkes-mks.ac.id](http://journal.poltekkes-mks.ac.id)

Internet Source

<1 %

---

11

[nopr.niscair.res.in](http://nopr.niscair.res.in)

Internet Source

<1 %

---

Exclude quotes  On

Exclude matches  < 5 words

Exclude bibliography  On