


GACETA MEDICA DE CARACAS

Fundada el 13 de marzo de 1893

por el

DR. LUIS RAZETTI

Organo de la Academia Nacional de Medicina
y del Congreso Venezolano de Ciencias Médicas



VOLUMEN 129 - N° 4

Octubre - Diciembre 2021

Caracas - Venezuela

Indizada en

AMELYCA, AURA, BASE, BASE DE DATOS PERIODICA, BIBLAT,
BIREME, CABELLS SCHOLARLY ANALYTICS, CROSS REF, DIRECTORY
OF OPEN ACCESS JOURNAL(DOAJ), Elsevier Scopus, EuroPub, GOOGLE
ANALYTICS, GOOGLE SCHOLAR, Latindex, LILACS, MIAR, ORCID,
PUBLONS, PubMed, REDIB, Research Gate, REPOSITORIO SABER UCV,
SCIELO, SCIMAGO, WEB OF SCIENCE, WORLDCAT

Gaceta Médica de Caracas

Órgano oficial de la Academia Nacional de Medicina
y del Congreso Venezolano de Ciencias Médicas

Fundada el 13 de marzo de 1893

por el

Dr. Luis Razetti

Primer número publicado el 15 de abril de 1893

Editor En Jefe

Dr. Enrique Santiago López-Loyo

Volumen 129

Nº 4

Octubre-Diciembre 2021

Í N D I C E

EDITORIAL

Aporte institucional de la Academia Nacional de Medicina de Venezuela en tiempos de pandemia: vacunación contra la COVID-19

Enrique Santiago López-Loyo, Huniades Urbina-Medina, José Esparza

801

ARTÍCULOS ORIGINALES

Terapia potencial de la planta de Sambiloto (*Andrographis paniculata*) mediante análisis de compuestos múltiples Handayani, Wiwik Winarningsih

814

Depresión, consumo de alcohol y calidad de vida en el adulto mayor con diabetes tipo 2
Humberto Frances-Salgado, Hayanín Cortés García, Jorge Escobedo de la Peña

823

Niveles de HbA1c con albuminuria en pacientes con diabetes mellitus
Prema Hapsari Hidayati, Indah Lestari Daeng Kanang, Dzulrizka Razak, Resky Pratiwi Lambang Basri

852

Factores influyentes en el riesgo de dependencia al ejercicio físico en usuarios de gimnasio colombianos Tatiana Castañeda Quirama, Jorge Emiro Restrepo

858

El efecto de la suplementación con aceite de Krill sobre los niveles de glucosa en sangre en ratas hiperglicémicas inducida por aloxano

Nesyana Nurmadilla, Anastasia Nugraha Pratiwi, Aryanti Bamahry, Nevi Sulvita, Yani Sodiqah

871

Niveles de fragilidad en el adulto mayor no institucionalizado y características sociodemográficas
María Victoria Quintero-Cruz, Yaneth Herazo-Beltrán, Miguel Terán-Martínez, Lay Viecco-Montero, Alexander Coronel-Verdecia, Yecid Villegas-Padilla, Jorgina Cure Manchego

877

Estímulo del agente activador Solanum melongena L. sobre la calidad del esperma a través del perfil lipídico de Rattus norvegicus con inducción hiperlipídica

Rinza Rahmawati Samsudin, Nur Vita Purwaningsih, Rahma Widystutti

884

Trastorno de ansiedad generalizada en los trabajadores del área de la salud de Salud Chacao
Daniel Ernesto Carvallo Ruiz, Elizabeth Natalia Martínez Núñez, Jesús Adolfo Monasterios Montero, María José Cotúa Acosta, Daniel Enrique Anato Moreno

893

Efectos del diálogo interno positivo sobre el nivel de ansiedad de los pacientes con enfermedad coronaria
Uswatun Hasanah, Dede Nasrullah, Rifma Yuniar, Nugroho Ari Wibowo, Reliani

906

ARTÍCULOS DE REVISIÓN

- Enfermedades tropicales y embarazo: revisión narrativa
Ana Carvajal, Carlos Cabrera Lozada, Jeiv Gómez 913

- Dispositivos no invasivos de neuromonitorización para la detección de isquemia cerebral intraoperatoria: una revisión exploratoria
Esteban García-Espitia, Juan Pablo Pacheco-Alba, Eduardo Tuta-Quintero, Daniel Botero-Rosas 933

- Medicamentos en el ambiente: un problema de salud pública. Revisión sistemática
Bartolomé Manzolillo 942

CASOS CLÍNICOS

- Angeitis de rama congelada en un paciente positivo al virus de la inmunodeficiencia humana con retinitis por citomegalovirus
Noel Roig-Marín, Pablo Roig-Rico, Vicente Navarro-Ibáñez 951

- Leiomiosarcoma vulvar. Una patología inusual
Franco Calderaro Di Ruggiero, Viviana García, Johatson Freytes, Carmen Silva, Lorenzo J Calderaro Fernández 956

- Lipoleiomioma uterino. Reporte de un caso
Viviana García, Franco Calderaro Di Ruggiero, Johatson Freytes 962

PERSPECTIVAS

- ¿Cómo terminan las pandemias? el futuro de la COVID-19
José Esparza, Gilberto Vizcaíno 968

HISTORIA Y FILOSOFÍA DE LA MEDICINA

- Manejo del “mal de costado”, una receta médica en el Nuevo Reino de Granada
Eduardo Tuta-Quintero, Julio César Martínez-Lozano, Ignacio Briceño-Balcázar, Diego Sierra-Barbosa, Alberto Gómez-Gutiérrez 975

- Serpentes, Insecta et Rabiem: manejo de las mordeduras de serpiente, una emergencia médica durante el período colonial
Jhonnatan Coronado-Sarmiento, Ignacio Briceño-Balcáza, Julio César Martínez-Lozano 981

BIOGRAFÍAS

- Semblanza de la Promoción de Médicos 1962 de la Universidad Central de Venezuela
Leopoldo Briceño-Iragorry 989

- Promoción de médicos-cirujanos “Bicentenario de los estudios médicos Dr. Lorenzo Campins y Ballester”, 1963.
Algunas anotaciones
Claudia Blandenier de Suárez, Luis Herrera García, Rosa Marcano de Cuenca, José Luis Cevallos, César Blanco Rengel, Leopoldo Moreno Brandt 996

VARIOS

- La Gaceta Médica de Caracas hace... 100... 50... 25 años.
Enrique Santiago López-Loyo 1012

- Vida de la Academia, Resúmenes de los trabajos presentados y Notas Bibliográficas
Enrique Santiago López-Loyo 1017

- Índice Volumen 129. Dr. Rogelio Pérez D' Gregorio 1036

Gaceta Médica de Caracas

Official Journal of the National Academy of Medicine
and The Venezuelan Congress of Medical Sciences

Founded March 13, 1893

By

Dr. Luis Razetti

First number published on April 15, 1893

Editor In Chief

Dr. Enrique Santiago López-Loyo

Volume 129

No. 4

October-December 2021

I N D E X

EDITORIAL

- Institutional contribution of the venezuelan academy of medicine in times of pandemic: Vaccination against COVID-19
Enrique Santiago Lopez-Loyo, Huniades Urbina-Medina, Jose Esperza 801

ORIGINAL ARTICLES

- Potential Therapy of Sambiloto Plant (*Andrographis paniculata*) Using Multi-Compounds Analysis
Handayani, Wiwik Winarningsih 814
- Depression, alcohol drinking and quality of life in older adults with type 2 diabetes
Humberto Frances-Salgado, Hayanin Cortes Garcia, Jorge Escobedo de la Pena 823
- HbA1c levels with albuminuria in diabetes mellitus Patients
Prema Hapsari Hidayati, Indah Lestari Daeng Kanang, Dzulrizka Razak, Resky Pratiwi Lambang Basri 852
- Influential factors in the risk of dependence on physical exercise in Colombian gym users
Tatiana Castaneda Quirama, Jorge Emiro Restrepo 858
- The effect of Krill oil supplementation on blood glucose levels in alloxan-induced hyperglycemic Rats
Nesyana Nurmadiilla, Anastasia Nugraha Pratiwi, Aryanti Bamahry, Nevi Sulvita, Yani Sodiqah 871
- Frailty levels in non-institutionalized older adults and sociodemographic characteristics
Maria Victoria Quintero-Cruz, Yaneth Herazo-Beltran, Miguel Teran-Martinez, Lay Viecco-Montero,
Alexander Coronel-Verdecia, Yecid Villegas-Padilla, Jorgina Cure Manchego 877
- Stimulus from the activator agent *Solanum melongena* L. on sperm quality through the lipid profile of *Rattus norvegicus* with hyperlipid induction
Rinza Rahmawati Samsudin, Nur Vita Purwaningsih, Rahma Widystuti 884
- Generalized anxiety disorder in the health workers of Salud Chacao
Daniel Ernesto Carvallo Ruiz, Elizabeth Natalia Martínez Núñez, Jesús Adolfo Monasterios Montero,
Maria Jose Cotua Acosta, Daniel Enrique Anato Moreno 893
- Effects of positive self-talk on the level of anxiety of patients with coronary heart disease
Uswatun Hasanah, Dede Nasrullah, Rifma Yuniar, Nugroho Ari Wibowo, Reliani 906

REVIEW ARTICLES	
Tropical diseases and pregnancy: Narrative review Ana Carvajal, Carlos Cabrera Lozada, Jeiv Gomez	913
Non-invasive neuromonitoring devices for the detection of intraoperative cerebral ischemia: A scoping review Esteban García-Espitia, Juan Pablo Pacheco-Alba, Eduardo Tuta-Quintero, Daniel Botero-Rosas	933
Medicines in the environment: A public health problem. Sistematic Review Bartolome Manzolillo	942
CASE REPORTS	
Frosted branch angiitis in a human immunodeficiency virus-positive patient with cytomegalovirus retinitis Noel Roig-Marin, Pablo Roig-Rico, Vicente Navarro-Ibanez	951
Vulvar leiomyosarcoma. An unusual pathology Franco Calderaro Di Ruggiero, Viviana Garcia, Johatson Freytes, Carmen Silva, Lorenzo J Calderaro Fernandez	956
Uterine lipoleiomioma. Case report Viviana Garcia, Franco Calderaro Di Ruggiero, Johatson Freytes	962
PERSPECTIVES	
How do pandemics end? The future of COVID -19 Jose Esperza, Gilberto Vizcaino	968
HISTORY AND PHILOSOPHY OF MEDICINE	
Management of “backache”, a medical prescription in the New Kingdom of Granada Eduardo Tuta-Quintero, Julio Cesar Martinez-Lozano, Ignacio Briceno-Balcazar, Diego Sierra-Barbosa, Alberto Gomez-Gutierrez	975
Serpentes, Insecta et Rabiem: management of snake bites, a medical emergency during the colonial period Jhonnatan Coronado-Sarmiento, Ignacio Briceno-Balcaza, Julio Cesar Martinez-Lozano, Eduardo Tuta-Quintero	981
BIOGRAPHIES	
Semblance of the Promotion of Physicians 1962 of the Universidad Central de Venezuela Leopoldo Briceno-Iragorry	989
Promotion of surgeons “Bicentennial Promotion of Medical Studies in Venezuela, Dr. Lorenzo Campins and Ballester”. Some annotations Claudia Blandenier de Suarez, Luis Herrera Garcia, Rosa Marcano de Cuenca, Jose Luis Cevallos, Cesar Blanco Rengel, Leopoldo Moreno Brandt	996
VARIOUS	
The Gaceta Medica de Caracas 100, 50, 25 years ago. Enrique Santiago Lopez-Loyo	1012
Life of the Academy, Summary of the papers presented and the Bibliographic Notes Enrique Santiago Lopez-Loyo	1017
Index Volume 129. Dr. Rogelio Perez D'Gregorio	1036

Academia Nacional de Medicina

Junta Directiva

2020-2022

Presidente: Dr. Enrique Santiago López Loyo

Vicepresidente: Dra. Isis Nézer de Landaeta

Secretario: Dr. Huníades Urbina-Medina

Tesorero: Dra. Lilia Cruz de Montbrun

Bibliotecario: Dr. Guillermo Colmenares Arreaza

Individuos de Número

Sillón I:	Dra. Aixa Müller de Soyano	Sillón XXI:	Dra. Claudia Blandenier de Suárez
Sillón II	Dr. Miguel José Saade Aure	Sillón XXII:	Dr. Huníades Urbina-Medina
Sillón III	Dr. Marco Sorgi Venturoni	Sillón XXIII:	Dr. José A. O'Daly Carbonell
Sillón IV:	Dr. Rafael Muci Mendoza	Sillón XXIV:	Vacante
Sillón V:	Dr. Miguel González Guerra	Sillón XXV:	Dr. Felipe Martín Piñate
Sillón VI:	Dr. Saúl Krivoy	Sillón XXVI:	Dr. Claudio Aoün Soulie
Sillón VII:	Dr. Guillermo Colmenares A	Sillón XXVII:	Dr. Mauricio Goihman Yahr
Sillón VIII:	Dr. Leopoldo Briceño-Iragorry	Sillón XXVIII:	Dr. Luis Ceballos García
Sillón IX:	Dr. Otto Rodríguez Armas	Sillón XXIX:	Dr. Julio Borges Iturriza
Sillón X:	Dr. José Ramón Poleo	Sillón XXX:	Dr. Félix José Amarista
Sillón XI:	Dr. Saúl Kizer Yorniski	Sillón XXXI:	Dr. Enrique López Loyo
Sillón XII:	Vacante	Sillón XXXII:	Dra. Ofelia Uzcátegui U.
Sillón XIII:	Dr. José Francisco	Sillón XXXIII:	Dr. Nicolas Bianco Colmenares
Sillón XIV:	Dr. Oscar Beaujón Rubín	Sillón XXXIV:	Dr. Otto Lima Gómez
Sillón XV:	Dr. Víctor Ruesta	Sillón XXXV:	Dr. Ítalo Marsiglia
Sillón XVI:	Dr. Harry Acquatella M	Sillón XXXVI:	Dr. Antonio Clemente H
Sillón XVII:	Dra. Isis Nézer de Landaeta	Sillón XXXVII:	Dr. Juan Antonio Yabur Tarrazzi
Sillón XVIII:	Dr. José M Guevara Iribarren	Sillón XXXVIII:	Dr. Rafael Apitz Castro
Sillón XIX:	Dra. Lilia Cruz de Montbrun	Sillón XXXIX:	Dra. Doris Perdomo de Ponce
Sillón XX:	Dr. Pedro Faneite Antigue	Sillón XL:	Dr. Horacio Vanegas

Miembros Correspondientes Nacionales

1. Dra. Eddy Verónica Mora (Carabobo)
2. Dr José Alberto Briceño Polacre (Trujillo)
3. Dr. Jorge García Tamayo (Zulia)*
4. Dr. José Luis Cevallos (Caracas)
5. Dr. Israel Montes de Oca (Caracas)
6. Dr. Carlos Rojas Malpica (Carabobo)
7. Dra. Laura C Vásquez de Ricciardi (Trujillo)
8. Dr. Jesús Enrique González Alfonso (Caracas)
9. Dr. Oswaldo Guerra Zagarzazu (Carabobo)
10. Vacante
11. Dr. José Alejandro Corado Ramírez (Carabobo)
12. Dra. Evelyn Figueroa de Sánchez (Carabobo)
13. Dr. Sergio Osorio Morales (Zulia)
14. Dr. Rafael María Rosales Acero (Táchira)
15. Dra. Myriam del Valle Marcano Torres (Carabobo)
16. Dr. Carlos Cabrera Lozada (Caracas)
17. Dr. Wilmar de Jesús Briceño Rondón (Barinas)
18. Dra. Emely Zoraida Karam Aguilar (Caracas)
19. Dr. Jesús Eduardo Meza Benítez (Carabobo)
20. Dra. Elsa Báez de Borges (Caracas)
21. Dr. Jesús Alfonso Osuna Ceballos (Mérida)
22. Dr. Felipe Díaz (Zulia)*
23. Dra. Nelly Petit (Zulia)*
24. Vacante
25. Dr. Alberto Paniz-Mondolfi (Lara)
26. Vacante
27. Dr. Raúl Díaz Castañeda (Valera)
28. Dr. Mariano Álvarez Álvarez (Maturín)
29. Dr. José Rodríguez Casas (Caracas)
30. Vacante
31. Dr. Nelson Urdaneta (Caracas)
32. Dr. Gastón Silva Cacavale (Caracas)
33. Dr. Eduardo Morales Briceño (Caracas)
34. Dra. Laddy Casanova de Escalona (Carabobo)
35. Dr. José Ramón Guzmán (Zulia)
36. Dra. Mercedes López de Blanco (Caracas)
37. Dr. José T. Nuñez Troconis (Zulia)
38. Dra. Enriqueta Sileo Giuseffi (Caracas)
39. Dr. Marino José González Reyes (Caracas)
40. Dr. Luzardo Canache Campos (Aragua)
41. Dr. Franco Calderaro Di Ruggiero (Caracas)
42. Vacante
43. Dr. José Manuel De Abreu D'Monte (Caracas)
44. Dr. José Andrés Octavio Seijas (Caracas)
45. Dr. Antonio De Santolo (Caracas)
46. Vacante
47. Dr. Andrés Soyano López (Caracas)
48. Dra. Janice Fernández de D'Pool (Zulia)
49. Dra. Rosa Cedeño de Rincón (Zulia)
50. Dr. Raúl Fachin Viso (Carabobo)

Miembros Correspondientes Extranjeros

1. Dr. Vladimir Hachinsky (Canadá)
2. Dr. Remigio Vela Navarrete (España)
3. Dr. Zoilo Cuellar Montoya (Colombia)
4. Dr. Alvaro Rodríguez González (Colombia)
5. Dr. Pedro Grases (Costa Rica)
6. Dr. Igor Palacios (Estados Unidos)
7. Dr. Otto Gago (Estados Unidos)
8. Dr. Francisco López Muñoz (España)
9. Dr. Eduardo Pretell Zárate (Perú)
10. Dr. Harold Zur Hausen (Alemania)
11. Dr. Henry Lynch (Estados Unidos)
12. Dr. Vicente Gutiérrez Maxwell (Argentina)
13. Dr. J. Aurelio Usón Calvo (España)
14. Dr. José Augusto Da Silva Messias (Brasil)
15. Dr. Gianfranco Parati (Italia)*
16. Dr. Juan del Rey Calero (España)
17. Dr. Jean Civatte (Francia)
18. Dra. Carmen Luisa García de Insausti (España)
19. Dr. Andrew V. Schally (Estados Unidos)
20. Dr. Terence J Ryan (Inglaterra)
21. Dr. Jean Pierre Delmont (Francia)
22. Vacante
23. Dr. Jörg G.D. Bikmayer (Austria)
24. Dr. John Uribe M. (Estados Unidos)
25. Dr. José Esparza (Estados Unidos)
26. Dr. Augusto Bonilla Barco (Ecuador)
27. Dr. Kenneth Kenyon (Estados Unidos)
28. Dr. Gabriel Carrasquilla (Colombia)*
29. Dr. Janis V. Klavins (Estados Unidos)
30. Vacante

Invitados de Cortesía

(Dado su carácter todavía no son Académicos)

Dra. Belkysyolé Alarcón de Noya
Dr. Rafael Arteaga Romero
Dra. Elvia Irene Badell Madrid
Dr. César Blanco Rengel
Dra. Susana Blanco Sobrino
Dra. Alba Cardozo
Dr. Antonio Cartolano
Dr. Pedro Ignacio Carvallo
Dr. Jaime Díaz Bolaños
Dra. Maritza Durán
Dr. Antonio A Eblen Zaijur
Dr. Mariano Fernández S.
Dr. Cutberto Guarapo Rodríguez
Dr. Peter Gunczler
Dra. Marienalla Herrera Cuenca

Dr. David Lobo
Dr. Aderito De Sousa
Dra. Ana María Martínez
Dra. María Eugenia Mondolfi Gudat
Dr. José Félix Oletta López
Dr. Saúl Peña Arciniegas
Dr. Rafael Rangel Aldao
Dr. Omar Reyes Morales
Dr. Francisco Alejandro Rískquez
Dr. Jesus Rodríguez Ramírez
Dr. Germán Rojas Loyola
Dr. Rafael Romero Reverón
Dr. Tomás José Sanabria Borjas
Dr. Daniel Sánchez
Dr. Herbert Stegemann
Dra. Marfa Yanes Herrera

Comité Editorial de la Gaceta Médica de Caracas (Editorial Board)

Editor en jefe (Editor in Chief)

Dr. Enrique Santiago López Loyo (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Editor Gerente (Senior Editor)

Dra. Anita Stern de Israel (UCV, Venezuela)

Editores Honorarios (Honorary Editors in Chief)

Dr. Antonio Clemente (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Rafael Muci Mendoza (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dra. Doris Perdomo de Ponce (ACADEMIA NACIONAL DE MEDICINA, México)

Editores Asociados (Associate Editors)

Dr. Harry Acquatella (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dra. Lissé Chiquinquirá Angarita Dávila (Universidad Andres Bello: Talcahuano, Concepción, Biobio, CL Chile)

Dr. Claudio Aoún (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Gustavo Aroca (Universidad Simón Bolívar, Colombia)

Dr. Leopoldo Briceño-Iragorry (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Franco Calderaro di Ruggiero (ACADEMIA NACIONAL DE MEDICINA, Clínica Razetti, Venezuela)

Dra. Lilia Cruz (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Jorge Escobedo (UNIVERSIDAD AUTÓNOMA DE MÉXICO, México)

Dr. Mariano Fernández (UCV, Venezuela)

Dr. José Francisco (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. José María Guevara (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Saúl Kízer (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Saúl Krivoy (ACADEMIA NACIONAL DE MEDICINA, EE. UU.)

Dr. Felipe Martín Piñate (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. José Parra (Universidad de Guadalajara, México)

Dra. Diana Marcela Rojas (Universidad Andrés Bello: Talcahuano, Concepción, Biobio, CL, Chile)

Dr. Tomás Sanabria (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dra. Enriqueta Sileo (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Gastón Silva (ACADEMIA NACIONAL DE MEDICINA y Policlínica Metropolitana, Caracas, Venezuela)

Dr. Marco Sorgi (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Andrés Soyano (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Huniades Urbina (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dra. Ofelia Uzcátegui (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Horacio Vanegas (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Dr. Juan Yabur (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Comité Editorial de la Gaceta Médica de Caracas (Editorial Board) Continuación

Editores Ejecutivos (Executive Guest Editors)

- Dr. Luis Alcocer (Universidad Autónoma de México, México)
Dr. Ezequiel Bellorin Font (Universidad de Cleveland, EE. UU)
Dr. Raúl Carlini (Hospital Universitario, Caracas, Venezuela)
Dr. Guillermo Colmenares (ACADEMIA NACIONAL DE MEDICINA, Venezuela)
Dr. Manuel Guzmán Blanco (Centro Médico de Caracas, Venezuela)
Dr. Zafar Israeli (Universidad de Emory, EE. UU)
Dra. Isis Nézer de Landaeta (ACADEMIA NACIONAL DE MEDICINA, Venezuela)
Dra. Alicia Ponte Sucre (IME-UCV, Venezuela)
Dr. Heberto Suarez Roca (LUZ, EE. UU)
Dr. Herbert Stegeman (ACADEMIA NACIONAL DE MEDICINA, Venezuela)

Editores de Sección (Section Editors)

- Dr. Oscar Aldrey (Instituto Médico, La Floresta, Venezuela)
Dr. Valmore Bermúdez (USB, Colombia)
Dra. Claudia Blandenier de Suárez (ACADEMIA NACIONAL DE MEDICINA, Venezuela)
Dr. Claudio Borghi (Universidad de Bologna, Italia)
Dr. Juan De Sanctis (Universidad de Olomouc, República Checa)
Dr. José Esparza (ACADEMIA NACIONAL DE MEDICINA, EE.UU)
Dr. Luis Juncos (J Robert Cade Foundation, Argentina)
Dr. Carlos Ferrario (Universidad de Carolina del Norte, EE. UU)
Dr. Claudio Ferri (Università degli Studi dell'Aquila, Italia)
Dr. Patricio López Jaramillo (Universidad Autónoma de Bucaramanga (UNAB), Colombia)
Dr. Héctor Marcano (Hospital Universitario de Caracas, Venezuela)
Dr. Oscar Noya (Instituto de Medicina Tropical, UCV, Venezuela)
Dr. José Andrés Octavio (Hospital de Clínicas Caracas, Venezuela)
Dr. José A O'Daly (ACADEMIA NACIONAL DE MEDICINA, Venezuela)
Dr. Stefano Omboni (Italian Institute of Telemedicine, Italia)
Dr. Gianfranco Parati (University of Milano-Bicocca, Italia)
Dr. José Ramón Poleo (ACADEMIA NACIONAL DE MEDICINA, Venezuela)
Dr. Bernardo Rodríguez Iturbe (Instituto Nacional Ignacio Chávez, México)
Dr. Félix Tapia (Instituto de Biomedicina-UCV, Venezuela)

Editores Asistentes (Assistant Editors)

- Dr. Henry Collet Camarillo (Clínica Ávila, Venezuela)
Dr. Freddy Contreras (UCV, Venezuela)
Dr. Giuseppe Crippa (Unidad de Hipertensión Arterial Piacenza, Italia)
Dra. Maricarmen Chacín (USB, Colombia)
Dra. María Sofía Martínez Cruz (Universidad de Virginia, EE. UU)
Dra. Dolores Moreno (Instituto de Biología Molecular -UCV, Venezuela)
Dr. Alexis García (Instituto de Inmunología, UCV, Venezuela)
Dra. Jenny Garmendia (Instituto de Biología Molecular-UCV, Venezuela)
Dr. Edward Rojas (Universidad de Virginia, EE. UU)
Dr. Juan Salazar (La Universidad del Zulia, Venezuela)
Dr. Francisco Tortoledo (España)

Stimulus from the activator agent *Solanum melongena L.* on sperm quality through the lipid profile of *Rattus norvegicus* with hyperlipid induction

Estímulo del agente activador *Solanum melongena L.* sobre la calidad del esperma a través del perfil lipídico de *Rattus norvegicus* con inducción hiperlipídica

Rinza Rahmawati Samsudin^{1a*}, Nur Vita Purwaningsih^{2b}, Rahma Widyastuti^{3b}

SUMMARY

Introduction: A high-fat diet and obesity, promoted by an unhealthy lifestyle, affect the structure of spermatozoa. The study aims to determine the effect of using purple eggplant on improving lipid profile and sperm quality as an alternative to traditional medicine of *Rattus norvegicus*. **Methods:** This type of research is a laboratory experiment using a post-test with a control group design. Twenty-eight *Rattus norvegicus* were divided into four groups. The number of spermatozoa was calculated by sucking the sperm stock solution using a hemocytometer suction. **Results:** The high cholesterol and distilled water (G1), high

cholesterol and purple eggplant 100 mg/kg BW (G2), and high cholesterol and purple eggplant 200 mg/kg BW (G3) groups had higher body weight compared to the standard feed and distilled water (G0) group. The G1 group had the highest body weight (136.25 ± 0.901 grams) and cholesterol levels (168.13 ± 1.797 mg/dL). Moreover, G3 group (159.38 ± 1.253 mg/dL) had lower cholesterol levels compared to G2 group (152.25 ± 10.31 mg/dL). The G3 group ($35.00 \times 10^6 \pm 0.500$) had the highest mean sperm count, followed by G1, G2, and G0 groups ($33.25 \times 10^6 \pm 0.675$, $32.75 \times 10^6 \pm 0.366$, $31.38 \times 10^6 \pm 0.680$, respectively). **Conclusion:** The purple eggplant extract was able to reduce cholesterol levels and increase sperm count.

Keywords: Purple eggplant, body weight, cholesterol levels, sperm count

DOI: <https://doi.org/10.47307/GMC.2021.129.4.8>

ORCID ID: 0000-0002-3183-4644¹

ORCID ID: 0000-0003-0828-7862²

ORCID ID: 0000-0002-3784-1466³

^aDivision of Biochemistry, Department of Medical Laboratory Technology, Faculty of Health Science, Muhammadiyah University of Surabaya, Surabaya 60113, Indonesia

^bDivision of Clinical Chemistry, Department of Medical Laboratory Technology, Faculty of Health Science, Muhammadiyah University of Surabaya, Surabaya 60113, Indonesia

Recibido: 18 de noviembre 2021

Aceptado: 1 de diciembre 2021

Corresponding Author: Rinza Rahmawati Samsudin, M.Sc.
Division of Biochemistry Department of Medical Laboratory Technology, Faculty of Health Science, Muhammadiyah University of Surabaya Indonesia
Sutorejo Street No. 59 Surabaya 60113, East Java, Indonesia
Tel: +628819000911; Fax: +6231-3811967
E-mail: rinzadiyanto@gmail.com

RESUMEN

Introducción: Una dieta alta en grasas y la obesidad, promovida por un estilo de vida poco saludable, afectan la estructura de los espermatozoides. El estudio tiene como objetivo determinar el efecto del uso de berenjena morada en la mejora del perfil lipídico y la calidad del esperma como alternativa a la medicina tradicional de *Rattus norvegicus*. **Métodos:** Este tipo de investigación es un experimento de laboratorio que utiliza pos-prueba con un diseño de grupo de control. Se dividieron veintiocho *Rattus norvegicus* en cuatro grupos. El número de espermatozoides se calculó succionando la solución madre de esperma usando una succión de hemocitómetro. **Resultados:** Los grupos fueron, colesterol alto y agua destilada (G1), colesterol alto y berenjena morada 100 mg/kg de peso corporal (G2), colesterol alto y berenjena morada de 200 mg/kg de peso corporal (G3), los cuales presentaron un mayor peso corporal en comparación con el alimento estándar y los grupos con agua destilada (G0). El grupo G1 tuvo el mayor peso corporal ($136,25 \pm 0,901$ gramos) y niveles de colesterol ($168,13 \pm 1,797$ mg/dL). Además, el grupo G3 ($159,38 \pm 1,253$ mg/dL) tuvo niveles de colesterol más bajos en comparación con el grupo G2 ($152,25 \pm 10,31$ mg/dL). El grupo G3 ($35,00 \times 10^6 \pm 0,500$) tuvo el recuento medio de espermatozoides más alto, seguido de los grupos G1, G2 y G0 ($33,25 \times 10^6 \pm 0,675$, $32,75 \times 10^6 \pm 0,366$, $31,38 \times 10^6 \pm 0,680$, respectivamente). **Conclusión:** El extracto de berenjena morada pudo reducir los niveles de colesterol y aumentar el recuento de espermatozoides.

Palabras clave: Berenjena morada, peso corporal, niveles de colesterol, recuento de espermatozoides.

INTRODUCTION

A high-fat diet and obesity, which are encouraged by an unhealthy lifestyle, affect the structure of spermatozoa and the development and health of offspring. Indeed, infertile men have been observed to have improper dietary patterns, including meal omissions, insufficient antioxidant intake, and a high energy density (1). An unhealthy hypercaloric diet, high saturated fat and trans fat intake, a high glycemic index, and low nutritional density may all be associated with increased oxidative stress, the underlying cause of obesity, intestinal dysbiosis, type 2 diabetes, and insulin resistance (2). The metabolic disorders mentioned above are associated with a decline

in fertility, primarily due to the generation of oxidative stress, which is regarded as a significant factor contributing to decreased sperm quality and an increased risk of infertility and hormonal and immunological disorders (3).

One of the leading causes of disruption of the process of spermatogenesis in the testes is the presence of free radicals (4). Free radicals can also inhibit the process of spermatogenesis because Leydig cells are disrupted to reduce levels of testosterone hormone secretion. The negative impact of free radicals on reproductive health can be overcome by giving antioxidants (5,6). Additionally, while sperm produces reactive oxygen species (ROS), an imbalance between ROS and antioxidants will harm sperm associated with male infertility, peroxy radicals, and peroxynitrite (7).

Food is the primary source of cholesterol. Dietary cholesterol is initially transported from the small intestine to the liver, where it is then redistributed to the organs that require it. The Scavenger receptor B type 1 (SR-B1) is a high-density lipoprotein (HDL) receptor on the cell surface that mediates the uptake of HDL-cholesterol ester (HDL-CE). Additionally, less than half of cholesterol is synthesized de novo. The synthesis of cholesterol is a multi-step enzymatic process. The synthesis of cholesterol begins with the transport of acetyl-CoA from the mitochondria to the cytosol. Hydroxy-methyl-glutaryl-CoA (HMG-CoA) is formed through a series of reactions and is then converted to mevalonate by HMG-CoA reductase. Finally, a series of steps will result in the formation of cholesterol (8,9). Cholesterol is a critical component of mammalian plasma membranes, where it is required for the proper permeability and fluidity of the membrane. Hypercholesterolemia is a risk factor for noninsulin-dependent diabetes, osteoarthritis, some types of cancer, and certain reproductive and metabolic disorders. Hypercholesterolemia is a lipoprotein metabolic disorder characterized by high serum low-density lipoprotein and blood cholesterol. Cholesterol is a steroid lipid found in the cell membranes and transported in the blood plasma of all animals (10).

Many studies have been conducted to identify natural substances that can protect the body from oxidative stress, such as the use of antioxidant compounds found in fruits, vegetables, and

grain plants (11-13). Antioxidants are needed to protect the body from free radical attack through protecting enzymes that repair DNA damage, therefore increasing our body's ability to regenerate itself (14). One of them is purple eggplant (*Solanum melongena L.*), one of Indonesia's medicinal plants. Purple eggplant contains alkaloid compounds in the form of glycosides, namely solanine, tomatin, and solasodine. Solasodine is a steroidal glycoalkaloid compound contained in purple eggplant and is thought to have an antifertility effect(15). The administration of purple eggplant ethanol extract could significantly reduce the percentage of rat spermatozoa motility compared to controls. Solasodine inhibits the expression of luteinizing hormone (LH) and spermatogenesis in mice (16) having high antioxidant content. Purple eggplant contains anthocyanin pigments which act as antioxidants. Anthocyanins are part of phenolic compounds, which are classified as flavonoids. Sadivola stated that the dominant anthocyanin content in purple eggplant skin is delphinidin 3-rutinoside, a purple pigment. Anthocyanin pigments are more stable under acidic conditions than basic and neutral conditions and are unstable with hot light and certain metals. Microencapsulation is one way to increase the stability of an anthocyanin compound (delphinidin 3-rutinoside)(17). This study aimed to determine the effect of using purple eggplant on the improvement of lipid profile and sperm quality as an alternative to traditional medicine.

METHODS

Research Type and Design

This study was a laboratory experiment using a post-test research design with a control group (*post-test control group design*). This study used experimental animals in the form of white male rats (*Rattus norvegicus*) Wistar strain. A total of 28 *Rattus norvegicus* were divided into four groups, including G0, G1, G2, and G3. G0 was only given standard feed and distilled water. G1 was only fed high cholesterol and distilled water diet. G2 was fed a high cholesterol diet and given purple eggplant with 100 mg/kg BW. G3 was fed a high-cholesterol diet with purple eggplant treatment with 200 mg/kg BW. This study was

carried out for 4 weeks.

Test Animals

The test animals selected were *Rattus norvegicus* Wistar under the inclusion criteria in the form of *Rattus norvegicus* Wistar strain, 3-4 months old, bodyweight of 150-200 g, healthy condition and no macro anatomical abnormalities, and never get any treatment. Exclusion criteria were sick rats and dead rats during the study.

Procedure for Making Purple Eggplant Extract

Fresh purple eggplant was dried in an oven at 70 °C for 24 hours and then mashed with a blender, macerated with 96 % ethanol until the macerated was clear. The results of the macerate are put into a vacuum rotatory evaporator until a thickened extract is obtained, then proceed with making a suspension at a dose of 100 mg/kg, BW and 200 mg/kg, BW, diluted by adding distilled water.

Procedure for Making High-Fat Feed

In this study, high-fat feeding was made from quail egg yolks, referring to the previous study. The yolk of the quail is separated from the white of the egg. Giving quail egg yolk to rats by induced using a gastric probe as much as 1.5 mL/head was carried out every morning for 4 weeks except for the negative control rat group, which was not induced by quail egg yolk.

Blood Sampling Procedure

The blood samples of experimental animals were taken from the hearts of rats. Before sampling, the experimental animals were fasted at night (12 hours fast). During fasting, the experimental animals were given water *ad libitum*. Animal sacrifices were made in the morning, starting with physical euthanasia techniques. To draw blood can be done directly by inserting a syringe directly into the heart and aspirating slowly.

Sperm Sampling Procedure

Spermatozoa collection of male rats begins with animal sacrifice, which is carried out in the morning, begins with physical euthanasia techniques, and then dissects their reproductive organs. The cauda epididymis was then removed and placed in a petri dish containing 0.9% sodium chloride. Additionally, the cauda epididymis was inserted into a watch glass containing 1 ml of 0.9 percent NaCl, and the proximal part of the cauda was cut slightly with scissors and the cauda was pressed slowly until secretory fluid emerged and was suspended in 0.9 percent NaCl. The suspension of spermatozoa from the cauda epididymis obtained can be used for observations, including motility, number, and morphology of spermatozoa.

Sperm Count Procedure

To determine the number of spermatozoa, the following method was used. The number of spermatozoa was calculated by sucking the sperm stock solution using a hemocytometer suction pipette to the 0.5 mark, then the physiological NaCl solution was sucked up to the 101 marks, and the pipette was shaken. Discard a few drops on tissue paper, then put them in a counting chamber which has been closed with a coverslip and has been prepared in a microscope, then examined under a microscope. Calculated using the formula for the number of spermatozoa counted (s) x dilution x 1 ml NaCl = s x 20,000 = million/mm³.

Sperm Motility Calculation

The rat sperm was taken from the *cauda epididymis* by slashing and pressing gently. One drop of sperm was placed on an object glass, plus one drop of 0.9 % NaCl physiological solution, mixed evenly and covered with an object glass. The percentage of motile spermatozoa was calculated in one field of view using a light microscope at a magnification of 100 times by estimating progressively moving spermatozoa from the entire field of view and the estimated area, then multiplied by 100 %. Assessment is done by calculating the percentage of spermatozoa whose movement is progressively

forward compared to those observed (moving and immobile).

Data Processing and Analysis

Data obtained from observations on sperm count, body weight, and cholesterol levels were previously carried out. From the dependent and independent variables, normality and homogeneity were carried out with the Shapiro-Wilk test because the number of samples was <50; if the data distribution was normal, then the average was homogeneous with the one-way ANOVA test. If the result means $p < 0.005$, the results were continued with the LSD test using Post Hock. If the data were not normal and not homogeneous, it would be tested with the Kruskal Walis test, followed by the Mann-Whitney Test. The results of data processing are displayed in the form of tables and graphs.

RESULTS

During the study, rats were weighed every week for 4 weeks. The weighing was done using a digital scale. The results of rat body weight, cholesterol level, and sperm count in each group are summarized in Table 1. The body weight, cholesterol level, and sperm count of rats were normally distributed ($p > 0.05$).

The results showed that the high-fat diet was successful in increasing the body weight of rats. The bodyweight of the G1, G2, and G3 groups was higher than the G0 group, which was only given standard feed. The G1 group showed the highest increase in body weight. Whilst G0 and G1 groups were the lowest and the highest cholesterol levels, respectively. Moreover, the G3 group had lower cholesterol levels compared to the G2 group. In addition, the G3 group had the highest mean sperm count, followed by G1, G2, and G0 groups (Table 1).

The microscopic observations, including sperm count, viability, and morphology, in male *Rattus norvegicus* Wistar strain with an objective lens magnification of 100x, had different results in each treatment. On sperm viability examination (Figure 1), G0 showed the sperm still alive (the

Table 1

The results of rat body weight, cholesterol level, and sperm count (N=28)

Group	Weight (Grams)*	Normality Test	Cholesterol level (mg/dL)*	Normality Test	Sperm Count*#	Normality Test
G0	131.38±0.822	0.945	129.38±2.719	0.210	31.38±0.680	0.603
G1	136.25±0.901	0.976	168.13±1.797	0.446	33.25±0.675	0.351
G2	130.13±0.675	0.626	159.38±1.253	0.959	32.75±0.366	0.408
G3	129.88±0.766	0.720	152.25±10.31	0.454	35.00±0.500	0.273

*Mean±SD

#Value multiplied by 106 mL

G0: Group was only fed standard feed and aquades

G1: Group was only fed high cholesterol and aquades

G2: Group was fed a diet high cholesterol and purple eggplant 100mg/kg, BW

G3: Group was fed a diet high cholesterol and purple eggplant 200mg/kg, BW

yellow arrow) because the sperm did not absorb the red color in the eosin negrosin reagent. At the same time, the red arrows indicate sperm that died because they absorbed the red color of the eosin-negrosin reagent. G1 to G3 only showed dead sperm.




Figure 1. Examination of sperm viability with an objective lens magnification of 100x.

DISCUSSION

This study demonstrated that giving a high-fat diet for 4 weeks could increase the body weight of rats and cholesterol levels compared to the control group. The administration of purple




Figure 2. Sperm morphology examination with an objective lens magnification of 100x.

Figure 2 shows the results of the sperm morphology examination. The normal sperm shapes from head to tail (yellow arrows) were shown in G0 and G1. G2 had the sperm in normal and abnormal forms, which only the sperm body (red arrow). While G3 had the sperm with only body shape (red arrow).

eggplant at 200 mg/kg BW was able to reduce cholesterol levels and increase the sperm count. Several diseases have been treated with eggplant, including asthma, bronchitis, diabetes, rheumatoid arthritis, and hypercholesterolemia (18). The clinical use of eggplant is due to its phenolic and alkaloid. The primary phenolic compounds in the skin and pulp of eggplant are delphinidin (an anthocyanin) and chlorogenic acid (a phenolic acid) (19). Moreover, delphinidin has exhibited inhibitory properties against the α -amylase enzyme, and thereby it may be useful in the treatment of diabetes and its complications such as overweight and obesity, and cardiovascular disease (20). It has been suggested that chlorogenic acids exhibit anti-obesity and anti-hyperlipidemic activities by alleviating the levels of free fatty acids and triglycerides (TG) (21). Additionally, several studies have reported the pharmacologic aspects of eggplant, such as antioxidant (22), obesity (23), hyperlipidemia (18). Hyperlipidemia is characterized by abnormally high levels of lipids in the blood, including fat, cholesterol, and TGs. It is widely recognized as the primary risk factor for cardiovascular disease and is one of the leading causes of death worldwide.

A previous study also showed increased high cholesterol levels after getting feed with a high fat composition (24). Eggs and egg products account for 25 % of daily total cholesterol intake in children and adults in the United States (25). Some studies related to man have shown the association of BMI with reproductive parameters like poor semen quality (26), decreased sperm concentration (27), decreased number of normal motile sperm cells (28), increased ROS, and increased DNA fragmentation index (29).

Male obesity has a less well-documented effect on reproduction than female obesity does. Several studies, however, indicate that overweight and obese men have lower sperm quality and fertility. Male obesity is suspected to alter sperm parameters, particularly sperm concentration, total progressively motile sperm count (29), total sperm count, total motile sperm count sperm morphology, and DNA fragmentation (30). However, the magnitude of this increase was small and had only a minor clinical significance, as there was no correlation between increased ROS production and decreased sperm DNA

integrity or motility. Additionally, increased BMI was found to be significantly associated with a decrease in sperm concentration, serum testosterone, and serum estradiol (31). In obese men, excessive ROS production and abnormal hormonal regulation result in suboptimal sperm quality. It is hypothesized that these patients' oxidative stress is caused by dysregulation of adipocytokine and ROS generation (32). The excessive production of ROS may be a result of obese men's increased metabolic rates and maintenance of hemostasis. Additionally, elevated levels of ROS and temperature in the testicles may denature enzymes involved in spermatogenesis. Increased scrotal skin temperature was associated with a decrease in sperm concentration (27).

The decrease in spermatozoa viability in the purple eggplant extract treatment group was influenced by the presence of alkaloid compounds in purple eggplant, namely solasodine which can interfere with the permeability of the spermatozoa membrane. The decrease in sperm viability is caused by disruption of the permeability of the spermatozoa membrane which can later interfere with the transportation of nutrients needed for the movement of spermatozoa (33), besides that the permeability of the spermatozoa membrane is closely related to cell metabolism and plays a role in the formation of energy so that in this case the permeability of the spermatozoa membrane is closely related to the motility and viability of spermatozoa.

The relationship between lipids and sperm production is complex and unknown in its etiology. In humans, the cholesterol content of sperm varies significantly, even between ejaculates (34). In the meantime, the amount of cholesterol in sperm membranes is directly proportional to the sperm morphology (35) and fertility potential (34). For several mammalian species, the lipid composition of the sperm plasma membrane has been determined (36). In comparison to other cell types, spermatozoa have a unique lipid composition. Spermatozoa contain a greater proportion of neutral lipids, particularly a high concentration of diacylglycerol (DAG) (37,38). Recently, the lipid component (O-acyl)- ω -hydroxy-fatty acids (OAHFA) with a carbon chain length of up to 52 was identified in spermatozoa for the first time. It was found in

the head of sperm rather than the tail region (38).

However, few studies are examining the detailed relationships between serum cholesterol or other lipid profiles and sperm quality. This cross-sectional study, like our previous ones, had some limitations (39,40). These included the lack of semen volume and total sperm count data in the initially designed protocol of the program, absence of detailed records of endocrine levels and lipid contents of semen, and lack of abstinence duration records for evaluation in the study. Moreover, the lipid-related pathological conditions, such as familial hypercholesterolemia or ApoE genotype, were not explored in our study because of the initially designed data requirement. The semen samples of our study were collected using home-collection kits; the quality may not be equal to that resulting from the on-site collection. Further studies, including those assessing the lipid content of semen and performing measurement of lipid-related nuclear receptors such as liver X receptors (LXRs) (41), peroxisome proliferator-activated receptors (PPARs), small heterodimer partner (SHP) (42), 27, and retinoid X receptors (RXRs) (43), should be designed further to explore the relationships between lipids and semen quality (34).

Reduced motility, intracellular enzyme damage, and damage to the structure of the plasma membrane are all characteristics of spermatozoa damaged by lipid peroxidation (44). Additionally, as spermatozoa mature in the epididymis, the composition of the components that comprise the spermatozoa plasma membrane changes. The plasma membrane of the spermatozoa will lose some cholesterol, increasing the ratio of unsaturated fatty acids to cholesterol (45).

The increase in abnormalities in the treatment group was caused by the purple eggplant extract containing tannins. Tannins function to bind proteins and ions contained in spermatozoa membranes so that the tyrosine enzyme and phosphorylation process in the spermatozoa membranes are disrupted and ultimately cause morphological abnormalities of spermatozoa (46). The increase in morphological abnormalities of spermatozoa can be caused by damage in the seminiferous tubules and when the spermatozoa leave the seminiferous tubules and during their journey through the epididymis. The male

reproductive system in normal conditions maintains a balance between ROS production and antioxidant activity. However, excessive ROS production in sperm or seminal plasma can impair the antioxidant defense mechanisms of the sperm or seminal plasma, resulting in oxidative stress. It is well established that oxidative stress damages sperm chromatin/DNA. Antioxidant therapy is widely accepted to improve sperm quality and male fertility by reducing oxidative stress. The findings of this study using purple eggplant corroborate those of Sabeti et al. (47).

CONCLUSION

Giving a high-fat diet for 4 weeks was able to increase the bodyweight of rats and cholesterol levels compared to the control group. At the same time, the administration of purple eggplant at 200 mg/kg, BW could reduce cholesterol levels and increase the sperm count.

REFERENCES

1. Giahi L, Mohammadmoradi S, Javidan A, Sadeghi MR. Nutritional modifications in male infertility: A systematic review covering 2 decades. *Nutr Rev.* 2016;74(2):118-130.
2. Varani J. Healthful Eating, the Western Style Diet and Chronic Disease. *Approaches Poultry, Dairy Vet Sci.* 2017;1(3):44-46.
3. Walczak-Jedrzejowska R, Wolski JK, Slowikowska-Hilczer J. The role of oxidative stress and antioxidants in male fertility. *Cent Eur J Urol.* 2013;66(1):60-67.
4. NingrumAG,Ratnawati R,Andarini S. Effect of sweet potato anthocyanin (*Ipomoea batatas* L.) on levels of follicle-stimulating hormone and folliculogenesis in *Rattus norvegicus* exposed to cigarette smoke. *Indian J Forensic Med Toxicol.* 2021;15(1):1820-1826.
5. Ernawati,NurlianiA. Antioxidant Effect of Eleutherine Americana Merr (Bawang Dayak) Ethanol Extract Towards the Tubule Seminiferous Microanatomy Structure Testis of Male Rats Exposed to Cigarette Smoke. *Sains dan Terap Kim.* 2012;6(2):93-100.
6. Hidayatik N, Purnomo A, Fikri F, Purnama MTE. Amelioration on oxidative stress, testosterone, and cortisol levels after administration of Vitamins C and e in albino rats with chronic variable stress. *Vet World.* 2021;14(1):137-143.

7. Guerriero G, Trocchia S, Abdel-Gawad FK, Ciarcia G. Roles of reactive oxygen species in the spermatogenesis regulation. *Front Endocrinol (Lausanne)*. 2014;5(APR):10-13.
8. Sèdes L, Thirouard L, Maqdasy S, Garcia M, Caira F. Cholesterol : A Gatekeeper of Male Fertility ? *2018;9(July):1-13.*
9. Kuang H, Yang F, Zhang Y, Wang T, Chen G. The Impact of Egg Nutrient Composition and Its Consumption on Cholesterol Homeostasis. *Cholesterol*. 2018;2018:6303810.
10. Souza UD, Biswas A, D'souza UJA, Bhat S. Dietary Hypercholesterolemia Induces Oxidative Stress Challenging Spermatogenesis in rat Model: A Link to Possible Infertility. *Int J Pharm Sci Res*. 2017;8(12):5065-5071.
11. Kusmiyati, Keman S, Amin M, Suwarno, Nugroho HSW. The role of Moringa oleifera leaves against oxidative stress and chronic inflammation: A review. *Indian J Public Heal Res Dev*. 2018;9(6):257-262.
12. Revianti S, Soetjipto, Rahayu RP, Parisihni K. Protective role of Sticophus hermanii ethanol extract supplementation to oxidative stress and oral hyperkeratosis in smoking exposed rats. *Int J ChemTech Res*. 2016;9(5):408-417.
13. Thant TM, Aminah NS, Kristanti AN, Ramadhan R, Phuwapraisirisan P, Takaya Y. A new pyrano coumarin from Clausena excavata roots displaying dual inhibition against α -glucosidase and free radical. *Nat Prod Res*. 2021;35(4):556-561.
14. Rosita N, Haryadi DM, Erawati T, Nanda RP, Soeratri W. Photostability study on character and antioxidant activity of tomato extract (*Solanum Lycopersicum l.*) in Nanostructured lipid carrier (NLC) and conventional cream. *Int J Drug Deliv Technol*. 2017;7(1):71-74.
15. Sholeh M, Isradji I, Oktaviyanti D, Fatmawati D. Pengaruh Ekstrak Terung Ungu (*Solanum melongena L.*) terhadap Motilitas dan Viabilitas Spermatozoa secara In Vitro. *J Wiyata*. 2020;7(1):78-85.
16. Wulansari D, Oktanella Y, Hendrawan VF, Agustina GC. Efektivitas Solasodine dan Gosipol sebagai Kandidat Kontrasepsi pada Hewan dalam Menghambat Reproduksi Tikus Putih (*Rattus norvegicus*) Jantan melalui Ekspresi LH dan Spermatogenesis as a Male Contraception Inhibit LH Expression and Spermatogenesi. *Vet Bio Clin J Vol*. 2019;1(2):51-59.
17. Lestari E, Sumarni NK, Mappiratu M. Kajian Aktivitas Antioksidan Mikrokapsul Ekstrak Kulit Terong Ungu (*Solanum melongena L.*). *KOVALEN J Ris Kim*. 2019;5(3):299-307.
18. Guimarães PR, Galvão AMP, Batista CM, Azevedo GS, Oliveira RD, Lamounier RP, et al. Eggplant (*Solanum melongena*) infusion has a modest and transitory effect on hypercholesterolemic subjects. *Brazilian J Med Biol Res*. 2000;33(9):1027-1036.
19. Niño-Medina G, Uriás-Orona V, Muy-Rangel MD, Heredia JB. Structure and content of phenolics in eggplant (*Solanum melongena*) - a review. *South African J Bot*. 2017;111:161-169.
20. Demir Y, Durmaz L, Taslimi PG. Antidiabetic properties of dietary phenolic compounds: Inhibition effects on α -amylase, aldose reductase, and α -glycosidase. *Biotechnol Appl Biochem*. 2019;66(5):781-786.
21. Sudeep HV, Venkatakrishna K, Dipak P, Shyamprasad K. Biomechanism of chlorogenic acid complex-mediated plasma free fatty acid metabolism in rat liver. *BMC Complement Altern Med*. 2016;16(1):4-8.
22. Gürbüz N, Uluişik S, Frary A, Frary A DS. Health benefits and bioactive compounds of eggplant. *Food Chem*. 2018;268:602-610.
23. Yarmohammadi F, Rahbardar MG, Hosseinzadeh H. Effect of eggplant (*Solanum melongena*) on the metabolic syndrome: A review. *Iran J Basic Med Sci*. 2021;24(4):420-427.
24. Samsudin RR, Kunsah B, Widayastuti R. The effect of pacitan's sweet oranges (*Citrus sinensis (L.) Osbeck*) peel powder on the lipid profile of males dyslipidemia rats (*Rattus norvegicus*). *Bali Med J*. 2017;6(3):51.
25. Keast DR, Fulgoni VL, Nicklas TA, O'Neil CE. Food sources of energy and nutrients among children in the United States: National Health and Nutrition Examination Survey 2003-2006. *Nutrients*. 2013;5(1):283-301.
26. Magnusdottir E V, Thorsteinsson T, Thorsteinsdottir S, Heimisdottir M, Olafsdottir K. Persistent organochlorines, sedentary occupation, obesity, and human male subfertility. *2005;20(1):208-215.*
27. Hjollund NHI, Bonde JPE, Jensen TK, Olsen J. Diurnal scrotal skin temperature and semen quality. *Int J Androl*. 2000;23(5):309-318.
28. Hajshafih M, Ghareaghaji R, Salemi S, Sadegh-Asadi N, Sadeghi-Bazargani H. Association of body mass index with some fertility markers among male partners of infertile couples. *Int J Gen Med*. 2013;6:447-451.
29. Kort HI, Massey JOEB, Elsner CW, Mitchell-Leef D, Shapiro DB, Witt MA, et al. Impact of Body Mass Index Values on Sperm Quantity and Quality. *2006;27(3):1150-1152.*
30. MacDonald AA, Herbison GP, Showell M, Farquhar CM. The impact of body mass index on semen parameters and reproductive hormones in human males: A systematic review with meta-analysis. *Hum Reprod Update*. 2009;16(3):293-311.
31. Tunc O, Bakos HW, Tremellen K. Impact of body mass index on seminal oxidative stress. *Andrologia*. 2011;43(2):121-128.
32. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, et al. Increased oxidative

STIMULUS FROM THE ACTIVATOR AGENT *SOLANUM MELONGENA L.*

- stress in obesity and its impact on metabolic syndrome. *J Clin Invest.* 2004;114(12):1752-1761.
33. Ferreira G, Costa C, Bassaizteguy V, Santos M, Cardozo R, Montes J, et al. Incubation of human sperm with micelles made from glycerophospholipid mixtures increases sperm motility and resistance to oxidative stress. *PLoS One.* 2018;13(6):1-32.
34. Liu CY, Chou YC, Lin SH, Wu ST, Cha TL, Chen HI, et al. Serum lipid profiles are associated with semen quality. *Asian J Androl.* 2017;19(6):633-638.
35. Meseguer M, Garrido N, Martínez-Conejero JA, Simón C, Pellicer A, Remohí J. Relationship between standard semen parameters, calcium, cholesterol contents, and mitochondrial activity in ejaculated spermatozoa from fertile and infertile males. *J Assist Reprod Genet.* 2004;21(12):445-451.
36. Samadian F, Towhidi A, Rezayazdi K, Bahreini M. Effects of dietary n-3 fatty acids on characteristics and lipid composition of ovine sperm. *Animal.* 2010;4(12):2017-2022.
37. Shan S, Xu F, Hirschfeld M, Brenig B. Sperm lipid markers of male fertility in mammals. *Int J Mol Sci.* 2021;22(16):8767.
38. Wood PL, Scoggin K, Ball BA, Troedsson MH, Squires EL. Lipidomics of equine sperm and seminal plasma: Identification of amphiphilic (O-acyl)- ω -hydroxy-fatty acids. *Theriogenology.* 2016;86(5):1212-1221.
39. Tsao CW, Liu CY, Chou YC, Cha TL, Chen SC, Hsu CY. Exploration of the association between obesity and semen quality in a 7 630 male population. *PLoS One.* 2015;10(3):1-13.
40. Liu CY, Chou YC, Chao JCJ, Hsu CY, Cha TL, Tsao CW. The association between dietary patterns and semen quality in a general Asian population of 7282 Males. *PLoS One.* 2015;10(7):1-12.
41. Volle DH, Mouzat K, Duggavathi R, Siddeek B, Déchelotte P, Sion B, et al. Multiple roles of the nuclear receptors for oxysterols liver X receptor to maintain male fertility. *Mol Endocrinol.* 2007;21(5):1014-1027.
42. Aquila S, Bonofoglio D, Gentile M, Middea E, Gabriele S, Belmonte M, et al. Peroxisome proliferator-activated receptor (PPAR) gamma is expressed by human spermatozoa: its potential role on the sperm physiology. *J Cell Physiol.* 2006;209(3):977.
43. Bowles J, Knight D, Smith C, Wilhelm D, Richman J, Mamiya S, et al. Retinoid signaling determines germ cell fate in mice. *Science (80-).* 2006;312(5773):596-600.
44. Lange Y, Ye J, Steck TL. Activation of Membrane Cholesterol by Displacement from Phospholipids. *J Biol Chem.* 2005;280(43):36126-36131.
45. Juyena NS, Stellella C. Seminal plasma: An essential attribute to spermatozoa. *J Androl.* 2012;33(4):536-551.
46. Wurlina W, Hariadi M, Safitri E, Susilowati S, Meles DK. The effect of crude guava leaf tannins on motility, viability, and intact plasma membrane of stored spermatozoa of Etawa crossbred goats. *Vet World.* 2020;13(3):530-534.
47. Sabeti P, Pourmasumi S, Rahiminia T, Akyash F, Talebi AR. Etiologies of sperm oxidative stress. *Int J Reprod Biomed.* 2016;14(4):231-240.