

LEMBAR PENILAIAN
PRE-TEST DAN POST-TEST

No	Variabel	Lingkup	Aspek yang diamati	Indikator	Penilaian			
					1	2	3	4
1.	Kemampuan berbahasa	Kemampuan berbahasa anak pada	Aktifitas anak kemampuan berbahasa	1. Anak mampu melakukan melengkapi kata/huruf seperti yang sudah dicontohkan oleh guru 2. Anak mampu melakukan mengulang kalimat sederhana yang sering didengar anak				
Jumlah								
Rata-rata								

Mojokerto,

Penilai

.....

Hasil output perhitungan SPSS 26

```

DATASET ACTIVATE DataSet0.
EXAMINE VARIABLES=Hasil_tes BY Kelompok
  /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL
  /COMPARE GROUPS
  /STATISTICS DESCRIPTIVES
  /CINTERVAL 95
  /MISSING LISTWISE
  /NOTOTAL.

```

Explore

Notes		
Output Created		26-OCT-2020 10:56:29
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.

Syntax	EXAMINE VARIABLES=Hasil_tes BY Kelompok /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL /COMPARE GROUPS /STATISTICS DESCRIPTIVES /CINTERVAL 95 /MISSING LISTWISE /NOTOTAL.
Resources	Processor Time 00:00:00,91 Elapsed Time 00:00:01,05

[DataSet0]

Kelompok

Case Processing Summary

Kelompok	Cases							
	Valid		Missing		Total			
	N	Percent	N	Percent	N	Percent	N	Percent
Hasil tes pre-test	15	100,0%	0	0,0%	15	100,0%	15	100,0%
post-test	15	100,0%	0	0,0%	15	100,0%	15	100,0%

Descriptives

Kelompok	Statistic	Std. Error
Hasil tes pre-test	Mean	2,40 ,214
	95% Confidence Interval for Lower Bound	1,94
	Mean Upper Bound	2,86
	5% Trimmed Mean	2,39
	Median	2,00
	Variance	,686

	Std. Deviation		,828
	Minimum		1
	Maximum		4
	Range		3
	Interquartile Range		1
	Skewness	-,070	,580
	Kurtosis	-,224	1,121
post-test	Mean		3,00
	95% Confidence Interval for Mean	Lower Bound	2,58
		Upper Bound	3,42
	5% Trimmed Mean		3,00
	Median		3,00
	Variance		,571
	Std. Deviation		,756
	Minimum		2
	Maximum		4
	Range		2
	Interquartile Range		2
	Skewness	,000	,580
	Kurtosis	-1,077	1,121

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Hasil tes	pre-test	,232	15	,029	,883	15	,052
	post-test	,233	15	,027	,823	15	,007


a. Lilliefors Significance Correction


Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	,878	1	28	,357
	Based on Median	,412	1	28	,526
	Based on Median and with adjusted df	,412	1	27,154	,526
	Based on trimmed mean	,867	1	28	,360

Hasil tes

Histograms





Stem-and-Leaf Plots

Hasil tes Stem-and-Leaf Plot for
Kelompok= pre-test

Frequency	Stem & Leaf
2,00	1 . 00
6,00	2 . 000000
6,00	3 . 000000
1,00	4 . 0

Stem width: 1
Each leaf: 1 case(s)


Hasil tes Stem-and-Leaf Plot for
Kelompok= post-test


Frequency	Stem & Leaf
-----------	-------------

4,00	2 .	0000
,00	2 .	
7,00	3 .	0000000
,00	3 .	
4,00	4 .	0000


Stem width: 1
Each leaf: 1 case(s)



Normal Q-Q Plots






Detrended Normal Q-Q Plots







```

ONEWAY Hasil_tes BY Kelompok
  /STATISTICS HOMOGENEITY
  /MISSING ANALYSIS.

```

Oneway

Notes

Output Created	26-OCT-2020 10:59:27	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.

Cases Used		Statistics for each analysis are based on cases with no missing data for any variable in the analysis.
Syntax		ONEWAY Hasil_tes BY Kelompok /STATISTICS HOMOGENEITY /MISSING ANALYSIS.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02

Test of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	,878	1	28	,357
	Based on Median	,412	1	28	,526
	Based on Median and with adjusted df	,412	1	27,154	,526
	Based on trimmed mean	,867	1	28	,360

ANOVA

Hasil tes

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	2,700	1	2,700	4,295	,048
Within Groups	17,600	28	,629		
Total	20,300	29			

```
T-TEST GROUPS=Kelompok(1 2)
/MISSING=ANALYSIS
/VARIABLES=Hasil_tes
/CRITERIA=CI (.95).
```

T-Test

Hasil tes	Equal variances assumed	,878	,357	- 2,073	28	,048	-,600	,289	- 1,193	-,007
	Equal variances not assumed			- 2,073	27,770	,048	-,600	,289	- 1,193	-,007

```

DATASET ACTIVATE DataSet1.
CORRELATIONS
  /VARIABLES=pre_test post_test
  /PRINT=TWOTAIL NOSIG
  /MISSING=PAIRWISE.

```

Correlations

Notes

Output Created		26-OCT-2020 11:03:13
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	15
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=pre_test post_test /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,05
	Elapsed Time	00:00:00,02

[DataSet1]

Correlations

		pre_test	post_test
pre_test	Pearson Correlation	1	,571*
	Sig. (2-tailed)		,026
	N	15	15
post_test	Pearson Correlation	,571*	1
	Sig. (2-tailed)	,026	
	N	15	15

*. Correlation is significant at the 0.05 level (2-tailed).

