

LEMBAR PENILAIAN
PRE-TEST DAN POST-TEST

No	Variabel	Lingkup	Aspek yang diamati	Indikator	Penilaian			
					1	2	3	4
1.	Perkembangan motorik halus	Perkembangan motorik halus anak pada	Aktifitas anak perkembangan motorik halus	1. Anak mampu melakukan melipat kertas origami sesuai dengan gambar alat komunikasi 2. Anak mampu melakukan mengurutkan arah pelipatan kertas origami sesuai petunjuk				
Jumlah								
Rata-rata								

Mojokerto,

Penilai

.....

Hasil output perhitungan SPSS 26

```

EXAMINE VARIABLES=Hasil_tes BY Kelompok
/PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL
/COMPARE GROUPS
/STATISTICS DESCRIPTIVES
/CINTERVAL 95
/MISSING LISTWISE
/NOTOTAL.

```

Explore

Notes		
Output Created		26-OCT-2020 18:07:22
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.

Syntax	EXAMINE VARIABLES=Hasil_tes BY Kelompok /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL /COMPARE GROUPS /STATISTICS DESCRIPTIVES /CINTERVAL 95 /MISSING LISTWISE /NOTOTAL.
Resources	Processor Time 00:00:01,13 Elapsed Time 00:00:01,22

Kelompok

Case Processing Summary

Kelompok	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Hasil tes pre-test	15	100,0%	0	0,0%	15	100,0%
post-test	15	100,0%	0	0,0%	15	100,0%

Descriptives

Kelompok	Statistic	Std. Error
Hasil tes pre-test	Mean	2,27
	95% Confidence Interval for Lower Bound	1,73
	Mean Upper Bound	2,80
	5% Trimmed Mean	2,24
	Median	2,00
	Variance	,924
	Std. Deviation	,961
	Minimum	1
	Maximum	4

	Range		3	
	Interquartile Range		2	
	Skewness		-,059	,580
	Kurtosis		-1,055	1,121
post-test	Mean		2,87	,192
	95% Confidence Interval for Mean	Lower Bound	2,46	
		Upper Bound	3,28	
	5% Trimmed Mean		2,85	
	Median		3,00	
	Variance		,552	
	Std. Deviation		,743	
	Minimum		2	
	Maximum		4	
	Range		2	
	Interquartile Range		1	
	Skewness		,227	,580
	Kurtosis		-,970	1,121

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Hasil tes	pre-test	,244	15	,017	,866	15	,030
	post-test	,238	15	,022	,817	15	,006


a. Lilliefors Significance Correction


Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	2,149	1	28	,154
	Based on Median	1,836	1	28	,186
	Based on Median and with adjusted df	1,836	1	27,813	,186
	Based on trimmed mean	2,043	1	28	,164

Hasil tes

Histograms





Stem-and-Leaf Plots

Hasil tes Stem-and-Leaf Plot for
Kelompok= pre-test

Frequency Stem & Leaf

```

4,00  1 . 0000
4,00  2 . 0000
6,00  3 . 000000
1,00  4 . 0

```

Stem width: 1
Each leaf: 1 case(s)


Hasil tes Stem-and-Leaf Plot for
Kelompok= post-test


Frequency Stem & Leaf

```
5,00 2 . 00000
,00 2 .
7,00 3 . 0000000
,00 3 .
3,00 4 . 000
```


Stem width: 1
Each leaf: 1 case(s)



Normal Q-Q Plots






Detrended Normal Q-Q Plots







```

ONEWAY Hasil_tes BY Kelompok
  /STATISTICS HOMOGENEITY
  /MISSING ANALYSIS.

```

Oneway

Notes

Output Created		26-OCT-2020 18:22:24
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each analysis are based on cases with no missing data for any variable in the analysis.

Syntax	ONEWAY Hasil_tes BY Kelompok /STATISTICS HOMOGENEITY /MISSING ANALYSIS.
Resources	Processor Time 00:00:00,00 Elapsed Time 00:00:00,01

Test of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	2,149	1	28	,154
	Based on Median	1,836	1	28	,186
	Based on Median and with adjusted df	1,836	1	27,813	,186
	Based on trimmed mean	2,043	1	28	,164

ANOVA

Hasil tes

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	2,700	1	2,700	3,658	,066
Within Groups	20,667	28	,738		
Total	23,367	29			

```
T-TEST GROUPS=Kelompok(1 2)
/MISSING=ANALYSIS
/VARIABLES=Hasil_tes
/CRITERIA=CI (.95).
```

T-Test

Notes

Output Created	26-OCT-2020 18:24:16
Comments	

Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User defined missing values are treated as missing.
	Cases Used	Statistics for each analysis are based on the cases with no missing or out-of-range data for any variable in the analysis.
Syntax	T-TEST GROUPS=Kelompok(1 2) /MISSING=ANALYSIS /VARIABLES=Hasil_tes /CRITERIA=CI(.95).	
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,02

Group Statistics

	Kelompok	N	Mean	Std. Deviation	Std. Error Mean
Hasil tes	pre-test	15	2,27	,961	,248
	post-test	15	2,87	,743	,192

Independent Samples Test

	Levene's Test for Equality of Variances		t-test for Equality of Means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
								Lower	Upper
Hasil tes Equal variances assumed	2,149	,154	-1,913	28	,066	-,600	,314	-1,243	,043

Equal variances not assumed			- 1,913	26,333	,067	-,600	,314	- 1,244	,044
-----------------------------	--	--	---------	--------	------	-------	------	---------	------

```

DATASET ACTIVATE DataSet1.
CORRELATIONS
/VARIABLES=pre_test post_test
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

Correlations

Notes

Output Created	26-OCT-2020 18:26:33	
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	15
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=pre_test post_test /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,08
	Elapsed Time	00:00:00,02

[DataSet1]

Correlations

		pre_test	post_test
pre_test	Pearson Correlation	1	,053
	Sig. (2-tailed)		,850
	N	15	15
post_test	Pearson Correlation	,053	1
	Sig. (2-tailed)	,850	
	N	15	15