

LEMBAR PENILAIAN
PRE-TEST DAN POST-TEST

No	Variabel	Lingkup	Aspek yang diamati	Indikator	Penilaian			
					1	2	3	4
1.	Kemampuan mengenali huruf awal	Kemampuan mengenal huruf awal anak pada	Aktifitas anak kemampuan mengenal huruf awal	1. Anak mampu melakukan menunjukkan huruf N-A-S-I secara terpisah 2. Anak mampu melakukan menunjukkan huruf awal benda dari unsur N-A-S-I				
Jumlah								
Rata-rata								

Mojokerto,

Penilai

.....

Hasil output perhitungan SPSS 26

```

DATASET ACTIVATE DataSet0.
EXAMINE VARIABLES=Hasil_tes BY Kelompok
  /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL
  /COMPARE GROUPS
  /STATISTICS DESCRIPTIVES
  /CINTERVAL 95
  /MISSING LISTWISE
  /NOTOTAL.

```

Explore

Notes

Output Created	24-OCT-2020 22:34:18	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	28
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.

Syntax	EXAMINE VARIABLES=Hasil_tes BY Kelompok /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL /COMPARE GROUPS /STATISTICS DESCRIPTIVES /CINTERVAL 95 /MISSING LISTWISE /NOTOTAL.
Resources	Processor Time 00:00:01,02 Elapsed Time 00:00:01,20

[DataSet0]

Kelompok

Case Processing Summary

Kelompok	N	Valid		Missing		Total	
		N	Percent	N	Percent	N	Percent
Hasil tes pre-test	14	14	100,0%	0	0,0%	14	100,0%
post-test	14	14	100,0%	0	0,0%	14	100,0%

Descriptives

Kelompok	Statistic	Std. Error
Hasil tes pre-test	Mean	2,43
	95% Confidence Interval for Lower Bound	1,89
	Mean Upper Bound	2,97
	5% Trimmed Mean	2,42
	Median	2,00
	Variance	,879

	Std. Deviation		,938	
	Minimum		1	
	Maximum		4	
	Range		3	
	Interquartile Range		1	
	Skewness		,240	,597
	Kurtosis		-,491	1,154
post-test	Mean		2,93	,195
	95% Confidence Interval for Mean	Lower Bound	2,51	
		Upper Bound	3,35	
	5% Trimmed Mean		2,92	
	Median		3,00	
	Variance		,533	
	Std. Deviation		,730	
	Minimum		2	
	Maximum		4	
	Range		2	
	Interquartile Range		1	
	Skewness		,113	,597
	Kurtosis		-,856	1,154

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Hasil tes	pre-test	,248	14	,020	,893	14	,088
	post-test	,253	14	,015	,821	14	,009


a. Lilliefors Significance Correction


Test of Homogeneity of Variance

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	1,821	1	26	,189
	Based on Median	,807	1	26	,377
	Based on Median and with adjusted df	,807	1	23,528	,378
	Based on trimmed mean	1,761	1	26	,196

Hasil tes

Histograms





Stem-and-Leaf Plots

Hasil tes Stem-and-Leaf Plot for
Kelompok= pre-test

Frequency	Stem & Leaf
2,00	1 . 00
6,00	2 . 000000
4,00	3 . 0000
2,00	4 . 00

Stem width: 1
Each leaf: 1 case(s)


Hasil tes Stem-and-Leaf Plot for
Kelompok= post-test


Frequency	Stem & Leaf
-----------	-------------

4,00	2 . 0000
,00	2 .
7,00	3 . 0000000
,00	3 .
3,00	4 . 000


Stem width: 1
Each leaf: 1 case(s)



Normal Q-Q Plots






Detrended Normal Q-Q Plots







```

ONEWAY Hasil_tes BY Kelompok
  /STATISTICS HOMOGENEITY
  /MISSING ANALYSIS.

```

Oneway

Notes

Output Created		24-OCT-2020 22:36:50
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	28
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.

Cases Used		Statistics for each analysis are based on cases with no missing data for any variable in the analysis.
Syntax		ONEWAY Hasil_tes BY Kelompok /STATISTICS HOMOGENEITY /MISSING ANALYSIS.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01

Test of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	1,821	1	26	,189
	Based on Median	,807	1	26	,377
	Based on Median and with adjusted df	,807	1	23,528	,378
	Based on trimmed mean	1,761	1	26	,196

ANOVA

Hasil tes

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	1,750	1	1,750	2,479	,127
Within Groups	18,357	26	,706		
Total	20,107	27			

```
T-TEST GROUPS=Kelompok(1 2)
/MISSING=ANALYSIS
/VARIABLES=Hasil_tes
/CRITERIA=CI (.95).
```

T-Test

Hasil tes	Equal variances assumed	1,821	,189	-	26	,127	-,500	,318	-	,153
	Equal variances not assumed			-	24,526	,128	-,500	,318	-	,155
				1,574					1,153	
				1,574					1,155	

```

DATASET ACTIVATE DataSet1.
CORRELATIONS
/VARIABLES=pre_test post_test
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

Correlations

Notes

Output Created	24-OCT-2020 22:40:13	
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	14
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=pre_test post_test /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,03
	Elapsed Time	00:00:00,01

[DataSet1]

Correlations

		pre_test	post_test
pre_test	Pearson Correlation	1	,385
	Sig. (2-tailed)		,174
	N	14	14
post_test	Pearson Correlation	,385	1
	Sig. (2-tailed)	,174	
	N	14	14