

LEMBAR PENILAIAN
PRE-TEST DAN POST-TEST

No	Variabel	Lingkup	Aspek yang diamati	Indikator	Penilaian			
					1	2	3	4
1.	Keterampilan berbahasa tulis	Keterampilan berbahasa tulis anak pada	Aktifitas anak Keterampilan berbahasa tulis	1. Anak mampu melakukan menulis huruf awal gambar kendaraan 2. Anak mampu melakukan mencari huruf awal yang sama pada gambar kendaraan				
Jumlah								
Rata-rata								

Mojokerto,

Penilai

.....

Hasil output perhitungan SPSS 26

```

DATASET ACTIVATE DataSet0.
EXAMINE VARIABLES=Hasil_tes BY Kelompok
  /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL
  /COMPARE GROUPS
  /STATISTICS DESCRIPTIVES
  /CINTERVAL 95
  /MISSING LISTWISE
  /NOTOTAL.

```

Explore

Notes		
Output Created		26-OCT-2020 11:46:31
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values for dependent variables are treated as missing.
	Cases Used	Statistics are based on cases with no missing values for any dependent variable or factor used.

Syntax		EXAMINE VARIABLES=Hasil_tes BY Kelompok /PLOT BOXPLOT STEMLEAF HISTOGRAM NPLOT SPREADLEVEL /COMPARE GROUPS /STATISTICS DESCRIPTIVES /CINTERVAL 95 /MISSING LISTWISE /NOTOTAL.
Resources	Processor Time	00:00:00,84
	Elapsed Time	00:00:00,85

[DataSet0]

Warnings

The regression slope of the spread vs.level plot for Hasil tes*Kelompok cannot be computed because there are not enough unique spread/level pairs.

Kelompok

Case Processing Summary

	Kelompok	Valid		Cases Missing		Total	
		N	Percent	N	Percent	N	Percent
Hasil tes	pre-test	15	100,0%	0	0,0%	15	100,0%
	post-test	15	100,0%	0	0,0%	15	100,0%

Descriptives

Kelompok	Statistic	Std. Error
----------	-----------	------------

Hasil tes	pre-test	Mean		2,60	,190
		95% Confidence Interval for Mean	Lower Bound	2,19	
			Upper Bound	3,01	
		5% Trimmed Mean		2,61	
		Median		3,00	
		Variance		,543	
		Std. Deviation		,737	
		Minimum		1	
		Maximum		4	
		Range		3	
		Interquartile Range		1	
		Skewness		-,396	,580
		Kurtosis		,425	1,121
		post-test	Mean		2,93
		95% Confidence Interval for Mean	Lower Bound	2,49	
			Upper Bound	3,38	
		5% Trimmed Mean		2,93	
		Median		3,00	
		Variance		,638	
		Std. Deviation		,799	
	Minimum		2		
	Maximum		4		
	Range		2		
	Interquartile Range		2		
	Skewness		,128	,580	
	Kurtosis		-1,348	1,121	

Tests of Normality

	Kelompok	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
		Statistic	df	Sig.	Statistic	df	Sig.
Hasil tes	pre-test	,306	15	,001	,846	15	,015
	post-test	,212	15	,068	,817	15	,006

a. Lilliefors Significance Correction


Test of Homogeneity of Variance


	Levene Statistic	df1	df2	Sig.

Hasil tes	Based on Mean	,003	1	28	,955
	Based on Median	,100	1	28	,754
	Based on Median and with adjusted df	,100	1	26,610	,754
	Based on trimmed mean	,008	1	28	,931

Hasil tes

Histograms





Stem-and-Leaf Plots

Hasil tes Stem-and-Leaf Plot for
Kelompok= pre-test

Frequency	Stem & Leaf
1,00	1 . 0
5,00	2 . 00000
8,00	3 . 00000000
1,00	4 . 0

Stem width: 1
Each leaf: 1 case(s)


Hasil tes Stem-and-Leaf Plot for
Kelompok= post-test


Frequency	Stem & Leaf
-----------	-------------

5,00	2 .	00000
,00	2 .	
6,00	3 .	000000
,00	3 .	
4,00	4 .	0000


Stem width: 1
Each leaf: 1 case (s)



Normal Q-Q Plots






Detrended Normal Q-Q Plots







```

ONEWAY Hasil_tes BY Kelompok
  /STATISTICS HOMOGENEITY
  /MISSING ANALYSIS.

```

Oneway

Notes

Output Created	26-OCT-2020 11:48:51	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.

Cases Used		Statistics for each analysis are based on cases with no missing data for any variable in the analysis.
Syntax		ONEWAY Hasil_tes BY Kelompok /STATISTICS HOMOGENEITY /MISSING ANALYSIS.
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01

Test of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
Hasil tes	Based on Mean	,003	1	28	,955
	Based on Median	,100	1	28	,754
	Based on Median and with adjusted df	,100	1	26,610	,754
	Based on trimmed mean	,008	1	28	,931

ANOVA

Hasil tes

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	,833	1	,833	1,411	,245
Within Groups	16,533	28	,590		
Total	17,367	29			

```
T-TEST GROUPS=Kelompok(1 2)
/MISSING=ANALYSIS
/VARIABLES=Hasil_tes
/CRITERIA=CI (.95).
```

T-Test

Hasil tes	Equal variances assumed	,003	,955	-1,188	28	,245	-,333	,281	-,908	,241
	Equal variances not assumed			-1,188	27,819	,245	-,333	,281	-,908	,242

```

DATASET ACTIVATE DataSet1.
CORRELATIONS
  /VARIABLES=pre_test post_test
  /PRINT=TWOTAIL NOSIG
  /MISSING=PAIRWISE.

```

Correlations

Notes

Output Created		26-OCT-2020 11:51:59
Comments		
Input	Active Dataset	DataSet1
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	15
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=pre_test post_test /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00,02
	Elapsed Time	00:00:00,01

[DataSet1]

Correlations

		pre_test	post_test
pre_test	Pearson Correlation	1	,437
	Sig. (2-tailed)		,103
	N	15	15
post_test	Pearson Correlation	,437	1
	Sig. (2-tailed)	,103	
	N	15	15