

Lampiran 1

Uji Normalitas

UJI NORMALITAS

NPar Tests

| Notes | | |
|----------------|--------------------------------------|--|
| Output Created | | 10-AUG-2020 18:46:44 |
| Comments | | |
| Input | Data | D:\Maret 2020\olah data\otw\listi 2.sav |
| | Active Dataset | DataSet1 |
| | Filter | <none> |
| | Weight | <none> |
| | Split File | <none> |
| | N of Rows in Working Data File | 5 |
| | Missing Value Handling | Definition of Missing |
| | Cases Used | Statistics for each test are based on all cases with valid data for the variable(s) used in that test. |
| Syntax | | NPAR TESTS /K-S(NORMAL)=RES_1 /MISSING ANALYSIS. |
| Resources | Processor Time | 00:00:00.02 |
| | Elapsed Time | 00:00:00.02 |
| | Number of Cases Allowed ^a | 786432 |

a. Based on availability of workspace memory.

One-Sample Kolmogorov-Smirnov Test

| | | Unstandardized Residual |
|----------------------------------|----------------|----------------------------|
| N | | 5 |
| Normal Parameters ^{a,b} | Mean | .0000000 |
| | Std. Deviation | 1.18188867 |
| Most Extreme Differences | Absolute | .226 |
| | Positive | .226 |
| | Negative | -.188 |
| Test Statistic | | .226 |
| Asymp. Sig. (2-tailed) | | .200 ^{c,d} |

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

d. This is a lower bound of the true significance.

Lampiran 2

Regresi Linier Sederhana

Regression

Notes

| | | |
|------------------------|--|---|
| Output Created | | 10-AUG-2020 18:45:48 |
| Comments | | |
| Input | Data | D:\Maret 2020\olah data\otw\listi 2.sav |
| | Active Dataset | DataSet1 |
| | Filter | <none> |
| | Weight | <none> |
| | Split File | <none> |
| | N of Rows in Working Data File | 5 |
| Missing Value Handling | Definition of Missing | User-defined missing values are treated as missing. |
| | Cases Used | Statistics are based on cases with no missing values for any variable used. |
| Syntax | <pre> REGRESSION /MISSING LISTWISE /STATISTICS COEFF OUTS R ANOVA /CRITERIA=PIN(.05) POUT(.10) /NOORIGIN /DEPENDENT ROA /METHOD=ENTER PP /RESIDUALS HISTOGRAM(ZRESID) NORMPROB(ZRESID) /SAVE RESID. </pre> | |
| Resources | Processor Time | 00:00:03.03 |
| | Elapsed Time | 00:00:02.06 |

| | | |
|-----------------------|--|-------------------------|
| | Memory Required | 2608 bytes |
| | Additional Memory Required for Residual Plots | 640 bytes |
| Variables Modified | Created or RES_1 | Unstandardized Residual |

Variables Entered/Removed^a

| Model | Variables Entered | Variables Removed | Method |
|-------|-------------------|-------------------|--------|
| 1 | PP ^b | . | Enter |

- a. Dependent Variable: ROA
b. All requested variables entered.

Model Summary^b

| Model | R | R Square | Adjusted R Square | Std. Error of the Estimate |
|-------|-------------------|----------|-------------------|----------------------------|
| 1 | .556 ^a | .309 | .079 | 1.36472748192 3861 |

- a. Predictors: (Constant), PP
b. Dependent Variable: ROA

ANOVA^a

| Model | | Sum of Squares | df | Mean Square | F | Sig. |
|-------|------------|----------------|----|-------------|-------|-------------------|
| 1 | Regression | 2.501 | 1 | 2.501 | 1.343 | .330 ^b |
| | Residual | 5.587 | 3 | 1.862 | | |
| | Total | 8.088 | 4 | | | |

- a. Dependent Variable: ROA
b. Predictors: (Constant), PP

Coefficients^a

| Model | | Unstandardized Coefficients | | Standardized Coefficients | t | Sig. |
|-------|------------|-----------------------------|------------|---------------------------|-------|------|
| | | B | Std. Error | Beta | | |
| 1 | (Constant) | -.133 | 3.267 | | -.041 | .970 |
| | PP | .931 | .804 | .556 | 1.159 | .330 |

- a. Dependent Variable: ROA

Residuals Statistics^a

| | Minimum | Maximum | Mean | Std. Deviation | N |
|----------------------|----------------------------|-----------------------|-----------------------|-----------------------|---|
| Predicted Value | 2.51966524124 1455 | 4.33212947845 4590 | 3.58569511797 5396 | .790721898493 133 | 5 |
| Residual | - 1.04661166667 9382 | 1.91070497035 9802 | .000000000000 001 | 1.18188866858 8832 | 5 |
| Std. Predicted Value | -1.348 | .944 | .000 | 1.000 | 5 |
| Std. Residual | -.767 | 1.400 | .000 | .866 | 5 |

a. Dependent Variable: ROA

Lampiran 3
Nilai Kritis t-Student

NILAI KRITIS T-STUDENT (TABLE t)

| Pr DF | 0.100 | 0.050 | 0.025 | 0.010 |
|------------------------|--------------|--------------|--------------|--------------|
| 1. | 6.314 | 12.706 | 25.452 | 63.657 |
| 2. | 2.920 | 4.303 | 6.205 | 9.925 |
| 3. | 2.353 | 3.182 | 4.177 | 5.841 |
| 4. | 2.132 | 2.776 | 3.495 | 4.604 |
| 5. | 2.015 | 2.571 | 3.163 | 4.032 |
| 6. | 1.943 | 2.447 | 2.969 | 3.707 |
| 7. | 1.895 | 2.365 | 2.841 | 3.499 |
| 8. | 1.860 | 2.306 | 2.752 | 3.355 |
| 9. | 1.833 | 2.262 | 2.685 | 3.250 |
| 10. | 1.812 | 2.228 | 2.634 | 3.169 |
| 11. | 1.796 | 2.201 | 2.593 | 3.106 |
| 12. | 1.782 | 2.179 | 2.560 | 3.055 |
| 13. | 1.771 | 2.160 | 2.533 | 3.012 |
| 14. | 1.761 | 2.145 | 2.510 | 2.977 |
| 15. | 1.753 | 2.131 | 2.490 | 2.947 |
| 16. | 1.746 | 2.120 | 2.473 | 2.921 |
| 17. | 1.740 | 2.110 | 2.458 | 2.898 |
| 18. | 1.734 | 2.101 | 2.445 | 2.878 |
| 19. | 1.729 | 2.093 | 2.433 | 2.861 |
| 20. | 1.725 | 2.086 | 2.423 | 2.845 |
| 21. | 1.721 | 2.080 | 2.414 | 2.831 |
| 22. | 1.717 | 2.074 | 2.405 | 2.819 |
| 23. | 1.714 | 2.069 | 2.398 | 2.807 |
| 24. | 1.711 | 2.064 | 2.391 | 2.797 |
| 25. | 1.708 | 2.060 | 2.385 | 2.787 |
| 30. | 1.697 | 2.042 | 2.360 | 2.750 |
| 35. | 1.690 | 2.030 | 2.342 | 2.724 |
| 40. | 1.684 | 2.021 | 2.329 | 2.704 |
| 45. | 1.679 | 2.014 | 2.319 | 2.690 |
| 50. | 1.676 | 2.009 | 2.311 | 2.678 |

Sumber : Microsoft Excell versi 2007