


Perpustakaan Um Surabaya

THE POTENCY OF WATER LEAVES EXTRACT OF BAY (*Syzygium polyanthum*) AND PAPAYA (*Carica papaya*) AS LARVICIDES T...

 Quick Submit

 Quick Submit

 Universitas Muhammadiyah Surabaya

Document Details

Submission ID

trn:oid::1:3095412786

Submission Date

Nov 28, 2024, 10:54 AM GMT+7

Download Date

Nov 28, 2024, 11:41 AM GMT+7

File Name

AY..._oktober_2022_-_penulis_ke_1_-_junal_bioma_-_sinta_3.pdf

File Size

351.1 KB

7 Pages

2,227 Words

12,302 Characters

16% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.





Filtered from the Report

- ▶ Small Matches (less than 15 words)
- ▶ Internet sources
- ▶ Publications




Exclusions

- ▶ 8 Excluded Sources

Match Groups

-  **8** Not Cited or Quoted 9%
Matches with neither in-text citation nor quotation marks
-  **0** Missing Quotations 0%
Matches that are still very similar to source material
-  **5** Missing Citation 7%
Matches that have quotation marks, but no in-text citation
-  **0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 0%  Internet sources
- 0%  Publications
- 16%  Submitted works (Student Papers)

Integrity Flags

0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

Match Groups

- 8** Not Cited or Quoted 9%
Matches with neither in-text citation nor quotation marks
- 0** Missing Quotations 0%
Matches that are still very similar to source material
- 5** Missing Citation 7%
Matches that have quotation marks, but no in-text citation
- 0** Cited and Quoted 0%
Matches with in-text citation present, but no quotation marks

Top Sources

- 0% Internet sources
- 0% Publications
- 16% Submitted works (Student Papers)

Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Student papers	
	Copperbelt University	2%
2	Student papers	
	Vrije Universiteit Amsterdam	2%
3	Student papers	
	Universitas Jember	2%
4	Student papers	
	Indian Institute of Technology Guwahati	2%
5	Student papers	
	The University of Manchester	2%
6	Student papers	
	Universiti Kebangsaan Malaysia	2%
7	Student papers	
	Universitas Islam Lamongan	2%
8	Student papers	
	Swinburne University of Technology	1%
9	Student papers	
	Universitas Airlangga	1%
10	Student papers	
	Universitas Tanjungpura	1%

11

Student papers

University of Arizona

1%



Available online at BIOMA: Jurnal Ilmiah Biologi
 Websites: <http://journal.upgris.ac.id/index.php/bioma/index>
BIOMA: Jurnal Ilmiah Biologi, 11(2), October 2022, 135-141
 Doi: <https://doi.org/10.26877/bioma.v11i2.10587>



THE POTENCY OF WATER LEAVES EXTRACT OF BAY (*Syzygium polyanthum*) AND PAPAYA (*Carica papaya*) AS LARVICIDES TO FILARIA VECTOR *Culex quequesciatus* (Diptera: Culicidae)

Vella Rohmayani^{1*}, Anindita Riesti Retno Arimurti², Rinza Rahmawati Samsudin²

¹Department of Medical Laboratory Technologist, Vocational Program, Faculty of Health Sciences, University of Muhammadiyah Surabaya.

Jl. Sutorejo No.59, Surabaya, East Java, Indonesia 60113

²Medical Laboratory Technology, Study Program DIII, Faculty of Health Sciences, University of Muhammadiyah Surabaya.

Jl. Sutorejo No.59, Surabaya, East Java, Indonesia 60113

*Corresponding author: vella_rohmayani@um-surabaya.ac.id

ARTICLE INFO

Article history
 Submission
 December 26th, 2021
 Revision
 March 14th, 2022
 Accepted
 June 9th, 2022

Keywords:

Bay leaves (*Syzygium polyanthum*)
Culex quequesciatus
 Larvacide
 Papaya leaves (*Carica papaya*)

ABSTRACT

Culex quequesciatus is a mosquito that acts as a vector of filariasis disease. In Indonesia filariasis is still one of the endemic diseases. This proves that the handling of vector *Culex quequesciatus* is still not done optimally. One way that can be done to control mosquito vectors one of them is to use larvacide. Larvacide which is widely used today is still synthetic. Although very efficient in causing mosquito mortality, larvacide synthesis harms the environment and can cause vectors resistance. Then another larvacide alternative is needed that is effective and does not cause negative effects. Plants can be a choice of larvacide from nature. The study aims to find out the potential of the water extract of bay leaves and papaya leaves in killing the larvae of the *Cx. quequesciatus* mosquito as well as the value of LC_{50} after 24-hour observation. The type of research used is experimental research. The concentration of the water extract of bay leaves and papaya leaves used is 10%, 30%, 50%, 70%, and 90% with three times replication. LC_{50} is calculated using the Finney formula (1971), the calculation is carried out using application SPSS statistical version 26. Based on the research conducted, the value of bay leaf water extract $LC_{50} = 62.89$ g/ml and papaya leaf water extract $LC_{50} = 45.10$ g/ml. So based on the LC_{50} value, it can be concluded that papaya leaf water extract is more effective in causing mortality *Cx. quequesciatus* compared with bay leaf water extract.

INTRODUCTION

Filariasis is a disease that transmitted through mosquito vector intermediaries. In Indonesia, filariasis disease is still one of the endemic diseases that cause health problems

Rohmayani, et al. The Potency of Water Leaves Extract Of Bay...

in the community until now. There were 10,681 cases in Indonesia in 2018 that were spread in 236 districts/cities from 28 provinces are declared as filariasis endemic. The highest cases of filariasis in Indonesia occurred in Papua Province with 3,615 cases, then East Nusa Tenggara with 1,542 cases, and followed by West Java province with the 985 cases (Indah, 2019).

The main step that can be done to prevent the transmission of transmitted diseases through vector intermediaries is controlling the disease-transmitting vector (Fouque & Reeder, 2019). One of the ways to control the disease vectors is chemical using larvicides. However, the larvacides that are widely used today are still synthetic larvacides that cause various negative effects such as causing environmental pollution, and resistance and can be toxic to non-target organisms (Kumar *et al.*, 2014; Princess & Kurnia, 2018). Therefore it is necessary to develop natural larvacides that are cheaper, eco-friendly does not toxic to non-target organisms, biodegradable, and do not destroy ecosystems (Astriani & Widawati, 2016; Kumar *et al.*, 2014).

Plants that may potentially be used as natural larvacides are bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*) because they contain secondary metabolite compounds. According to Utami & Puspaningtyas (2014), papaya leaves contain essential oils, lactones, phenols, saponins, and polyphenols. According to Cahyati *et al.* (2019), papaya leaves contain the enzymes papain, saponins, tannins, flavonoids, steroids, karpaina alkaloids, glycosids, karposids, saccharides, dextrose, and fructose.

Tannin compounds, saponins, flavonoids, and alkaloids have toxic properties for animals, so they can be used as larvicides (Ariesta, 2013; Gutierrez *et al.*, 2014). Tannins can damage the digestive part of insects, as well as cause disruptions in the function of organism water absorption (Mardiana *et al.*, 2009; Pradani, 2009). Saponins can inhibit growth, and development and interfere insect reproductive system. Alkaloids and flavonoids can inhibit insect-eating, besides essential oils and flavonoids can work as respiratory toxins (Pradani, 2009).

Research of potential testing by bay leaf extract and papaya leaves as natural larvacides has been done by researchers before. This research aims to increase information about the potential of bay leaves and papaya leaves using the water extract method, the use of squeeze method can be directly applied by the general public, in

Rohmayani, *et al.* The Potency of Water Leaves Extract Of Bay...

contrast to extract method which must be carried out in laboratory. Therefore, it can be acknowledged the potential or effectiveness of each water extract of bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*) towards the death rate of *Cx. quenequefasciatus* mosquitoes.

MATERIALS AND METHODS

Tools and Materials

The tools used in this study are blenders, measuring cups, digital scales, beaker glass, petri dishes, filters, stirrers, rubber, stationery, and la bel paper. While the ingredients used are bay leaves (*Syzygium polyanthum*), papaya leaves (*Carica papaya*), aquades, and *Cx. quenequefasciatus* mosquito larvae. As for the leaves used are dark green leaves.

Research Procedures

First, make sharpening bay leaves (*Syzygium polyanthum*) and papaya (*Carica papaya*). Papaya leaves (*Syzygium polyanthum*) 100 grams. Second, washed thoroughly, dried, and cut into small pieces to facilitate the smoothing process. Then, papaya leaves (*Syzygium polyanthum*) are put in a blender and added 100 ml of akuades. The blender results are water extractd and filtered. Furthermore, the result of this water extract is a 100% concentration. The result of this water extract is diluted using the formula:

$$V_1 \times M_1 = V_2 \times M_2$$

Description:

V1 = Initial volume

M1 = Initial concentration

V2 = Final volume

M2 = Final concentration

(Nurhaifah & Sukesu, 2015)

The way to make a water extract of papaya leaf (*Carica papaya*) is similar to how to make a water extract of bay leaves (*Syzygium polyanthum*). Next, make a water extract on each leave with concentrations of 10% (P1), 30% (P2), 50% (P3), 70% (P4), and 90% (P5). After that, the *Cx. quenequefasciatus* instar III larvae are put into a petri dish with as many as 25 individu for each treatment. Then, the petri dish treatment was closed

Rohmayani, et al. The Potency of Water Leaves Extract Of Bay...

and incubated for 24 hours after treatment. The experiment are repeated 3 times. After 24 hours, the number of dead larvae was calculated and recorded for each treatment on the data tabulation table. Mosquito larvae are claimed as dead larvae are larvae if the condition is not moving anymore and/or the body is destroyed.

Data Analysis

Calculated LC50 following to Probit analyssis by formula Finney 1971). The calculating using SPSS aplication version 26.

RESULTS AND DISCUSSION

The deaths number of *Cx. quinquefasciatus* larvae after being treated by bay leaf and papaya leaves can be seen in **Figure 1**. Based on **Figure 1** it can be observed that in treatment 1, 10% of the water extract of bay leaves and papaya leaves causes the death of larvae of *Cx. quinquefasciatus*, the percentage of death in mosquito larvae will increase proportionally to the amount of concentration of water extract given.




Figure 1. Death average of *Cx. quinquefasciatus* larvae after 24 hours giving a water extract of bay leaves (*Syzygium polyanthum*) and papaya leaves (*Carica papaya*)

Rohmayani, *et al.* The Potency of Water Leaves Extract Of Bay...

The lowest percentage of deaths occurred in the treatment with a 10% concentration and the highest percentage of *Cx. quinquefasciatus* mosquito larvae deaths occurred in the treatment with a 90% concentration both in the treatment of bay leaf water extract or papaya leaf water extract. Based on these results, it can be seen that the higher the concentration or content of active compounds in bay leaves (*Syzygium polyanthum*) and papaya leaves, the mortality rate in *Cx. quinquefasciatus* larvae will increase too.

LC₅₀ for bay leaves = 62.89 g/ml and LC₅₀ for papaya leaves = 45.10 g/ml. Based on these results, the LC₅₀ papaya leaves is smaller than the LC₅₀ bay leaves, it can be concluded that the papaya leaf water extract is more effective than the bay leaf water extract. Both Bay leaf (*Syzygium Polyanthum*) and papaya leaves (*Carica papaya*) contain various secondary metabolite compounds, such as flavonoids, tannins, saponins, phenolic hydroquinone alkaloids, steroids, and triterpenoids that are toxic to animals (Cahyati *et al.*, 2019; Sulistiyani *et al.*, 2014; Utami & Puspaningtyas, 2014).

Based on the results of the research, tannin compounds can damage the digestive to some parts of insects and able to make disturb the function of water absorption in organisms (Mardiana *et al.*, 2009; Pradani, 2009). Saponins can cause a decrease in insect food intake, causing stunted growth, development, and disruption of the insect reproductive system. In addition, other active compounds such as alkaloids and flavonoids are also able to act as compounds that inhibit eating insects, essential oils and flavonoids can also work as respiratory toxins (Pradani, 2009).

CONCLUSION

The LC₅₀ of papaya leaves (*Carica papaya*) is smaller than the LC₅₀ of bay leaves (*Syzygium polyanthum*). It can be concluded that the papaya leaves water extract is more effective than the bay leaf water extract as larvacide to *Cx. quinquefasciatus*.

REFERENCES

- 3 Ariesta, A. A. (2013). *Uji Efektifitas Larutan Daun Pepaya (Carica papaya) Sebagai Larvasida Terhadap Kematian Larva Nyamuk Aedes aegypti di Laboratorium B2P2VRP*. Balai Besar Litbang Vektor dan Reservoir Penyakit. Salatiga.
- 3 Astriani, Y., & Widawati, M. (2016). *Potensi tanaman di Indonesia sebagai larvasida*

Rohmayani, et al. The Potency of Water Leaves Extract Of Bay...

alami untuk *Aedes aegypti*. *SPIRAKEL*, 8(2), 37–46.
<https://doi.org/10.22435/spirakel.v8i2.6166>

6 Cahyati, W. H., Asmara, W., Umniyati, S. R., & Mulyaningsih, B. (2017). The phytochemical analysis of hay infusions and papaya leaf juice as an attractant containing insecticide for *Aedes aegypti*. *Jurnal Kesehatan Masyarakat*, 12(2), 218–224. <https://doi.org/10.15294/kemas.v12i2.6223>

1 Cahyati, W. H., Asmara, W., Umniyati, S. R., & Mulyaningsih, B. (2019). Biolarvicidal effects of papaya leaves juice against *Aedes aegypti* Linn larvae. *Journal of International Dental and Medical Research*, 12(2), 780–785. http://www.jidmr.com/journal/wp-content/uploads/2019/07/70_M17_449_Widya_Hary_Cahyati_Layout.pdf

7 Dwiyanti, R. D., Dediq, R., & Thuraidah, A. (2017). Daya bunuh ekstrak air daun salam (*Syzygium polyanthum*) terhadap larva *Aedes* sp. *Medical Laboratory Technology Journal*, 3(1), 17. <https://doi.org/10.31964/mltj.v3i1.151>

2 Fouque, F., & Reeder, J. C. (2019). Impact of past and on-going changes on climate and weather on vector-borne diseases transmission: A look at the evidence. *Infectious Diseases of Poverty*, 8(1), 1–9. <https://doi.org/10.1186/s40249-019-0565-1>

4 Gutierrez, P. M., Antepuesto, A. N., Eugenio, B. A. L., & Santos, M. F. L. (2014). Larvicidal activity of selected plant extracts against the dengue vector *Aedes aegypti* mosquito. *International Research Journal of Biological Sciences*, 3(4), 23–32.

Indah, I. S. (2019). *Situasi Filariasis di Indonesia*. Pusat Data dan Informasi Kementerian Kesehatan RI (p. 12). <https://pusdatin.kemkes.go.id/article/view/20010200002/infodatin-situasi-filariasis-di-indonesia.html>

5 Kumar, S., Mishra, M., Wahab, N., & Warikoo, R. (2014). Larvicidal, repellent, and irritant potential of the seed-derived essential oil of *Apium graveolens* against dengue vector, *Aedes aegypti* L. (Diptera: Culicidae). *Frontiers in Public Health*, 2(SEP), 1–6. <https://doi.org/10.3389/fpubh.2014.00147>

9 Mardiana, Supraptini, & Aminah, N. S. (2009). *Datura metel* Linnaeus sebagai insektisida dan larvasida botani serta bahan baku obat tradisional. *Media Penelitian dan Pengembangan Kesehatan*, XIX. <https://agris.fao.org/agris-search/search.do?recordID=DJ2020000225>

10 Pradani, F. Y. (2009). Indeks pertumbuhan larva *Aedes aegypti* L. yang terdedah dalam ekstrak air kulit jengkol (*Pithecellobium lobatum*). *Aspirator: Journal of Vector-borne Disease Studies*, 1(2). <https://ejournal2.litbang.kemkes.go.id/index.php/aspirator/issue/archive>

8 Sulistiyani, Falah, S., Wahyuni, W. T., Sugahara, T., Tachibana, S., & Syaefudin. (2014). Cellular mechanism of the cytotoxic effect of extracts from *Syzygium polyanthum* leaves. *American Journal of Drug Discovery and Development*, 4(2), 90–101. <https://doi.org/10.3923/ajdd.2014.90.101>

Rohmayani, et al. The Potency of Water Leaves Extract Of Bay...

Utami, P., & Puspaningtyas, D. E. (2014). *The Miracle Of Herbs*. Agro Media Pustaka.

Waskito, P. E., & Cahyati, W. H. (2018). Efektivitas granul daun salam (*Eugenia polyantha* Wight) sebagai larvasida nyamuk *Aedes aegypti*. *SPIRAKEL*, 10(1), 12–20. <http://ejournal2.litbang.kemkes.go.id/index.php/spirakel/article/view/546>