


**Dr. Irwan Syahrir**

# The Prevalence and Cost Impication of bloodstream Infection Producing

 Quick Submit

 Quick Submit

 Universitas Muhammadiyah Surabaya

---

## Document Details

Submission ID

trn:oid:::1:3328823471

Submission Date

Sep 4, 2025, 9:15 AM GMT+7

Download Date

Sep 4, 2025, 9:19 AM GMT+7

File Name

lence\_and\_Cost\_Impication\_of\_bloodstream\_Infection\_Producing.pdf

File Size

854.4 KB

7 Pages

2,773 Words

16,631 Characters





# 12% Overall Similarity

The combined total of all matches, including overlapping sources, for each database.




## Filtered from the Report

- Bibliography
- Quoted Text
- Cited Text
- Small Matches (less than 8 words)

## Match Groups

-  **24 Not Cited or Quoted 12%**  
Matches with neither in-text citation nor quotation marks
-  **0 Missing Quotations 0%**  
Matches that are still very similar to source material
-  **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 0%  Internet sources
- 12%  Publications
- 0%  Submitted works (Student Papers)

## Integrity Flags

### 0 Integrity Flags for Review

No suspicious text manipulations found.

Our system's algorithms look deeply at a document for any inconsistencies that would set it apart from a normal submission. If we notice something strange, we flag it for you to review.

A Flag is not necessarily an indicator of a problem. However, we'd recommend you focus your attention there for further review.

## Match Groups

- 24 Not Cited or Quoted** 12%  
Matches with neither in-text citation nor quotation marks
- 0 Missing Quotations** 0%  
Matches that are still very similar to source material
- 0 Missing Citation** 0%  
Matches that have quotation marks, but no in-text citation
- 0 Cited and Quoted** 0%  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 0% Internet sources
- 12% Publications
- 0% Submitted works (Student Papers)

## Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Publication	Annisa Fira Salsabila, Juniastuti, Dominicus Husada, Dwiyantri Puspitasari. "HEM...	2%
2	Publication	Jesús Rodríguez-Baño, Belén Gutiérrez-Gutiérrez, Isabel Machuca, Alvaro Pascual...	1%
3	Publication	M. Tumbarello, M. Sanguinetti, E. Montuori, E. M. Trecarichi et al. "Predictors of M...	1%
4	Publication	Lutifta Hilwana, Ninik Asmaningsih Soemyarso, Atika Atika. "Clinical Profile of Chi...	<1%
5	Publication	Brahmana Askandar Tjokroprawiro, Khoirunnisa Novitasari, Wita Saraswati, Indr...	<1%
6	Publication	"Abstracts of the 18th Congress of the International Pediatric Nephrology Associ...	<1%
7	Publication	Bernadette G. Pfang, Joaquín García-Cañete, Julia García-Lasheras, Antonio Blanc...	<1%
8	Publication	Rose M. Collis, Sara A. Burgess, Patrick J. Biggs, Anne C. Midwinter, Nigel P. Frenc...	<1%
9	Publication	Neissya Nastiti Firmanto, Maulydia, Pungky Mulawardhana, Mariza Fitriati. "Seve...	<1%
10	Publication	Patrick N.A. Harris, M. Diletta Pezzani, Belén Gutiérrez-Gutiérrez, Pierluigi Viale et...	<1%

11	Publication	Apichai Srijan, Katie R. Margulieux, Sirigade Ruekit, Erik Snestrud et al. " Genomic ...	<1%
12	Publication	F. Márquez-Friedrichs, M. B. Nolly, A. Ferreyra, L. Zuloaga et al. " Shifts in genes a...	<1%
13	Publication	"Uncovering the secrets of resistance: An introduction to computational methods...	<1%
14	Publication	Falagas, M.E.. "Extended-spectrum @b-lactamase-producing organisms", Journal ...	<1%
15	Publication	Lisa L Maragakís, Eli N Perencevich, Sara E Cosgrove. "Clinical and economic burd...	<1%
16	Publication	M. Tumbarello, M. Sali, E. M. Trecarichi, F. Leone et al. "Bloodstream Infections Ca...	<1%
17	Publication	Ton Duc Thang University	<1%



# QANUN MEDIKA

## JURNAL KEDOKTERAN FKUM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



### Research Article

## The Prevalence and Cost Implications of Bloodstream Infection Producing *Extended-Spectrum Beta-Lactamase (ESBL)* in Adult Patients at Dr. Soetomo General Academic Hospital

Neneng Dewi Kurniati<sup>1,2\*</sup>, Irwan Syahrir<sup>3,4</sup>, Musa Ghufro<sup>4</sup>

1) Hospital Administration Magister, Faculty of Medicine, Universitas Muhammadiyah Surabaya

2) Department of Clinical Microbiology, Dr. Soetomo General Academic Hospital, Surabaya

3) Senior Lecturer in Administration Magister Technic Faculty, Universitas Muhammadiyah Surabaya

4) Senior Lecturer in Administration Magister Medical Faculty, Universitas Muhammadiyah Surabaya

### ARTICLE INFO

Submitted : 17<sup>th</sup> February 2025

Accepted : 18<sup>th</sup> July 2025

Published : 25<sup>th</sup> July 2025

#### Keywords:

ESBL, BSI, MDRO, hospitalization

#### \*Correspondence:

neneng.dewi.kurniati-2023@

fk.um-surabaya.ac.id

This is an Open Access article under the CC-BY license



### ABSTRACT

Bloodstream infections (BSIs) caused by extended-spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae represent a growing concern in healthcare settings, particularly in tertiary hospitals. These infections are associated with limited therapeutic options, increased mortality, and rising healthcare costs. This study aimed to describe the clinical, microbiological, and economic characteristics of adult inpatients diagnosed with hospital-acquired (HA) BSIs due to ESBL-producing Enterobacteriaceae at Dr. Soetomo General Academic Hospital from January 1 to December 31, 2024. A cross-sectional study was conducted using retrospective data from adult inpatients ( $\geq 18$  years) with confirmed HA-BSIs caused by ESBL-producing *Escherichia coli*, *Klebsiella pneumoniae*, or *Proteus mirabilis*. Only the first positive blood culture per patient was analyzed. Community-acquired infections and non-Enterobacteriaceae organisms were excluded. The most frequent pathogen was *Escherichia coli* ESBL, especially among patients from medical wards. BSIs occurred predominantly in medical and intensive care units. Notably, microbiological confirmation of ESBL-producing organisms led to higher claim reimbursements through the national health insurance system, yielding a positive cost margin despite longer hospital stays and expensive treatment regimens. In conclusion, ESBL-producing Enterobacteriaceae remain a major challenge in managing HA-BSIs at Dr. Soetomo General Academic Hospital. While clinically burdensome, accurate microbiological diagnosis also supports optimal resource utilization and insurance claim efficiency in the hospital setting.



# QANUN MEDIKA

## JURNAL KEDOKTERAN FKUM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



### INTRODUCTION

Bloodstream infections (BSIs) remain a significant cause of morbidity and mortality among hospitalized patients worldwide, especially in critical care and immunocompromised populations. The emergence and spread of antimicrobial-resistant organisms, particularly extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae*, have significantly complicated the clinical management of BSIs. ESBL enzymes confer resistance to a broad range of beta-lactam antibiotics, including penicillins, third-generation cephalosporins, and aztreonam. This limits treatment options and increases dependence on last-line antibiotics such as carbapenems (CDC, 2021).

The global prevalence of ESBL-producing pathogens has been rising, with marked increases reported in both community and healthcare settings, especially in low- and middle-income countries. Hospital-acquired (HA) BSIs caused by ESBL-producing organisms are particularly concerning due to their association with prolonged hospital stays, increased healthcare costs, higher rates of intensive care unit (ICU) admission, and increased mortality (Schwaber & Carmeli, 2007; Jean et al., 2022). Identifying these infections' clinical and epidemiological characteristics is critical for improving infection control, guiding empirical antibiotic therapy, and informing antimicrobial stewardship strategies (NICE, 2016).

Despite the growing clinical importance of ESBL-BSIs, there is limited data specific to Indonesian tertiary care centers, including Dr. Soetomo General Academic Hospital. This study aims to characterize the microbiological features of adult inpatients diagnosed with hospital-acquired bloodstream infections due to ESBL-producing *Enterobacteriaceae*

and the cost implications over one year in Dr. Soetomo General Academic Hospital, Surabaya.

### METHODS

This was a retrospective cohort study conducted at Dr. Soetomo General Academic Hospital, a tertiary care and teaching hospital in Surabaya, Indonesia. The study period covered one full calendar year, from 1st January to 31st December 2024. The study aimed to describe the characteristics of adult inpatients with hospital-acquired (HA) bloodstream infections (BSIs) caused by extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae*.

Data were obtained from hospitalized adult patients ( $\geq 18$  years old) who developed a laboratory-confirmed BSI due to *Enterobacteriaceae* during their admission. Only HA-BSI cases were included in the analysis. A BSI was classified as hospital-acquired if the first positive blood culture was obtained  $\geq 48$  hours after hospital admission. The date of BSI onset was defined as the date of collection of the first blood sample that yielded an *Enterobacteriaceae* isolate.

Patients were excluded from the study if the BSI was classified as community-acquired, defined as detection of BSI within the first 48 hours of admission. In addition, BSIs caused by non-*Enterobacteriaceae* pathogens were excluded. If the same *Enterobacteriaceae* strain was isolated more than once in the same patient, only the first isolate was included in the analysis to avoid duplication.

Microbiological data were collected from the hospital's clinical microbiology laboratory records. Blood cultures were processed according to standard procedures, and identification of *Enterobacteriaceae* and ESBL production was performed using the BD Phoenix automated system and confirmed by



phenotypic confirmatory tests according to Clinical and Laboratory Standards Institute (CLSI) guidelines.

The information, including ward of hospitalization and medical cost, was extracted from electronic medical records. Data were analyzed using descriptive statistics to determine the frequency and distribution of ESBL-producing *Enterobacteriaceae* BSI characteristics during the study period. This study has been ethically approved by the Ethics Committee of Dr. Soetomo General Academic Hospital with reference number 1294/KEPK/IV/2025.

## RESULTS

Table 1 shows the extended-spectrum beta-lactamase (ESBL)-producing *Enterobacteriaceae* that caused hospital-acquired (HA) bloodstream infections (BSIs) in Dr. Soetomo General Academic Hospital in 2024 were *Escherichia coli* ESBL, *Klebsiella pneumoniae* ESBL, and *Proteus mirabilis* ESBL. The total of these isolates was 170.

**Table 1.** Total ESBL bacteria

PATHOGENS	ESBL (%)	Number of isolates
<i>Escherichia coli</i> ESBL	61.53 %	105
<i>Klebsiella pneumoniae</i> ESBL	33.13 %	57
<i>Proteus mirabilis</i> ESBL	4.75 %	8
Total	100	170

**Table 2.** Distribution of ESBL Bacteria

ROOMS	AMOUNTS	%
<b>Emergency Department</b>	<b>12</b>	<b>8.29</b>
<i>Klebsiella pneumoniae</i> ESBL	4	2.37
<i>Escherichia coli</i> ESBL	8	5.92
<b>Surgery Wards</b>	<b>11</b>	<b>6.51</b>
<i>Klebsiella pneumoniae</i> ESBL	3	1.78
<i>Escherichia coli</i> ESBL	7	4.14
<i>Proteus mirabilis</i> ESBL	1	0.59
<b>Medical Wards</b>	<b>72</b>	<b>42.60</b>
<i>Klebsiella pneumoniae</i> ESBL	24	13.60
<i>Escherichia coli</i> ESBL	45	26.63
<i>Proteus mirabilis</i> ESBL	3	1.78
<b>Intensive Care Unit</b>	<b>72</b>	<b>42.60</b>
<i>Klebsiella pneumoniae</i> ESBL	26	15.38
<i>Escherichia coli</i> ESBL	42	24.85
<i>Proteus mirabilis</i> ESBL	4	2.37
<b>Total</b>	<b>170</b>	<b>100</b>





# QANUN MEDIKA

## JURNAL KEDOKTERAN FKUM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



**Table 3. Distribution of Loss and Profit by ESBL Bacteria Type**

ROOMS	LOSS (Patients)	PROFIT (Patients)
<b>Emergency Department</b>	<b>5</b>	<b>7</b>
<i>Escherichia coli ESBL</i>	5	3
<i>Klebsiella pneumoniae ESBL</i>	0	4
<b>EMERGENCY DEPARTMENT INTENSIVE ROOM</b>	<b>16</b>	<b>27</b>
<i>Escherichia coli ESBL</i>	12	22
<i>Klebsiella pneumoniae ESBL</i>	3	3
<i>Proteus mirabilis ESBL</i>	1	2
<b>SURGERY WARDS</b>	<b>1</b>	<b>10</b>
<i>Escherichia coli ESBL</i>	0	7
<i>Klebsiella pneumoniae ESBL</i>	0	3
<i>Proteus mirabilis ESBL</i>	1	
<b>MEDICAL WARDS</b>	<b>20</b>	<b>52</b>
<i>Escherichia coli ESBL</i>	12	33
<i>Klebsiella pneumoniae ESBL</i>	7	17
<i>Proteus mirabilis ESBL</i>	1	2
<b>INTENSIVE CARE UNIT</b>	<b>2</b>	<b>30</b>
<i>Escherichia coli ESBL</i>	2	9
<i>Klebsiella pneumoniae ESBL</i>	0	20
<i>Proteus mirabilis ESBL</i>	0	1
<b>Total Patients</b>	<b>44</b>	<b>126</b>

Table 2 shows the most common distribution of ESBL-producing *Enterobacteriaceae* that caused HA-BSIs from medical wards and intensive care units. The most common ESBL-producing *Enterobacteriaceae* that caused HA-BSIs were *Escherichia coli ESBL* in medical wards.

Table 3 shows the distribution of medical costs' loss and profit by ESBL-producing *Enterobacteriaceae* that caused HA-BSIs in Dr. Soetomo General Academic Hospital in 2024. It shows that the microbiological culture that resulted in ESBL-producing *Enterobacteriaceae* from HA-BSIs gave more profit in the claim of medical insurance.

## DISCUSSION

This study highlights the epidemiological and clinical characteristics of adult inpatients diagnosed with bloodstream infections (BSIs) caused by extended-spectrum beta-lactamase (ESBL)-producing organisms at Dr. Soetomo General Academic Hospital. The high incidence of ESBL-producing pathogens, particularly *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis*, reflects an ongoing challenge in managing hospital-acquired (HA) infections in tertiary care settings (Lee et al., 2019; Rodríguez-Baño et al., 2018). These infections predominantly originated from medical wards and intensive care units (ICUs), where patients often present with complex comorbidities and are exposed to broad-spectrum antibiotics and





# QANUN MEDIKA

## JURNAL KEDOKTERAN FKUM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



invasive procedures (Yap et al., 2021). Notably, the most frequent isolate was *Escherichia coli* ESBL, especially in patients from medical wards, indicating a possible link to urinary tract-related bloodstream infections or post-abdominal surgery complications (Russo & Johnson, 2003).

The presence of ESBL-producing bacteria significantly complicates BSI management due to resistance to third-generation cephalosporins and monobactams, necessitating the use of last-resort antibiotics such as carbapenems (Paterson & Bonomo, 2005; Bush & Bradford, 2016). Although most isolates in this study remained carbapenem-susceptible, reliance on these agents raises concerns about the rise of carbapenem-resistant *Enterobacteriaceae* (CRE) (Meletis, 2016; Tamma et al., 2021).

Compared with regional data, the prevalence of ESBL-BSIs in this study is consistent with findings from other tertiary hospitals in Southeast Asia (Jean et al., 2022). However, antimicrobial resistance patterns may vary due to differences in antimicrobial use, diagnostic resources, and infection control policies (World Health Organization, 2017). Continuous local surveillance and antimicrobial stewardship are essential to ensure empirical treatment remains effective and to reduce selective pressure for resistance (Tamma et al., 2021; Indonesian Ministry of Health, 2023).

Economically, ESBL-related infections are known to increase healthcare resource use, including longer hospital stays and higher drug costs (Russo & Johnson, 2003). Interestingly, analysis of national health insurance claims at Dr. Soetomo General Academic Hospital in 2024 revealed that HA-BSI cases caused by ESBL-producing *Enterobacteriaceae* yielded higher reimbursement due to classification as high-severity cases (Direktorat Jenderal Pelayanan Kesehatan, 2022). This allowed

hospitals to claim for comprehensive diagnostics and advanced antimicrobial treatments. Consequently, although ESBL infections impose a greater clinical burden, they also result in a net positive hospital margin due to successful claim approvals, demonstrating the dual value of microbiological confirmation (Ritchie & Roser, 2019). These findings reinforce the importance of infection control bundles, including hand hygiene, antimicrobial de-escalation, catheter care, and early removal of invasive devices (Kaye et al., 2021; Indonesian Ministry of Health, 2023). Special attention should be paid to high-risk units like medical wards and ICUs, where ESBL-producing *Enterobacteriaceae* are most frequently isolated.

This study has several limitations, including its retrospective design and single-center scope, which limit generalizability. Additionally, molecular analysis of ESBL genes was not performed, which could have offered further insights into resistance mechanisms and transmission (Rodríguez-Baño et al., 2018). Nonetheless, this study contributes meaningful data on the clinical, microbiological, and economic aspects of ESBL-BSI in a high-volume Indonesian tertiary hospital. Future prospective multicenter studies are needed to assess molecular resistance patterns and evaluate cost-effective prevention strategies in similar healthcare settings.

## CONCLUSION

Hospital-acquired bloodstream infections caused by ESBL-producing *Enterobacteriaceae* remain a significant clinical and economic concern at Dr. Soetomo General Academic Hospital. In 2024, the predominant pathogens were *Escherichia coli*, *Klebsiella pneumoniae*, and *Proteus mirabilis*, with *E. coli* ESBL being most commonly isolated from medical ward



# QANUN MEDIKA

## JURNAL KEDOKTERAN FKUM SURABAYA

<http://journal.um-surabaya.ac.id/index.php/qanunmedika>



patients. These infections were associated with prolonged hospitalization and complex clinical management, yet they also generated higher insurance claim reimbursements due to their classification as severe cases requiring intensive diagnostics and high-cost treatments. Strengthening infection control measures, antimicrobial stewardship, and early microbiological diagnostics is essential to mitigate the burden of ESBL-related BSIs while optimizing both clinical outcomes and hospital resource utilization.

### REFERENCES

- Bush, K., & Bradford, P. A. (2016).  $\beta$ -Lactams and  $\beta$ -lactamase inhibitors: An overview. *Cold Spring Harbor Perspectives in Medicine*, 6(8), a025247. <https://doi.org/10.1101/cshperspect.a025247>
- Busse, R. (2012). Do diagnosis-related groups explain variations in hospital costs and length of stay? - Analyses from the EURODRG project for 10 episodes of care across 10 European countries. *Health Economics*, 21(Suppl.2), 1–5. <https://doi.org/10.1002/hec>
- CDC. (2021). Background of the Management of Multidrug- Resistant Organism in Healthcare Settings Guideline, <https://www.cdc.gov/infection-control/hcp/mdro-management/background.html>
- Direktorat Jenderal Pelayanan Kesehatan Kemenkes RI. (2022). Panduan pengajuan klaim Jaminan Kesehatan Nasional (JKN) untuk RS rujukan kelas A dan B. Jakarta: Kemenkes RI.
- Indonesian Ministry of Health. (2023). Pedoman Nasional Pengendalian Resistensi Antimikroba di Rumah Sakit. Jakarta: Kemenkes RI.
- Jean, S. S., Hsueh, P. R., Lee, W. S., & Yu, C. J. (2022). Hospital-acquired bloodstream infections: Current trends and challenges in the Asia-Pacific region. *Journal of Global Antimicrobial Resistance*, 29, 321–329. <https://doi.org/10.1016/j.jgar.2021.08.011>
- Kaye, K. S., Pogue, J. M., Tran, T. B., Nation, R. L., & Li, J. (2021). Agents of last resort: The role of polymyxins and newer antibiotics in treating resistant Gram-negative infections. *Current Opinion in Infectious Diseases*, 34(6), 519–528. <https://doi.org/10.1097/QCO.0000000000000792>
- Lee, C. H., Lee, M. H., Ko, W. C., & Chang, C. M. (2019). Epidemiology and clinical significance of bloodstream infections caused by ESBL-producing Enterobacteriaceae. *Journal of Microbiology, Immunology and Infection*, 52(4), 515–524. <https://doi.org/10.1016/j.jmii.2018.12.005>
- Meletis, G. (2016). Carbapenem resistance: Overview of the problem and future perspectives. *Therapeutic Advances in Infectious Disease*, 3(1), 15–21. <https://doi.org/10.1177/2049936115621709>
- NICE. (2016). Suspected sepsis: recognition, diagnosis and early management. *NICE guideline*. [www.nice.org.uk/guidance/ng51](http://www.nice.org.uk/guidance/ng51).
- Paterson, D. L., & Bonomo, R. A. (2005). Extended-spectrum  $\beta$ -lactamases: A clinical update. *Clinical Microbiology Reviews*, 18(4), 657–686. <https://doi.org/10.1128/CMR.18.4.657-686.2005>
- Ritchie, H., & Roser, M. (2019). Causes of death. *Our World in Data*. <https://ourworldindata.org/causes-of-death>
- Rodríguez-Baño, J., Gutiérrez-Gutiérrez, B., Machuca, I., & Pascual, Á. (2018).



- Treatment of infections caused by extended-spectrum-beta-lactamase-, AmpC-, and carbapenemase-producing Enterobacteriaceae. *Clinical Microbiology Reviews*, 31(2), e00079-17. <https://doi.org/10.1128/CMR.00079-17>
- Russo, T. A., & Johnson, J. R. (2003). Medical and economic impact of extraintestinal infections due to *Escherichia coli*: Focus on an increasingly important endemic problem. *Microbes and Infection*, 5(5), 449–456. [https://doi.org/10.1016/S1286-4579\(03\)00049-2](https://doi.org/10.1016/S1286-4579(03)00049-2)
- Schwaber, M. J., & Carmeli, Y. (2007). Mortality and delay in effective therapy associated with extended-spectrum  $\beta$ -lactamase production in Enterobacteriaceae bacteraemia: A systematic review and meta-analysis. *Journal of Antimicrobial Chemotherapy*, 60(5), 913–920. <https://doi.org/10.1093/jac/dkm318>
- Tamma, P. D., Aitken, S. L., Bonomo, R. A., Mathers, A. J., van Duin, D., & Clancy, C. J. (2021). Infectious Diseases Society of America Guidance on the treatment of extended-spectrum  $\beta$ -lactamase-producing Enterobacterales (ESBL-E), carbapenem-resistant Enterobacterales (CRE), and *Pseudomonas aeruginosa* with difficult-to-treat resistance (DTR-P. *aeruginosa*). *Clinical Infectious Diseases*, 72(7), e169–e183. <https://doi.org/10.1093/cid/ciaa1478>
- World Health Organization. (2017). Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. <https://www.who.int/publications/i/item/WHO-EMP-IAU-2017.12>
- Yap, P. S. X., Ahmad Kamar, A., Chong, C. W., & Yap, I. K. S. (2021). Risk factors of bloodstream infections caused by multidrug-resistant organisms in Asia. *Infectious Disease Reports*, 13(1), 80–91. <https://doi.org/10.3390/idr13010010>