



**DAFTAR PUSTAKA**

## DAFTAR PUSTAKA

- Arefian, H. *et al.* (2017) 'Hospital-related cost of sepsis: A systematic review', *Journal of Infection*, 74(2), pp. 107–117. Available at: <https://doi.org/10.1016/j.jinf.2016.11.006>.
- Arora, J., Mendelson, A.A. and Fox-Robichaud, A. (2023) 'Sepsis: network pathophysiology and implications for early diagnosis', *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 324(5), pp. R613–R624. Available at: <https://doi.org/10.1152/ajpregu.00003.2023>.
- Barbara, P. *et al.* (2018) 'The quick sequential organ failure assessment (qSOFA) identifies septic patients in the out-of-hospital setting', *The American Journal of Emergency Medicine*, 36(6), pp. 1022–1026. Available at: <https://doi.org/https://doi.org/10.1016/j.ajem.2018.01.073>.
- Bou Chebl, R. *et al.* (2022) 'The association between the neutrophil to lymphocyte ratio and in-hospital mortality among sepsis patients: A prospective study', *Medicine*, 101, p. e29343. Available at: <https://doi.org/10.1097/MD.00000000000029343>.
- Cowland, J.B. and Borregaard, N. (2016) 'Granulopoiesis and granules of human neutrophils', *Immunological Reviews*, 273(1), pp. 11–28. Available at: <https://doi.org/10.1111/imr.12440>.
- Darkwah, S. *et al.* (2024) 'Sepsis-Related Lung Injury and the Complication of Extrapulmonary Pneumococcal Pneumonia', *Diseases*, 12(4), p. 72. Available at: <https://doi.org/10.3390/diseases12040072>.
- Djuang, M.H. *et al.* (2022) *FAKTOR RESIKO SEPSIS PADA PASIEN LANSIA DI RUMAH SAKIT UMUM ROYAL PRIMA MEDAN SEPSIS RISK FACTORS IN ELDERLY PATIENTS AT ROYAL PRIMA MEDAN GENERAL HOSPITAL*. Oktober. Available at: <https://ejournal.ung.ac.id/index.php/jjhsr/index>.
- Dugani, S., Veillard, J. and Kissoon, N. (2017) 'Reducing the global burden of sepsis', *Canadian Medical Association Journal*, 189(1), pp. E2–E3. Available at: <https://doi.org/10.1503/cmaj.160798>.
- Evans, L. *et al.* (2021) 'Surviving sepsis campaign: international guidelines for management of sepsis and septic shock 2021', *Intensive Care Medicine*, 47(11), pp. 1181–1247. Available at: <https://doi.org/10.1007/s00134-021-06506-y>.
- Fisher, A. *et al.* (2016) 'The Neutrophil to Lymphocyte Ratio on Admission and Short-Term Outcomes in Orthogeriatric Patients', *International Journal of Medical Sciences*, 13(8), pp. 588–602. Available at: <https://doi.org/10.7150/ijms.15445>.
- Gauer, R., Forbes, D., & Boyer, N. (2020). Sepsis: Diagnosis and Management. *American family physician*, 101(7), 409–418.
- Hotchkiss, R.S. *et al.* (2016) 'Sepsis and septic shock', *Nature Reviews Disease Primers*, 2(1), p. 16045. Available at: <https://doi.org/10.1038/nrdp.2016.45>.

- Hsieh, M.-S. *et al.* (2024) 'Utilizing the National Early Warning Score 2 (NEWS2) to confirm the impact of emergency department management in sepsis patients: a cohort study from taiwan 1998–2020', *International Journal of Emergency Medicine*, 17(1), p. 42. Available at: <https://doi.org/10.1186/s12245-024-00614-4>.
- Huang, Z. *et al.* (2020) 'Prognostic value of neutrophil-to-lymphocyte ratio in sepsis: A meta-analysis', *The American Journal of Emergency Medicine*, 38(3), pp. 641–647. Available at: <https://doi.org/10.1016/j.ajem.2019.10.023>.
- Ince, C. *et al.* (2016) 'The Endothelium in Sepsis', *Shock*, 45(3), pp. 259–270. Available at: <https://doi.org/10.1097/SHK.0000000000000473>.
- de Jager, C.P.C. *et al.* (2012) 'The Neutrophil-Lymphocyte Count Ratio in Patients with Community-Acquired Pneumonia', *PLoS ONE*, 7(10), p. e46561. Available at: <https://doi.org/10.1371/journal.pone.0046561>.
- Jarczak, D., Kluge, S. and Nierhaus, A. (2021) 'Sepsis—Pathophysiology and Therapeutic Concepts', *Frontiers in Medicine*, 8. Available at: <https://doi.org/10.3389/fmed.2021.628302>.
- Jevon, P. and Shamsi, S. (2020) 'Using National Early Warning Score (NEWS) 2 to help manage medical emergencies in the dental practice', *British Dental Journal*, 229(5), pp. 292–296. Available at: <https://doi.org/10.1038/s41415-020-2024-6>.
- Jiang, J. *et al.* (2019) 'Nonviral infection-related lymphocytopenia for the prediction of adult sepsis and its persistence indicates a higher mortality', *Medicine*, 98(29), p. e16535. Available at: <https://doi.org/10.1097/MD.00000000000016535>.
- Jung, M. *et al.* (2022) 'Analysis of Infection Transmission Routes through Exhaled Breath and Cough Particle Dispersion in a General Hospital', *International Journal of Environmental Research and Public Health*, 19(5). Available at: <https://doi.org/10.3390/ijerph19052512>.
- Kaushik, R. *et al.* (2018) 'Diagnostic and prognostic role of neutrophil-to-lymphocyte ratio in early and late phase of sepsis', *Indian Journal of Critical Care Medicine*, 22(9), pp. 660–663. Available at: [https://doi.org/10.4103/ijccm.IJCCM\\_59\\_18](https://doi.org/10.4103/ijccm.IJCCM_59_18).
- Kita, Y. *et al.* (2015) 'Fever is mediated by conversion of endocannabinoid 2-arachidonoylglycerol to prostaglandin E2', *PLoS ONE*, 10(7). Available at: <https://doi.org/10.1371/journal.pone.0133663>.
- Kurniati, I., Arisqan, F.S. and Mutiara, U.G. (2023) 'Differences in Neutrophil Lymphocyte Ratio (NLR) between Sepsis and Septic Shock Patients in a Tertiary Hospital in Indonesia', *Majalah Kedokteran Bandung*, 55(3), pp. 149–153. Available at: <https://doi.org/10.15395/mkb.v55n3.3149>.
- Kuswoyo, N.H. and Natadidjaja, R.I. (2019) 'Korelasi antara kadar procalcitonin dengan serum transaminase pada pasien sepsis: sebuah studi pendahuluan',

*Jurnal Biomedika dan Kesehatan*, 2(1), pp. 15–19. Available at: <https://doi.org/10.18051/JBiomedKes.2019.v2.15-19>.

- Lahoz-Beneytez, J. *et al.* (2016) ‘Human neutrophil kinetics: modeling of stable isotope labeling data supports short blood neutrophil half-lives’, *Blood*, 127(26), pp. 3431–3438. Available at: <https://doi.org/10.1182/blood-2016-03-700336>.
- Lambden, S. *et al.* (2019) ‘The SOFA score—development, utility and challenges of accurate assessment in clinical trials’, *Critical Care*, 23(1), p. 374. Available at: <https://doi.org/10.1186/s13054-019-2663-7>.
- Levi, M., Scully, M. and Singer, M. (2018) ‘The role of ADAMTS-13 in the coagulopathy of sepsis’, *Journal of Thrombosis and Haemostasis*, 16(4), pp. 646–651. Available at: <https://doi.org/10.1111/jth.13953>.
- Liu, X. *et al.* (2016) ‘Prognostic Significance of Neutrophil-to-Lymphocyte Ratio in Patients with Sepsis: A Prospective Observational Study’, *Mediators of Inflammation*, 2016, pp. 1–8. Available at: <https://doi.org/10.1155/2016/8191254>.
- Orakpoghenor, O. *et al.* (2019) ‘Lymphocytes: A Brief Review’, 3, pp. 4–8.
- Pantzaris, N.-D. *et al.* (2018) ‘Neutrophil-to-lymphocyte ratio relation to sepsis severity scores and inflammatory biomarkers in patients with community-acquired pneumonia: A case series’, *Journal of Translational Internal Medicine*, 6(1), pp. 43–46. Available at: <https://doi.org/10.2478/jtim-2018-0009>.
- Park, S. *et al.* (2020) ‘Normothermia in Patients With Sepsis Who Present to Emergency Departments Is Associated With Low Compliance With Sepsis Bundles and Increased In-Hospital Mortality Rate\*’, *Critical Care Medicine*, 48(10), pp. 1462–1470. Available at: <https://doi.org/10.1097/CCM.0000000000004493>.
- Rada, B. (2019) ‘Neutrophil Extracellular Traps’, in, pp. 517–528. Available at: [https://doi.org/10.1007/978-1-4939-9424-3\\_31](https://doi.org/10.1007/978-1-4939-9424-3_31).
- Riché, F. *et al.* (2015) ‘Reversal of neutrophil-to-lymphocyte count ratio in early versus late death from septic shock’, *Critical Care*, 19(1), p. 439. Available at: <https://doi.org/10.1186/s13054-015-1144-x>.
- Rosales, C. (2018) ‘Neutrophil: A Cell with Many Roles in Inflammation or Several Cell Types?’, *Frontiers in Physiology*, 9. Available at: <https://doi.org/10.3389/fphys.2018.00113>.
- Royal College of Physicians (2017) *National Early Warning Score (NEWS) 2 Standardising the assessment of acute-illness severity in the NHS*. Available at: [www.rcplondon.ac.uk](http://www.rcplondon.ac.uk).
- Rubio, I. *et al.* (2019) ‘Current gaps in sepsis immunology: new opportunities for translational research’, *The Lancet Infectious Diseases*, 19(12), pp. e422–e436. Available at: [https://doi.org/10.1016/S1473-3099\(19\)30567-5](https://doi.org/10.1016/S1473-3099(19)30567-5).
- Rudd, K.E. *et al.* (2020) ‘Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study’,

- The Lancet*, 395(10219), pp. 200–211. Available at: [https://doi.org/10.1016/S0140-6736\(19\)32989-7](https://doi.org/10.1016/S0140-6736(19)32989-7).
- Sakr, Y. *et al.* (2013) ‘The influence of gender on the epidemiology of and outcome from severe sepsis’, *Critical Care*, 17(2), p. R50. Available at: <https://doi.org/10.1186/cc12570>.
- Saliccioli, J.D. *et al.* (2015) ‘The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: an observational cohort study’, *Critical Care*, 19(1), p. 13. Available at: <https://doi.org/10.1186/s13054-014-0731-6>.
- Schupp, T. *et al.* (2023) ‘The Neutrophil-to-Lymphocyte-Ratio as Diagnostic and Prognostic Tool in Sepsis and Septic Shock’, *Clinical Laboratory*, 69(05/2023). Available at: <https://doi.org/10.7754/Clin.Lab.2022.220812>.
- Seymour, C.W. *et al.* (2016) ‘Assessment of Clinical Criteria for Sepsis’, *JAMA*, 315(8), p. 762. Available at: <https://doi.org/10.1001/jama.2016.0288>.
- Singer, M. *et al.* (2016) ‘The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)’, *JAMA*, 315(8), p. 801. Available at: <https://doi.org/10.1001/jama.2016.0287>.
- Song, M. *et al.* (2021) ‘Neutrophil-to-lymphocyte ratio and mortality in the United States general population’, *Scientific Reports*, 11(1), p. 464. Available at: <https://doi.org/10.1038/s41598-020-79431-7>.
- Tang, Y. *et al.* (2017) ‘Systolic blood pressure variability in patients with early severe sepsis or septic shock: a prospective cohort study’, *BMC Anesthesiology*, 17(1), p. 82. Available at: <https://doi.org/10.1186/s12871-017-0377-4>.
- Tigner, A., Ibrahim, S. and Murray, I. (2020) ‘Histology, White Blood Cell’, in: World Health Organization (2020) *Current evidence, identifying gaps and future directions GLOBAL REPORT ON THE EPIDEMIOLOGY AND BURDEN OF SEPSIS*. Available at: <http://apps.who.int/bookorders>.
- Ye, W. *et al.* (2020) ‘The association between neutrophil-to-lymphocyte count ratio and mortality in septic patients: a retrospective analysis of the MIMIC-III database’, *Journal of Thoracic Disease*, 12(5), pp. 1843–1855. Available at: <https://doi.org/10.21037/jtd-20-1169>.
- Zahorec, R. (2021) ‘Neutrophil-to-lymphocyte ratio, past, present and future perspectives’, *Bratislava Medical Journal*, 122(07), pp. 474–488. Available at: [https://doi.org/10.4149/BLL\\_2021\\_078](https://doi.org/10.4149/BLL_2021_078).
- Zorio, V. *et al.* (2017) ‘Assessment of sepsis-induced immunosuppression at ICU discharge and 6 months after ICU discharge’, *Annals of Intensive Care*, 7(1), p. 80. Available at: <https://doi.org/10.1186/s13613-017-0304-3>.