



# DAFTAR PUSTAKA



## DAFTAR PUSTAKA

- Adisasmito, A.W., & Hadinegoro, S.R.S. (2016). 'Infeksi bakteri Gram negatif di ICU anak: Epidemiologi, manajemen antibiotik dan pencegahan'. *Sari Pediatri*, 6(1), pp. 32-35.
- Aliviameita, A. & Puspitasari, P., 2024. *Pemeriksaan Hematologi Rutin*. Sidoarjo: Umsida Press.
- Anies, A. (2016). 'Bakteremia: Definisi, Diagnosis, dan Penanganan'. *Jurnal Kesehatan Masyarakat*, 10(2), pp. 123-130.
- Atmawati, T.U., Faisal, M.A. & Rahmiati, R. (2017). 'Pola kepekaan isolat bakteri aerob pada konjungtivitis terhadap antibiotik terpilih di poliklinik mata rsud ulin Banjarmasin'. *Berkala Kedokteran*, 13(1), pp. 15.
- Balkhair, A., Saadi, K.A. & Adawi, B.A. (2023). 'Epidemiology and mortality outcome of carbapenem- and colistin-resistant *Klebsiella pneumoniae*, *Escherichia coli*, *Acinetobacter baumannii*, and *Pseudomonas aeruginosa* bloodstream infections', *IJID Regions*, 7, pp. 1–5.
- Buonacera, A., Stancanelli, B., Colaci, M., Malatino, L., & Neri, S. (2022). 'Neutrophil to lymphocyte ratio: An emerging marker of systemic inflammation'. *Clinical and Experimental Medicine*, 22, pp. 1–11.
- Bozdemir, Ş. E., Altıntop, Y. A., Uytun, S., Aslaner, H., Torun, Y. A. (2017). 'Diagnostic role of mean platelet volume and neutrophil to lymphocyte ratio in childhood brucellosis'. *Korean J Intern Med*, 32(6), pp. 1054-1060.
- Brendel, M., Köhler, T. P., Neufend, J. V., Puppe, A., Gisch, N., & Hammerschmidt, S. (2024). 'Lipoteichoic Acids Are Essential for Pneumococcal Colonization and Membrane Integrity'. *Journal of Innate Immunity*, 16(1), 370–384.
- Cahyaningtyas, D. E., Gaina, C. D., & Tangkonda, E. (2024). 'Isolasi dan identifikasi bakteri *Escherichia coli*, *Klebsiella* sp., dan *Staphylococcus aureus* pada ambing dan susu kambing Peranakan Etawa'. *Jurnal Veteriner Nusantara*, 7(1), pp. 41–52.
- Chen, J., Li, H., & Wang, Y. (2020). 'Neutrophil-to-Lymphocyte Ratio as a Predictor of Clinically Significant Bacterial Infections: Threshold Value and Clinical Utility'. *Journal of Infection and Public Health*, 13(11), pp. 1690–1697.
- European Centre for Disease Prevention and Control (ECDC)., 2019. *Annual epidemiological report for 2019: Healthcare-associated infections acquired in intensive care units*. Stockholm: ECDC
- Gomes, L.T., Morato-Conceição, Y.T., Gambati, A.V.M., Maciel-Pereira, C.M. & Fontes, C.J.F. (2020). 'Diagnostic value of neutrophil-to-lymphocyte ratio in patients with leprosy reactions'. *Heliyon*, 6(2), e03369.

- Gonzalez MD, Chao T, Pettengill MA. (2020). 'Modern Blood Culture: Management Decisions and Method Options'. *Clin Lab Med*, 40(4), pp. 605-617
- Gürol, G., Çiftci, İ. H., Terizi, H. A., Atasoy, A. R., Özbek, A., & Köroğlu, M. (2015). 'Are there standardized cutoff values for neutrophil-lymphocyte ratios in bacteremia or sepsis?'. *Journal of Microbiology and Biotechnology Research*, 5(4), 50–53.
- Hatman, F.A., Semedi, B.P. & Budiono, B. (2021). 'Analisis Faktor Risiko terhadap Lama Perawatan Pasien Sepsis yang Meninggal di Ruang Perawatan Intensif RSUD Dr. Soetomo Surabaya'. *JAI (Jurnal Anestesiologi Indonesia)*, 13, pp. 78–87.
- Hayati, Z., Widyastuti, E., Nurjannah, N., Mudatsir, M. & Saputra, I. (2022). 'Hubungan Kualitas Penggunaan Antibiotik dengan Luaran Klinis Pasien Bakteremia yang Disebabkan Methicillin Resistant Staphylococcus aureus (MRSA)'. *Jurnal Kedokteran Syiah Kuala*, 22(1), pp. 37-43.
- Hoffbrand, A.V., Moss, P.A.H. and Pettit, J.E., 2019. *Essential Haematology 8th ed.* Hoboken, NJ: Wiley-Blackwell
- Holmes, C.L., Anderson, M.T., Mobley, H.L.T. and Bachman, M.A. (2021). 'Pathogenesis of Gram-negative bacteremia'. *Clinical Microbiology Reviews*, 34(2), e00103-20.
- Hotchkiss, R. S., *et al.* (2016). 'Sepsis-induced immunosuppression'. *Nature Reviews Immunology*, 13(12), pp. 862–874.
- IDAI., 2016. *Pedoman Nasional Pelayanan Kedokteran IDAI Diagnosis dan Tatalaksana Sepsis Pada Anak*. Jakarta: IDAI
- Jones, L.K., White, M.S. & Green, T.H. (2019). 'Effects of Long-Term Corticosteroid Therapy on Peripheral Blood Neutrophil and Lymphocyte Counts'. *Immunology Letters*, 215, pp. 40-47.
- Kaukonen, K.-M., Bailey, M., Pilcher, D., Cooper, D.D. & Bellomo, R. (2018). 'The systemic inflammatory response syndrome criteria and their differential association with mortality'. *Journal of Critical Care*, 46, pp. 29–36.
- Kaushik, R., Gupta, M., Sharma, M., Jash, D., Jain, N., Sinha, N., Chaudhry, A., & Chaudhry, D. (2018). 'Diagnostic and Prognostic Role of Neutrophil-to-Lymphocyte Ratio in Early and Late Phase of Sepsis'. *Indian Journal of Critical Care Medicine*, 22(9), pp. 660–663.
- Kescandra, N., Rosa, E.M. & Sundari, S. (2020). 'E-Surveilans Bloodstream Infection pada Pasien di Nitipuran Health Centre'. *Jurnal Ilmiah Universitas Batanghari Jambi*, 20(1), pp. 263.
- Kumar, S., Raina, S., & Mehta, A. (2021). 'Neutrophil-to-lymphocyte ratio in viral infections: Clinical significance and diagnostic implications'. *Infectious Diseases Journal*, 29(2), pp. 112–119.
- Lagunas-Rangel, F. A. (2020). 'Neutrophil-to-lymphocyte ratio and lymphocyte-to-CRP ratio in patients with severe infections: A meta-analysis'. *Journal of Medical Virology*, 92(10), pp. 1733–1734.

- Lamy, B., Dargère, S., Arendrup, M.C., Parienti, J.-J. and Tattevin, P. (2016). 'How to optimize the use of blood cultures for the diagnosis of bloodstream infections? A state-of-the-art'. *Frontiers in Microbiology*, 7, pp.697.
- Lebinson & David, P. (2021). Profil Bakteri Penyebab Bakteremia dan Pola Kepekaan terhadap Antibiotik serta Faktor Risikonya di RSUP H. Adam Malik Medan Tahun 2020. Skripsi Sarjana. Universitas Sumatera Utara
- Lee, H., Kim, J., & Park, S. (2022). 'Neutrophil-to-Lymphocyte Ratio as a Predictor of Gram-Negative Bacterial Infection in Hospitalized Patients'. *BMC Infectious Diseases*, 22(1), pp. 567.
- Laukemann, S., *et al.* (2018). 'Early diagnosis of bloodstream infections and sepsis in adults: A clinical microbiology perspective'. *Clinical Microbiology and Infection*, 24(10), pp. 1121–1128.
- Li, X., Zhang, Y., Wang, J. *et al.* (2025). 'Diagnostic value of neutrophil-to-lymphocyte ratio combined with biomarkers for bloodstream infection in children'. *BMC pulmonary medicine*, 25(1), pp. 134.
- Liu, X., Shen, Y., Wang, H., Ge, Q., Fei, A. & Pan, S. (2016). 'Prognostic significance of neutrophil-to-lymphocyte ratio in patients with sepsis: a prospective observational study'. *Mediators of Inflammation*, 8191254.
- Lubis, B., Hasby, A.Y., Putra, A.O., Yanni, G.N. and Amelia, P. (2021). 'Hubungan Neutrophil–Lymphocyte Ratio (NLR) terhadap mortalitas pasien sepsis di Unit Perawatan Intensif RSUP Haji Adam Malik pada tahun 2018'. *Journal of Medicine*, 6(1), pp.1–5.
- Malik, C.P & Mustafa, M.I. (2017). Identifikasi mikroba metode pewarnaan Gram, Jurnal Praktikum Mikrobiologi Umum. Available at: <https://id.scribd.com/document/363076632/Jurnal-Pewarnaan-Gram> (Accessed: 02 Januari 2026)
- Martinez, J.M., Espírito Santo, A., Ramada, D., Fontes, F. & Medeiros, R. (2024) 'Diagnostic accuracy of neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio, and neutrophil–lymphocyte-to-platelet ratio biomarkers predicting bacteremia and sepsis in immunosuppressive patients with cancer: literature review'. *Porto Biomedical Journal*, 9(1), pp. e000254.
- Mudge, A.M., Banks, M.D., Barnett, A.G., Blackberry, I., Graves, N., Green, T., Harvey, G., Hubbard, R.E., Inouye, S.K., Kurrle, S., Lim, K., McRae, P., Peel, N.M., Suna, J. & Young, A.M. (2017). 'CHERISH (collaboration for hospitalised elders reducing the impact of stays in hospital): protocol for a multi-site improvement program to reduce geriatric syndromes in older inpatients'. *BMC Geriatrics*, 17(1), pp. e399.
- Nur, M.I., Wisudarti, C.F.R. & Rahardjo, S. (2023). 'Procalcitonin Sebagai Biomarker Diagnostik pada Pasien Sepsis yang Dirawat di ICU RSUP Dr. Sardjito'. *Jurnal Komplikasi Anestesi*, 10(3), pp. 20–28.
- Pamungkas, K. S., Suardana, I. W., & Adnyana, I. B. W. (2025). 'Identification of Gram-positive bacteria and their virulence characteristics from pig tonsil

- isolates based on catalase test and hemolytic properties'. *Buletin Veteriner Udayana*, 17(3), pp. 1002–1011.
- PKK, Prof.Dr. dr. A., M. Kes., n.d, 2016. *Ensiklopedia Penyakit*. Jakarta: PT Kanisius.
- Pramonodjati, F., Prabandari, A. S., Angelo, F., & Sudjono, E. (2019). 'Pengaruh Perokok Terhadap Adanya C-Reaktif Protein (Crp)'. *In Infokes*, 9(2), pp. 45-52.
- Prawira Ichsan, R. & Dwi Wahyu Widodo, A. (2023). 'Hospital-acquired Bacteremia: A Comprehensive Review on Risk Factors, Pathophysiology, Diagnosis and Management'. *International Journal of Research Publications*, 140(11), pp. e245978.
- Premandari, I.G.A.Ag.W.S., Abadi, M.F. & Arwidiana, D.P. (2023) 'Bakteri penyebab bakteremia dan pola resistensi terhadap antibiotik pada kultur darah', *Jurnal Insan Cendekia*, 10(3), pp. 189–200.
- Rahajeng, E.P. (2020) 'Analisis Laktat, Albumin dan Rasio Laktat Albumin Sebagai Prediktor Luaran Pada Pasien Sepsis dan Syok Septik di RSUP Dr. Wahidin Sudirohusodo Makassar', *Jurnal Kesehatan Andalas*, 9(1), pp. 26.
- Rhee, C., & Klompas, M. (2020). 'Sepsis trends, epidemiology, and definitions'. *Critical Care Clinics*, 36(1), pp. 1-19.
- Roldgaard, M., Benfield, T. and Tingsgård, S. (2024). 'Blood neutrophil to lymphocyte ratio is associated with 90-day mortality and 60-day readmission in Gram-negative bacteremia: a multi-center cohort study'. *BMC Infectious Diseases*, 24, pp.255.
- Rosadini, C. V., & Kagan, J. C. (2017). 'Bacterial lipopolysaccharide: Structure, function, and recognition'. *Cold Spring Harbor Perspectives in Biology*, 9(6), a015180.
- Rosales, C. (2020). 'Neutrophil roles in inflammation'. *Frontiers in Physiology*, 11, pp. 112.
- Santoso, M., Wijaya, R., & Pratama, R. (2020). 'Epidemiology of Gram-Negative Bacteremia in Hospitalized Patients: Trends and Risk Factors.' *Indonesian Journal of Clinical Microbiology*, 9(1), pp. 12–20.
- Saputra, I.M.Y., Gustawan, W., Utama, M.D. & Arhana, B. (2019). 'Rasio Neutrofil dan Limfosit (NLCR) Sebagai Faktor Risiko Terjadinya Infeksi Bakteri di Ruang Rawat Anak RSUP Sanglah Denpasar', *Sari Pediatri*, 20(6), pp. 354–359.
- Schuetz, P., Birkhahn, R., Sherwin, R., Jones, A. E., Singer, A., Kline, J. A., Beishuizen, A. (2017). 'Procalcitonin (PCT)-guided antibiotic stewardship: An international experts consensus on optimizing antibiotic use in acute respiratory infections and sepsis'. *BMC Medicine*, 15(1), pp. 1–16.
- Sforzini, L., Cattaneo, A., Ferrari, C., Turner, L., Mariani, N., Enache, D., Hastings, C., Lombardo, G., Nettis, M. A., Nikkheslat, N., Worrell, C., Zajkowska, Z., Kose, M., Cattane, N., Lopizzo, N., Mazzelli, M., Pointon, L., Cowen, P. J., Cavanagh, J., Pariante, C. M. (2023). 'Higher immune-related gene expression

- in major depression is independent of CRP'. *Molecular Psychiatry*, 28(3), pp. 1234–1245.
- Shen, X., Wang, Y., & Li, H. (2020). 'Immunosenescence and Susceptibility to Infection in the Elderly'. *Frontiers in Immunology*, 11, pp. 1682.
- Singer, M., Deutschman, C. S., Seymour, C. W., *et al.* (2016). 'The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3)' . *JAMA*, 315(8), pp. 801–810.
- Smith, J.P. & Doe, A.B. (2022) 'The Neutrophil-to-Lymphocyte Ratio as a New Prognostic Factor in Cancers: A Narrative Review'. *Journal of Oncologic Biomarkers*, 15(3), pp. 100-115.
- Soejono, C. H., *et al.* (2020). 'Length of stay of geriatric patients with gram-negative bacteremia: An analytical observation'. *Acta Medica Indonesiana*. 52(3), pp.245-251.
- Sogaard, K. K., Farkas, D. K., Pedersen, L., & Sorensen, H. T. (2022). 'Risk and prognosis of bloodstream infection in patients with cancer: A population-based cohort study'. *BMC Infectious Diseases*, 22, pp. 123.
- Sproston, N.R. & Ashworth, J.J. (2018) 'Role of C-reactive protein at sites of inflammation and infection', *Frontiers in Immunology*, 9, pp. 754.
- Sriwaningsi, L., Suryawan, I.W.B. and Suarca, I.K. (2021). 'Korelasi neutrofil-limfosit rasio dengan kadar C-reactive protein sebagai biomarker sepsis neonatus pada bayi prematur di Rumah Sakit Wangaya Denpasar'. *Medicina*, 52(3), pp.180–184.
- Sudiartha, I. P. G., Wiargitha, I. K., & Mahadewa, T. G. B. (2020). 'Perbedaan nilai Neutrophil Lymphocyte Ratio (NLR) terhadap pemeriksaan kultur darah dalam mendiagnosis sepsis pada pasien peritonitis di RSUP Sanglah, Bali, Indonesia'. *Intisari Sains Medis*, 11(1), pp. 165–171.
- Sun, L., Wang, Y., & Tang, H. (2023). 'Variability of NLR Across Different Patient Populations With Bacteremia'. *International Journal of Clinical Practice*, 77(4), e14852.
- Temkin E, Biran D, Braun T, Schwartz D, Carmeli Y. (2020). 'Analysis of Blood Culture Collection and Laboratory Processing Practices in Israel'. *JAMA Netw Open*, 3(10), e2023649.
- Wahdaniah & Tumpuk, S. (2018). 'Perbedaan penggunaan antikoagulan K2EDTA dan K3EDTA terhadap hasil pemeriksaan indeks eritrosit'. *Jurnal Laboratorium Khatulistiwa*, 5(2), pp. 45–52.
- Wahyuni, S., & Hasanah, F. (2023). 'Hubungan hs-CRP (High Sensitivity CReactive Protein) dengan APTT pada Diabetes mellitus tipe 2'. *Jurnal Sintesis: Penelitian Sains, Terapan Dan Analisisnya*, 4(1), pp 40–45.
- Weinstein, M. P., & Murphy, P. M. (2017). 'Clinical importance of blood cultures'. *Emerging Infectious Diseases*, 23(6), e161433.

- Wijaya, R., Santoso, M., & Pratama, R. (2021). 'Risk Factors for Gram-Negative Bacteremia in Hospitalized Patients: The Role of Inappropriate Antibiotic Use'. *Indonesian Journal of Clinical Pathology*, 10(2), pp. 45–52.
- WHO. (2020). *Health topics: Infectious diseases*. Available at: [https://www.who.int/topics/infectious\\_diseases/en/](https://www.who.int/topics/infectious_diseases/en/) (Accessed: 04 Januari 2026).
- Wong, M.M.Y., Lam, F. & Mok, C.C. (2024) 'The neutrophil-to-lymphocyte ratio as a surrogate marker for disease activity in systemic lupus erythematosus: a correlation study using the SLE-DAS', *Therapeutic Advances in Musculoskeletal Disease*, 17, 1759720X251347272.
- Zahorec, R. (2021). 'Neutrophil-to-lymphocyte ratio, past, present and future perspectives'. *Bratislava Medical Journal*, 122(7), pp. 474–488.
- Zahroh, N., Wicaksono, A.P. & Deharja, A. (2020). 'Analisis Manajemen Risiko Kesehatan dan Keselamatan Kerja di Bagian Filing RSUP dr. Soeradji Tirtonegoro Klaten', *J-REMI: Jurnal Rekam Medik dan Informasi Kesehatan*, 1(3), pp. 148–154.
- Zhang, G.-M. & Gu, Y.-Y. (2024). 'Diagnostic value of Procalcitonin, C-reactive protein-to-lymphocyte ratio (CLR), C-reactive protein and neutrophil-to-lymphocyte ratio (NLR) for predicting patients with Bacteraemia in the intensive care unit'. *Journal of Critical Care*, 81, pp. 154538.
- Zhang, L., Wang, Y., & Liu, H. (2021). 'Neutrophil-to-Lymphocyte Ratio as a Predictor of Severity and Mortality in Gram-Negative Sepsis: A Prospective Cohort Study'. *Frontiers in Immunology*, 12, 685321.
- Zhang, Z., Xie, P., Zhou, Y., & Zhang, Y. (2022). 'Prognostic value of neutrophil-to-lymphocyte ratio in patients with sepsis: A meta-analysis. *Inflammation*, 45, pp. 29–39.
- Zhou, T., Zheng, N., Li, X., Zhu, D. & Han, Y. (2021) 'Prognostic value of neutrophil-lymphocyte count ratio (NLCR) among adult ICU patients in comparison to APACHE II score and conventional inflammatory markers: A multi center retrospective cohort study', *BMC Emergency Medicine*, 21(1), pp. 418.
- Ziesmann, M.T. & Marshall, J.C. (2018). 'Multiple Organ Dysfunction: The Defining Syndrome of Sepsis'. *Surgical Infections*, 19(2), pp. 184–190.