



DAFTAR PUSTAKA



DAFTAR PUSTAKA

- Auffret, N., Leccia, M.-T., Ballanger, F., Claudel, J. P., Dahan, S., & Dréno, B. (2025). Acne-induced Post-inflammatory Hyperpigmentation: From Grading to Treatment. *Acta Dermato-Venereologica*, *105*, adv42925. <https://doi.org/10.2340/actadv.v105.42925>
- Bae, Y. C., Rettig, S., Weiss, E., Bernstein, L., & Geronemus, R. (2020). Treatment of Post-Inflammatory Hyperpigmentation in Patients With Darker Skin Types Using a Low Energy 1,927 nm Non-Ablative Fractional Laser: A Retrospective Photographic Review Analysis. *Lasers in Surgery and Medicine*, *52*(1), 7–12. <https://doi.org/10.1002/lsm.23173>
- Bolognia, J. L., Schaffer, J. V, & Cerroni, L. (2018). Vitiligo and other disorders of hypopigmentation. *Dermatology*. 4th Ed. Mosby/Elsevier, 1087–1114.
- Feng, W., & Yang, L. (2025). Low-Fluence Q-Switch 1064 Nm Laser Combined With Oral Tranexamic Acid: A Quicker Treatment for Laser-Induced Postinflammatory Hyperpigmentation. *Journal of Cosmetic Dermatology*, *24*(2). <https://doi.org/10.1111/jocd.70018>
- Huang, C.-Y., Lin, C.-H., Tsai, Y.-J., Lin, K.-C., Kuo, S. C.-H., Wong, L.-S., Lin, H.-P., & Hsieh, C.-H. (2024). Evaluating the efficacy of combining fractional CO2 laser and topical tranexamic acid for burn scar and hyperpigmentation treatment: A prospective, randomized, single-blinded, split-body study. *Dermatologica Sinica*, *42*(4), 265–272. <https://doi.org/10.4103/ds.DS-D-24-00104>
- Lai, D., Cheng, S., Zhou, S., Hao, J., Chen, H., Jia, K., Liu, H., & Cui, Y. (2024). 755-nm picosecond laser plus topical 20% azelaic acid compared to topical 20% azelaic acid alone for the treatment of melasma: a randomized, split-face and controlled trial. *Lasers in Medical Science*, *39*(1), 113. <https://doi.org/10.1007/s10103-024-04052-9>
- Lee, H.-J., Lee, S. H., Yoon, S. Y., Lee, J. W., Kim, Y. K., & Choi, E. H. (2018). Postinflammatory Hyperpigmentation Successfully Treated with 1,064-nm Picosecond-Domain Neodymium:Yttrium-Aluminum-Garnet Laser. *Medical Lasers*, *7*(1), 52–54. <https://doi.org/10.25289/ML.2018.7.1.52>
- Lindgren, A., Austin, A., & Welsh, K. (2021). The Use of Tranexamic Acid to Prevent and Treat Post-Inflammatory Hyperpigmentation. *Journal of Drugs in Dermatology*, *20*(3), 344–345. <https://doi.org/10.36849/JDD.5622>
- Nayak, C. S., Ansari, S. M. M., Salve, V., & Patil, S. (2019). Effectiveness of a combination of anti-pigmentary products in facial post-inflammatory hyperpigmentation. *International Journal of Research in Dermatology*, *6*(1), 1. <https://doi.org/10.18203/issn.2455-4529.IntJResDermatol20195199>

- Negishi, K., Kushikata, N., Tezuka, Y., Takeuchi, K., Miyamoto, E., & Wakamatsu, S. (2004). Study of the Incidence and Nature of “Very Subtle Epidermal Melasma” in Relation to Intense Pulsed Light Treatment. *Dermatologic Surgery*, *30*(6), 881–886. <https://doi.org/10.1111/j.1524-4725.2004.30257.x>
- Park, J.-H., Kim, J.-I., & Kim, W.-S. (2016). Treatment of Persistent Facial Postinflammatory Hyperpigmentation With Novel Pulse-in-Pulse Mode Intense Pulsed Light. *Dermatologic Surgery*, *42*(2), 218–224. <https://doi.org/10.1097/DSS.0000000000000627>
- Pulumati, A., Jaalouk, D., Algarin, Y. A., & Nouri, K. (2023). The role of 755-nm alexandrite picosecond laser in melasma management. *Archives of Dermatological Research*, *316*(1), 60. <https://doi.org/10.1007/s00403-023-02794-0>
- Ramírez-Oliveros, J. F., de Abreu, L., Tamler, C., Vilhena, P., & de Barros, M. H. (2020). Microneedling with drug delivery (hydroquinone 4% serum) as an adjuvant therapy for recalcitrant Melasma. *Skinmed*, *18*(1), 38–40. <https://pubmed.ncbi.nlm.nih.gov/32167455/>
- Ren, R., Bao, S., Qian, W., & Zhao, H. (2021). 755-nm Alexandrite Picosecond Laser with a Diffractive Lens Array or Zoom Handpiece for Post-Inflammatory Hyperpigmentation: Two Case Reports with a Three-Year Follow-Up. *Clinical, Cosmetic and Investigational Dermatology, Volume 14*, 1459–1464. <https://doi.org/10.2147/CCID.S323872>
- Ren, R., & Zhao, H. (2022). A case report on the use of 755-nm alexandrite picosecond laser in the treatment of post-inflammatory hyperpigmentation in the nose. *Journal of Cosmetic Dermatology*, *21*(5), 2046–2048. <https://doi.org/10.1111/jocd.14332>
- Roggenkamp, D., Dlova, N., Mann, T., Batzer, J., Riedel, J., Kausch, M., Zoric, I., & Kolbe, L. (2021). Effective reduction of post-inflammatory hyperpigmentation with the tyrosinase inhibitor isobutylamido-thiazoly-resorcinol (Thiamidol). *International Journal of Cosmetic Science*, *43*(3), 292–301. <https://doi.org/10.1111/ics.12694>
- Shucheng, H., Zhou, X., Du, D., Li, J., Yu, C., & Jiang, X. (2024). Effects of 15% Azelaic Acid Gel in the Management of Post-Inflammatory Erythema and Post-Inflammatory Hyperpigmentation in Acne Vulgaris. *Dermatology and Therapy*, *14*(5), 1293–1314. <https://doi.org/10.1007/s13555-024-01176-2>
- Sobhan, M., Talebi-Ghane, E., & Poostiyani, E. (2023). A comparative study of 20% azelaic acid cream versus 5% tranexamic acid solution for the treatment of postinflammatory hyperpigmentation in patients with acne vulgaris: A single-blinded randomized clinical trial. *Journal of Research in Medical Sciences*, *28*(1). https://doi.org/10.4103/jrms.jrms_443_22

- Tirthani, S., Srinivas, R. T., Kuppaswamy, S. K., Keloji, H., Venkatesh, V. B., & Chandrashekar, M. K. (2024). Comparison of Q-Switched Nd:YAG LASER with 10% Azelaic Acid versus Fractional CO₂ LASER with 10% Azelaic Acid in the treatment of Post Inflammatory Hyperpigmentation secondary to Acne. *South Eastern European Journal of Public Health*, 1742–1750. <https://doi.org/10.70135/seejph.vi.2660>
- Vashi, N. A., Wiryaa, S. A., Inyang, M., & Kundu, R. V. (2017). Facial Hyperpigmentation in Skin of Color: Special Considerations and Treatment. *American Journal of Clinical Dermatology*, 18(2), 215–230. <https://doi.org/10.1007/s40257-016-0239-8>
- Wang, J., Luo, H., Zhao, X., Wang, C., & Cheng, F. (2025). Clinical study on the effectiveness of laser combined with topical tranexamic acid in the treatment of melasma. *Lasers in Medical Science*, 40(1), 368. <https://doi.org/10.1007/s10103-025-04639-w>
- Wilson, M. J. V., Jones, I. T., Bolton, J., Larsen, L., & Fabi, S. G. (2018). The Safety and Efficacy of Treatment With a 1,927-nm Diode Laser With and Without Topical Hydroquinone for Facial Hyperpigmentation and Melasma in Darker Skin Types. *Dermatologic Surgery*, 44(10), 1304–1310. <https://doi.org/10.1097/DSS.0000000000001521>