

Phytochemical Screening and Evaluation of Red Ginger Extracts on Aphrodisiac Activity

by Ellies Tunjung Sm Dosen Fik

Submission date: 06-Feb-2023 11:20AM (UTC+0700)

Submission ID: 2007275230

File name: Extracts_on_Aphrodisiac_Activity-_internasional_Q4-_mei_2022.pdf (147.04K)

Word count: 3374

Character count: 18294

Phytochemical Screening and Evaluation of Red Ginger Extracts on Aphrodisiac Activity

DetECCIÓN Y EVALUACIÓN FITOQUÍMICA DE EXTRACTOS DE JENGIBRE ROJO SOBRE LA ACTIVIDAD AFRODISÍACA

Subhan Rullyansyah^{1a*}, Fuad Muzakky^{2a}, Karima Samlan^{3a}, Ria Hanistya^{4a},
Ellies Tunjung Sari Maulidiyanti^{5a}

SUMMARY

Introduction: One of the breeding processes is influenced by reproduction factors. Herbal therapy can relieve male infertility, such as red ginger. This study aimed to determine the effect of red ginger extract on aphrodisiac activity.

Methods: This study was designed to evaluate the effect of red ginger extract on aphrodisiac activity in Wistar rats. Twenty-five male Wistar rats and fifteen two-month-old females weighing 150-200 g were purchased and placed in cages at the Faculty of Health Sciences, University of Muhammadiyah Surabaya. Wistar rats were given light and dark cycles for 12 hours, fed pellet starter broiler, and drank water ad libitum. Mice were acclimatized for 14 days and randomly divided into five groups A, B, C, D, and E.

Groups B, C, and D were the treatment groups, while groups A and E were negative and positive controls, each with five rats. In each group and each kept singly in a separate cage. Groups A and E received 1 mL of distilled water and 5 mg of sildenafil citrate orally for the positive control group. Groups B, C, and D received red ginger extract suspension orally at doses of 100, 200, and 300 mg/kg, respectively, between 9:00 - and 10:00 daily for 21 days. Female mice were paired with male mice in a 1:1 ratio, and mating behaviour was recorded.

Results: Male rats in groups C and E each showed a significant increase ($p < 0.05$) in mounting frequency (MF), intromission frequency (IF), and ejaculatory frequency (EF).

Conclusion: Red ginger extract has activity as an aphrodisiac, but it is necessary to conduct tests related to toxicity tests.

Keywords: Aphrodisiac, mating, rat, red ginger, sex stimulant.

DOI: <https://doi.org/10.47307/GMC.2022.130.s1.26>

ORCID ID: 0000-0002-2701-6235¹

ORCID ID: 0000-0002-4805-7191²

ORCID ID: 0000-0002-8910-0678³

ORCID ID: 0000-0001-5759-5314⁴

ORCID ID: 0000-0001-9241-5104⁵

^aUniversitas Muhammadiyah Surabaya, Indonesia

*Corresponding Author: Subhan Rullyansyah
E-mail: subhanfarm.umsurabaya@gmail.com

Recibido: 1 de mayo 2022

Aceptado: 6 de mayo 2022

RESUMEN

Introducción: Uno de los procesos de crianza está influenciado por factores de reproducción. La terapia a base de hierbas puede aliviar la infertilidad masculina, como el jengibre rojo. Este estudio tuvo como objetivo determinar el efecto del extracto de jengibre rojo sobre la actividad afrodisíaca.

Métodos: Este estudio fue diseñado para evaluar el efecto del extracto de jengibre rojo sobre la actividad afrodisíaca en ratas Wistar. Se compraron veinticinco ratas Wistar macho y quince hembras de dos meses de edad que pesaban entre 150 y 200 g y se colocaron

en jaulas en la Facultad de Ciencias de la Salud de la Universidad de Muhammadiyah Surabaya. A las ratas Wistar se les dieron ciclos de luz y oscuridad durante 12 horas, se alimentaron con pollos de engorde iniciados con gránulos y bebieron agua a 6° bitum. Los ratones se aclimataron durante 14 días y se dividieron aleatoriamente en cinco grupos A, B, C, D y E. Los grupos B, C y D fueron los grupos de tratamiento, mientras que los grupos A y E fueron controles negativos y positivos, cada uno con cinco ratas. en cada grupo y cada uno mantenido individualmente en una jaula separada. Los grupos A y E recibieron 1 mL de agua destilada y 5 mg de citrato de sildenafil por vía oral para el grupo de control positivo. Los grupos B, C y D recibieron suspensión de extracto de jengibre rojo por vía oral en dosis de 100, 200 y 300 mg/kg, respectivamente, entre las 9:00 y las 10:00 diariamente durante 21 días. Los ratones hembra se emparejaron con ratones macho en una proporción de 1:1 y se registró el comportamiento.

Resultados: Las ratas macho de los grupos C y E mostraron un aumento significativo ($p < 0,05$) en la frecuencia de monta (MF), frecuencia de intromisión (IF) y frecuencia eyaculatoria (EF).

Conclusión: El extracto de jengibre rojo tiene actividad como afrodisíaco, pero es necesario realizar pruebas relacionadas con pruebas de toxicidad.

Palabras clave: Afrodisíaco, apareamiento, rata, jengibre rojo, estimulante sexual

INTRODUCTION

Reproduction is one of the most important factors influencing the breeding process, and its success is 10° ghly dependent on factors such as genetics, physical environment, nutrition, and management (1,2). Evidence from the literature suggests that nutritional factors are the most vital ingredients that directly 10° ffect reproductive performance and potentially moderate the effects of other factors (3-6). Herbal therapy can relieve 7° male infertility, regardless of the etiology (7). A large number of plants have been tested for their 7° ossible fertility regulating properties (8,9). Some medicinal plants are widely used as aphrodisiacs to relieve sexual dysfunction or fertility-enhancing agents (10). Stress interferes with reproductive 3° function, especially oxidative stress, due to the production of oxygen free radicals that exceed the antioxidant 3° capacity of stressed tissues (11). Each oxidizing radical is a potential agent of oxidative stress (11). The main

3° antioxidant enzymes in mammals are superoxide dismutase (SOD), catalase and glutathione peroxidase, all of which are endogenous (12). Substances such as vitamins A, C, flavonoids and carotenoids are examples of exogenous antioxidants found in food (13).

2° Red ginger is a well-known spice in Asia for its medicinal properties. The antioxidant and free radical scavenging properties of the ginger extract have been clinically proven (14,15), as well as anticancer (16,17), antibacterial (2° ,19), and to treat rheumatoid arthritis (20). The ginger 2° extract contains oleoresin and essential oils, yellow, oily, and pungent. 2° oleoresin contains phenolic compounds that play a role in antioxidant activity, such as gingerol, shogaol, and zingerone (21,22). Gingerol, of which 6-gingerol is the main constituent of oleoresin, is susceptible to heat, light, and oxygen. However, there is still a lack of information about red ginger extract in Wistar rats as an aphrodisiac 6° ve-enhancing agent. Therefore, it is necessary to determine 4° the effect of red ginger extract on the mounting frequency (MF), intromission frequency (IF) and ejaculation frequency (EF) in Wistar rats.

MATERIALS AND METHODS

Plant Material

Fresh plants were collected from Bilitro (Plant Research Center for Spices and Medicines). Red Ginger Rhizome washed with running water and cut into small pieces. It was air-dried for ten days, and the dried material was then ground into a powder and sifted using a 40 mesh sieve. The cold maceration method carried out Extraction with 96 % ethanol as solvent.

Making Red Ginger Extract

Red Ginger Simplicia powder was weighed as much as 100 g and then extracted with 96 % ethanol 1 000 mL using the maceration method for seven days. With occasional shaking and heating. At the end of the seventh day, a clear filtrate was obtained by filtering through a Buchner funnel. The filtrate was further concentrated using a rotary evaporator, transferred to a Petri dish, and dried

in an oven at 60 °C for five minutes. Finally, the extract was stored in a desiccator for 15 days to reduce humidity (23).

Phytochemical Qualitative Analysis

The Extract of Red Ginger was subjected to different chemical tests to detect phytoconstituents such as alkaloids, saponins, flavonoids, phlobatanins, reducing sugars, and anthranoids, cardiac glycosides, anthraquinones, and polyphenols were carried out according to the method adopted by Harborne (24-26).

Animal Testing

This study used male and female rats aged three months with a bodyweight of 150-200 g. They were obtained from the animal center laboratory of the Faculty of Pharmacy, Universitas Airlangga. Animal testing is designed according to ethical standards for animal use.

Libido Test

The libido test was carried out according to the method described by (27). Twenty-five male Wister rats weighing 150 - 200 g which exhibited faster sexual activity, were selected for this study. They were divided in five groups with five rats each, and each was kept singly in separate cages during the experimental period. Groups A and E were the negative and positive control groups, each of which the negative control received 1 mL/kg aquadest and the positive control 5 mg of sildenafil citrate orally. Groups B, C, and D received red ginger ethanol extract suspension orally at doses of 100, 200, and 300 mg/kg, respectively, daily for 21 days between 9:00 - 10:00 am. Male Wistar rats were not tested under unusual circumstances. The animals were brought to the laboratory from cages and exposed to low light between 16:00 and 18:00 every day for six days before the experiment. Fifteen female rats were made receptive by oral administration of ethyl estradiol suspension (Lynoral® tablets Organon Pharm) at a dose of 100 g/rat 48 hours before the experiment. Acceptance of female

animals was confirmed before testing by exposing them to male Wister rats. In addition to controls, the animals were familiarised with the test conditions. Each male Wister rat was placed individually in one cage with receptive female rats in the same cage. Receptive female mice were put into male cages with a ratio of 1 female to 1 male. Observation of mating behaviour started for 3 hours (28). The test will be stopped if the male fails to show sexual interest. If the female does not show a receptive attitude, it will be replaced by another female from the female Wistar rat cluster. Male Wister rats will be observed for mounting frequency (MF), intromission frequency (IF), and ejaculation frequency (EF).

Statistical Analysis

The data obtained were expressed as the mean \pm standard error mean (SEM) of the six animals in each group. Data from all groups were analysed using a one-way analysis of variance (ANOVA) followed by Dunnet's t-test.

RESULTS

A total of 100 g of the dried red Ginger powder yielded 28.8 g of the crude extract. The extracts of Red ginger were subjected to qualitative phytochemical screening for the detection of phytoconstituents like alkaloids, saponins, flavonoids, phlobatanins, reducing sugars, anthranoids, cardiac glycosides, anthraquinones, and polyphenols. The results revealed the presence of cardiac glycosides, alkaloids, saponin, flavonoids, and polyphenols (Table 1).

The results obtained for sexual behaviour between groups B, C, and D were treated with aqueous seed extract of *M. oleifera* at 100 mg, 200 mg, and 300 mg, respectively. Meanwhile, groups A and E were given distilled water and sildenafil citrate (Embragra-100®) at 1 mL/kg and 5 mg/kg, respectively, as shown in (Table 5). As a result, groups B, C, and D showed a significant increase in mounting frequency (MF), intromission frequency (IF), and ejaculatory frequency (EF) compared to control groups A and E, as presented in (Table 2).

PHYTOCHEMICAL SCREENING AND EVALUATION

Table 1

Phytochemical screening of extract Red Ginger	
Chemicals Constituents	Extract Red Ginger
Cardiac glycosides	+
Alkaloids	+
Saponins	+
Tannins	-
Flavonoids	+
Polyphenols	+
Reducing sugars	+
Phlobatanins	-
Anthraquinones	-
Anthranoids	-

Legend: + = Present; - = Absent

Table 2

Evaluation of Aphrodisiac Effects of Red Ginger on Wistar Rats			
Treatment Group	Mounting Frequency	Intromission Frequency	Ejaculation Frequency
A	1.0 ± 1.0a	0 ± 0a	0 ± 0a
B	4.6 ± 0.8b	3.2 ± 0.8b	2.2 ± 0.4b
C	9.0 ± 0.7b	6.8 ± 0.8b	3.4 ± 0.5b
D	18.8 ± 0.8b	13.0 ± 1.0b	7.8 ± 0.8b
E	13.2 ± 0.8b	11.4 ± 1.1b	7.0 ± 0.7b

Mean ± SE with different superscripts a, b within columns are significantly different at P < 0.05 compared to the negative control group.

DISCUSSION

It has been reported that steroids and saponins present in many plants have fertility potentiating properties and are useful in treating impotence that occurs due to an imbalance of reproductive hormones (29). Saponins found mainly in the leaves of *Tribulis Terrestris L.* have been used as aphrodisiac agents in rats (30). It is also known to stimulate testicular androgenesis and is important for testicular differentiation, integrity, and steroidogenic function steroidogenic (31,32) in goats and rams, respectively. This study also agrees with the findings reported by Mu¹¹allad et al. (2009), who studied the effect of *Nigella sativa* on spermatogenesis and fertility activity

of male albino rats (33).

This study indicates that the aqueous extract of Moringa seeds at 300 mg/kg can¹² increase the libido of male Wistar rats. This is in agreement with the findings reported by Wattanathorn et al., who reported the aphrodisiac effect induced by *Allium tuberosum* seed extract in male rats¹⁴ treated at a dose of 500 mg/kg for 21 days, which significantly increased the frequency of MF, frequency of IF intromission and frequency of EF ejaculation (34).

The significant increase in EF indicated that⁸ the extract and standard drug prolonged interest in intercourse, which is an indicator of increased sexual arousal as reported in rats Wattanathorn et al. Therefore, the increase in spermatogenic activity is a reflection of the pharmacologic¹³ properties of red ginger extract that may play a role in stimulating the secretion of testosterone which allows the availability of the hormone along the pituitary-gonadal axis. In addition to the intensity of orgasm and ejaculation, which is in accordance with the conclusions noted by Watcho et al. when working with *Mondia whitei* hexane extract on the reproductive performance of male rats. Similar findings were observed in rats given red ginger extract in this study (35).

CONCLUSION

Phytochemical screening of red⁴ ginger rhizome showed the presence of cardiac glycosides, alkaloids, saponins, flavonoids, polyphenols, and reducing sugars. As a rich source of phytochemicals and minerals, Zingiber officinale can be considered a potential traditional medicine source. Furthermore, the results of this study confirm that the consumption of red ginger extract has the potential to improve the libido quality of male rats. This also supports the herbalist's claim that red ginger has sexual arousal properties. Thus, red ginger extract proves to be an effective and safe alternative for increasing libido.

ACKNOWLEDGEMENT

I really appreciate the support from the Dean of the Faculty of Health Sciences, University of Muhammadiyah Surabaya. Personally Dr.

Mundakir, M. Kep., N.S. The same appreciation was also given to the Majelis Pendidikan Tinggi Penelitian dan Pengembangan Pimpinan Pusat Muhammadiyah. Thank you to all of you.

REFERENCES

1. Thornton PK. Livestock production: Recent trends, future prospects. *Philos Trans R Soc B Biol Sci.* 2010;365(1554):2853-2867.
2. Saifuddin AB, Affandi B, Djajadilaga, Widohariadi, Bimo, Ajello C. Strengthening Indonesia's reproductive health education and training system: Progress to date. *Med J Indones.* 1996;5(2):99-103.
3. Kheradmand A, Babaei H, Ali Batavani R. Effect of improved diet on semen quality and scrotal circumference in the ram. *Vet Arh.* 2006;76(4):333-341.
4. Diskin MG, Kenny DA. Optimising reproductive performance of beef cows and replacement heifers. *Animal.* 2014;8 Suppl 1:27-39.
5. Ibtisham F, Nawab A, Li G, Xiao M, An L, Naseer G. Effect of nutrition on reproductive efficiency of dairy animals. *Med Weter.* 2018;74(6):356-361.
6. Bennett LR, Wiweko B, Bell L, Shafira N, Pangestu M, Adayana IBP, et al. Reproductive knowledge and patient education needs among Indonesian women infertility patients attending three fertility clinics. *Patient Educ Couns.* 2015;98(3):364-369.
7. Ojekale AB, Lawal OA, Lasisi AK, Adeleke TI. Phytochemistry and spermatogenic potentials of aqueous extract of *Cissus populnea* (Guill. and Per) stem bark. *Sci World J.* 2006;6:2140-2146.
8. Bhatia DK, Sharma AK, Pathania PC, Khanduri NC. Antifertility effects of crude different of *Adiantum lunulatum* Burm. on Reproductive Organs of male albino rats. In: *Biological Forum-An International Journal.* Citeseer; 2010.p.88-93.
9. Sabdoningrum EK, Hidanah S, Chusniati S. Characterization and Phytochemical Screening of Meniran (*Phyllanthus niruri* Linn) Extract's Nanoparticles Used Ball Mill Method. *Pharmacogn J.* 2021;13(6s):1568-1572.
10. Fatoba T, Faleyimu O, Adebayo A. Effects of Increasing Aqueous Root Extract of *Moringa oleifera* on Sperm Production of Albino Rats. *Agrosearch.* 2013;13(1):29.
11. Afolabi AO, Aderoju HA, Alagbonsi IA. Effects of methanolic extract of *Moringa oleifera* leaves on semen and biochemical parameters in cryptorchid rats. *African J Tradit Complement Altern Med AJTCAM.* 2013;10(5):230-235.
12. Ighodaro O, Akinloye OA. First line defense antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defense grid. *Alexandria J Med.* 2018;54(4):287-293.
13. Siddiq A, Anwar F, Manzoor M, Fatima A. Antioxidant activity of different solvent extracts of *Moringa oleifera* leaves under accelerated storage of sunflower oil. *Asian J Plant Sci.* 2005;4(6):630-635.
14. Mošovská S, Nováková D, Kaliňák M. Antioxidant activity of ginger extract and identification of its active components. *Acta Chim Slovaca.* 2015;8(2):115-119.
15. Kusumawati N, Bahar A, Setiarso P, Muslim S, Auliya ARS. Ginger and temulawak based herbal tea as potential functional drink products in the era of covid-19. *Rasayan J Chem.* 2021;1920-1926.
16. Zhu Y, Warin RF, Soroka DN, Chen H, Sang S. Metabolites of Ginger Component [6]-Shogaol Remain Bioactive in Cancer Cells and Have Low Toxicity in Normal Cells: Chemical Synthesis and Biological Evaluation. *PLoS One.* 2013;8(1):e54677.
17. Aryanti PI, Haryanto J, Ulfiana E. Red Ginger (*Zingiber officinale* var. *rubrum*) massage reduces stiffness and functional Disability in Elderly with Osteoarthritis. *J Ners.* 2018;13(2):200-206.
18. Nikolić M, Vasić S, Đurđević J, Stefanović O, Čomić L. Antibacterial and anti-biofilm activity of ginger (*Zingiber officinale* (Roscoe)) ethanolic extract. *Kragujev J Sci.* 2014;(36):129-136.
19. Dewi C, Mahasri G, Subekti S. Anthelmintic effect of red ginger to control myxobolus-infected in Koi fish. *Indian Vet J.* 2018;95(7):43-44.
20. Al-Nahain A, Jahan R, Rahmatullah M. *Zingiber officinale*: A Potential Plant against Rheumatoid Arthritis. *Arthritis.* 2014;2014:159089.
21. Nwaoha M, Elizabeth I, Okafor GI, Apochi VO. Production of oleoresin from ginger (*Zingiber officinale*) peels and evaluation of its antimicrobial and antioxidative properties. *Afr J Microbiol Res.* 2013;7(42):4981-4989.
22. Anisa AK, Permata FS, Rantam FA, Ernawati R. Jejunal histomorphometry of salmonella enteritidis infected broiler chickens supplemented with red ginger and antibiotic. *Vet Pract.* 2020;21(2):474-478.
23. Mulia K, Risqi UY, Pane IF, Krisanti EA. Formulation, characterization, and release property of antioxidant supplement capsule with red ginger oleoresin extract-loaded chitosan microparticles. In: *J Physics: Conference Series.* IOP Publishing; 2019.p.62008.
24. Crews SK. Practical pharmacognosy. *The Analyst.* 1943;68:266-267.
25. Harborne JB. *Phytochemical Methods.* Phytochemical Methods. 1980.

PHYTOCHEMICAL SCREENING AND EVALUATION

26. Sofowora A. Research on medicinal plants and traditional medicine in Africa. *J Altern Complement Med.* 1996;2(3):365-372.
27. Akmar K, Noor MM. Effect of combination of *Gynura procumbens* aqueous extract and *Trigona* spp. honey on fertility and libido of streptozotocin-induced hyperglycaemic male rats. *Asian Pacific J Reprod.* 2019;8(2):56.
28. Kimchi NA, Broide E, Scapa E, Birkenfeld S. Antiplatelet therapy and the risk of bleeding induced by gastrointestinal endoscopic procedures. A systematic review of the literature and recommendations. *Digestion.* 2007;75(1):36-45.
29. Shukla YN, Khanuja SPS. Chemical, Pharmacological and Botanical Studies on *Petalium Murex*. *J Med Aromat Plant Sci.* 2004;1:64-69.
30. Goel B, Maurya NK. Aphrodisiac Herbal therapy for Erectile Dysfunction. *Arch Pharm Pract.* 2020;11(1):1-6.
31. Iliyasu D, Paulline C, Sankey R, Ate I, Joseph A, Olayinka O, et al. Effects of Experimental *Fasciola Gigantica* Infection on Serum Testosterone Profiles and their Relationship with Semen Characteristics of Yankasa Ram. *J Vet Adv.* 2015;5(5):938.
32. Salem MH, Kamel KI, Yousef MI, Hassan GA, El-Nouty FD. Protective role of ascorbic acid to enhance semen quality of rabbits treated with sublethal doses of aflatoxin B1. *Toxicology.* 2001;162(3):209-218.
33. Mohammad MA, Mohamad MM, Dradka H. Effects of black seeds (*Nigella sativa*) on spermatogenesis and fertility of male albino rats. *Res J Med Med Sci.* 2009;4(2):386-390.
34. Wattanathorn J, Pangpookiew P, Muchimapura S, Sripanidkulchai K, Sripanidkulchai B. Aphrodisiac activity of *Kaempferia parviflora*. *Am J Agric Biol Sci.* 2012; 7(2):114-120.
35. Watcho P, Donfack MM, Zelefack F, Nguelafack TB, Wansi S, Nguola F, et al. Effects of the hexane extract of *Mondia whitei* on the reproductive organs of male rat. *African J Tradit Complement Altern Med.* 2005;2(3):302-311.

Phytochemical Screening and Evaluation of Red Ginger Extracts on Aphrodisiac Activity

ORIGINALITY REPORT

18%

SIMILARITY INDEX

%

INTERNET SOURCES

18%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Tajuddin, Shamshad Ahmad, Abdul Latif, Iqbal Ahmad Qasmi, Kunwar Mohammad Yusuf Amin. "An experimental study of sexual function improving effect of *Myristica fragrans* Houtt. (nutmeg)", *BMC Complementary and Alternative Medicine*, 2005
Publication 3%
- 2 K Mulia, U Y Risqi, I F Pane, E A Krisanti. "Formulation, Characterization, and Release Property of Antioxidant Supplement Capsule with Red Ginger Oleoresin Extract-loaded Chitosan Microparticles", *Journal of Physics: Conference Series*, 2019
Publication 3%
- 3 Afolabi, AO, HA Aderoju, and IA Alagbonsi. "Effects of methanolic extract of *Moringa Oleifera* leave on semen and biochemical parameters in cryptorchid rats", *African Journal of Traditional Complementary and Alternative Medicines*, 2013. 2%

4

Yakubu Musa Toyin, Ajiboye Taofeek Olakunle, Akanji Musbau Adewunmi. "Toxicity and Beneficial Effects of Some African Plants on the Reproductive System", Elsevier BV, 2014

Publication

2%

5

Tajuddin, Shamshad Ahmad, Abdul Latif, Iqbal Ahmad Qasmi. "Effect of 50% ethanolic extract of *Syzygium aromaticum* (L.) Merr. & Perry. (clove) on sexual behaviour of normal male rats", *BMC Complementary and Alternative Medicine*, 2004

Publication

1%

6

M Simarmata, B Halim, D Ardinata. "Effects of red ginger capsule supplementin reducing PGF2 α concentrations and pain intensity in primary dysmenorrhea", *IOP Conference Series: Earth and Environmental Science*, 2018

Publication

1%

7

Jimoh Olatunji Abubakar, Oyeyemi Wahab Adekunle, Okin-Aminu Hafsat Ololade, Oyeyemi Bolaji Fatai. "Reproductive characteristics, semen quality, seminal oxidative status, steroid hormones, sperm production efficiency of rabbits fed herbal supplements", *Theriogenology*, 2021

Publication

1%

8

Submitted to University of Limpopo

Student Paper

1 %

9

Andrés Cáceres Barreno, Carlos Cava Vergiú, José Robello Malatto, Erika Alberca Ramos, Jaime Rodríguez Chessa. "Biocompatibilidad del cianoacrilato de butilo en suturas en piel en comparación con las suturas convencionales", Revista Odontológica Mexicana, 2013

Publication

1 %

10

Anna Wilkanowska, Dariusz Kokoszyński. "Effect of Diet and Physical Activity of Farm Animals on their Health and Reproductive Performance", Elsevier BV, 2015

Publication

1 %

11

Saeed Al-Zahrani. "Thymoquinone and vitamin E supplementation improve the reproductive characteristics of heat stressed male mice", Journal of Medicinal Plants Research, 2012

Publication

1 %

12

Sonmez, M.. "The effect of ascorbic acid supplementation on sperm quality, lipid peroxidation and testosterone levels of male Wistar rats", Theriogenology, 20050415

Publication

1 %